

Nonlinear Gyrokinetics:

**A powerful tool for the description of
microturbulence in magnetized plasmas^a**

John A. Krommes
Princeton University

Presented at
“Turbulent Mixing and Beyond”
ICTP, Trieste, Italy
July 29, 2009

^aWork supported by U.S. Dept. of Energy Contract No. DE-AC02-09CH11466.

Table of Contents

- **Introduction: Heuristic Gyrokinetics**
- **Systematic Gyrokinetics, Part I: The Gyrokinetic Equations of Motion**
- **Systematic Gyrokinetics, Part II: The Gyrokinetic Poisson Equation**
- **The Current Status of Gyrokinetics**

Introduction: Heuristic Gyrokinetics

“Canonical” equations for turbulence

 Fluids — The Navier–Stokes equation (for $\vec{\nabla} \cdot \vec{u} = 0$):

$$\partial_t \vec{u}(\vec{x}, t) + \vec{u} \cdot \vec{\nabla} \vec{u} = -\vec{\nabla} P + \mu \nabla^2 \vec{u} \quad (1)$$

(3 dimensions + time).

“Canonical” equations for turbulence


- Fluids — The **Navier–Stokes equation** (for $\vec{\nabla} \cdot \vec{u} = 0$):

$$\partial_t \vec{u}(\vec{x}, t) + \vec{u} \cdot \vec{\nabla} \vec{u} = -\vec{\nabla} P + \mu \nabla^2 \vec{u} \quad (1)$$

(3 dimensions + time).

- Plasmas: The **Vlasov equation** (+ collision operator):

$$\frac{\partial f(\vec{x}, \vec{v}, t)}{\partial t} + \vec{v} \cdot \vec{\nabla} f + \frac{q}{m} (\vec{E} + \underbrace{c^{-1} \vec{v} \times \vec{B}}_{\text{rapid gyration}}) \cdot \frac{\partial f}{\partial \vec{v}} = -C[f] \quad (2)$$



[6 dimensions (!!!) + time]. A generalization of **Boltzmann’s eq’n**.

“Canonical” equations for turbulence


- Fluids — The **Navier–Stokes equation** (for $\vec{\nabla} \cdot \vec{u} = 0$):

$$\partial_t \vec{u}(\vec{x}, t) + \vec{u} \cdot \vec{\nabla} \vec{u} = -\vec{\nabla} P + \mu \nabla^2 \vec{u} \quad (1)$$

(3 dimensions + time).

- Plasmas: The **Vlasov equation** (+ collision operator):

$$\frac{\partial f(\vec{x}, \vec{v}, t)}{\partial t} + \vec{v} \cdot \vec{\nabla} f + \frac{q}{m} (\vec{E} + \underbrace{c^{-1} \vec{v} \times \vec{B}}_{\text{rapid gyration}}) \cdot \frac{\partial f}{\partial \vec{v}} = -C[f] \quad (2)$$



[6 dimensions (!!!) + time]. A generalization of **Boltzmann’s eq’n**.

- kinetic* — details of f in velocity space may be important (for linear growth rates and kinetic dissipation).
- electromagnetic* — self-consistent $\vec{E}[f]$ and $\vec{B}[f]$.

“Canonical” equations for turbulence


- Fluids — The **Navier–Stokes equation** (for $\vec{\nabla} \cdot \vec{u} = 0$):

$$\partial_t \vec{u}(\vec{x}, t) + \vec{u} \cdot \vec{\nabla} \vec{u} = -\vec{\nabla} P + \mu \nabla^2 \vec{u} \quad (1)$$

(3 dimensions + time).

- Plasmas: The **Vlasov equation** (+ collision operator):

$$\frac{\partial f(\vec{x}, \vec{v}, t)}{\partial t} + \vec{v} \cdot \vec{\nabla} f + \frac{q}{m} (\vec{E} + \underbrace{c^{-1} \vec{v} \times \vec{B}}_{\text{rapid gyration}}) \cdot \frac{\partial f}{\partial \vec{v}} = -C[f] \quad (2)$$




[6 dimensions (!!!) + time]. A generalization of **Boltzmann’s eq’n**.

- kinetic* — details of f in velocity space may be important (for linear growth rates and kinetic dissipation).
- electromagnetic* — self-consistent $\vec{E}[f]$ and $\vec{B}[f]$.

Plasmas motions span greatly disparate length and time scales.

Consider the ITER device...



- Cost $\sim E 10^{10}$.
- $B_T = 5 \text{ Tesla} = 5 \times 10^4 \text{ Gauss}$
- a (minor radius) $> 2 \text{ meters}$
 \gg gyroradius ρ_i
- Pulse length $> 1000 \text{ sec}$
 \gg turbulence autocorrelation time.

Fig. 5. The ITER device being constructed in Cadarache, France.

The **disparate time and space scales** in this device present a very difficult challenge for numerical simulation.

In a strong magnetic field, the large gyrofrequency complicates direct numerical solution of the plasma kinetic equation.

The gyrofrequency for species s ($s = e$ or i) is

$$\omega_{cs} = (qB/mc)_s. \quad (3)$$

Typical gyroperiods:

$$2\pi/\omega_{ce} < 10^{-10} \text{ sec}; \quad 2\pi/\omega_{ci} < 10^{-7} \text{ sec} = 0.1 \mu\text{sec}. \quad (4)$$


In a strong magnetic field, the large gyrofrequency complicates direct numerical solution of the plasma kinetic equation.

The gyrofrequency for species s ($s = e$ or i) is

$$\omega_{cs} = (qB/mc)_s. \quad (3)$$

Typical gyroperiods:

$$2\pi/\omega_{ce} < 10^{-10} \text{ sec}; \quad 2\pi/\omega_{ci} < 10^{-7} \text{ sec} = 0.1 \mu\text{sec}. \quad (4)$$

 Even with modern supercomputers, it is still not possible to follow all details (6D + time) of an ensemble of gyrating particles in ITER-scale devices for interesting lengths of time.

In a strong magnetic field, the large gyrofrequency complicates direct numerical solution of the plasma kinetic equation.

The gyrofrequency for species s ($s = e$ or i) is

$$\omega_{cs} = (qB/mc)_s. \quad (3)$$

Typical gyroperiods:

$$2\pi/\omega_{ce} < 10^{-10} \text{ sec}; \quad 2\pi/\omega_{ci} < 10^{-7} \text{ sec} = 0.1 \mu\text{sec}. \quad (4)$$

- Even with modern supercomputers, it is still not possible to follow all details (6D + time) of an ensemble of gyrating particles in ITER-scale devices for interesting lengths of time.
- Fortunately, for microturbulence in magnetically confined fusion devices, the collective modes of interest (e.g., *drift waves*) are **low frequency**:

$$\frac{\omega}{\omega_{ci}} \lesssim \frac{\rho_i}{L_n} \sim \frac{0.2 \text{ cm}}{2 \times 10^2 \text{ cm}} \approx 10^{-3} \ll 1. \quad (5)$$

In a strong magnetic field, the large gyrofrequency complicates direct numerical solution of the plasma kinetic equation.

The gyrofrequency for species s ($s = e$ or i) is

$$\omega_{cs} = (qB/mc)_s. \quad (3)$$

Typical gyroperiods:


$$2\pi/\omega_{ce} < 10^{-10} \text{ sec}; \quad 2\pi/\omega_{ci} < 10^{-7} \text{ sec} = 0.1 \mu\text{sec}. \quad (4)$$

- Even with modern supercomputers, it is still not possible to follow all details (6D + time) of an ensemble of gyrating particles in ITER-scale devices for interesting lengths of time.
- Fortunately, for microturbulence in magnetically confined fusion devices, the collective modes of interest (e.g., *drift waves*) are **low frequency**:

$$\frac{\omega}{\omega_{ci}} \lesssim \frac{\rho_i}{L_n} \sim \frac{0.2 \text{ cm}}{2 \times 10^2 \text{ cm}} \approx 10^{-3} \ll 1. \quad (5)$$

- Therefore, one can make some analytical reductions by averaging the motion over the rapid gyration (gyrophase ζ).

Gyrokinetics is the central formalism that describes the low-frequency physics of magnetized plasma.



For $\omega/\omega_{ci} \ll 1$ and $k_{\perp}\rho_i \ll 1$,
 particles move on average via the **drift equations**.

$$\frac{d\vec{X}}{dt} = v_{\parallel} \hat{\mathbf{b}} + \vec{V}_E + \vec{V}_d, \quad (6a)$$

$$\frac{dv_{\parallel}}{dt} = \frac{q}{m} E_{\parallel} + \dots, \quad (6b)$$

$$\frac{d\mu}{dt} = 0 \quad (\text{magnetic moment is conserved!}). \quad (6c)$$

Here the **guiding-center drifts** are

$$\vec{V}_E \doteq c \vec{E} \times \hat{\mathbf{b}} / B \quad (\vec{E} \times \vec{B} \text{ drift}), \quad (7a)$$

$$\vec{V}_d \doteq \frac{v_{\perp}^2/2}{\omega_c} \hat{\mathbf{b}} \times \vec{\nabla} \ln B + \frac{v_{\parallel}^2}{\omega_c} \hat{\mathbf{b}} \times (\hat{\mathbf{b}} \cdot \vec{\nabla} \hat{\mathbf{b}}) \quad (\text{magnetic drifts}), \quad (7b)$$

and the **magnetic moment**^a is $\mu \approx \frac{1}{2} m v_{\perp}^2 / \omega_c(\vec{x})$. Note

$$\mu \propto \oint \vec{p} \cdot d\vec{q}$$

(special case of general theory of “adiabatic invariants”).

^aThis formula for μ is the lowest-order approximation to the true magnetic moment $\bar{\mu}$. See later discussion.

For a distribution of particles and $k_{\perp}\rho_i \ll 1$,
we may write the **drift kinetic equation**:

$$\frac{\partial F(\vec{X}, v_{\parallel}, \mu, t)}{\partial t} + \underbrace{v_{\parallel} \nabla_{\parallel} F}_{\text{parallel streaming}} + \underbrace{(\vec{V}_E + \vec{V}_d) \cdot \vec{\nabla} F}_{\text{guiding-center drifts}} + \underbrace{\frac{q}{m}(E_{\parallel} + \dots) \frac{\partial F}{\partial v_{\parallel}}}_{\text{parallel acceleration}} = 0.$$

(8)

For a distribution of particles and $k_{\perp}\rho_i \ll 1$,
we may write the **drift kinetic equation**:

$$\frac{\partial F(\vec{X}, v_{\parallel}, \mu, t)}{\partial t} + \underbrace{v_{\parallel} \nabla_{\parallel} F}_{\text{parallel streaming}} + \underbrace{(\vec{V}_E + \vec{V}_d) \cdot \vec{\nabla} F}_{\text{guiding-center drifts}} + \underbrace{\frac{q}{m}(E_{\parallel} + \dots) \frac{\partial F}{\partial v_{\parallel}}}_{\text{parallel acceleration}} = 0. \quad (8)$$

- No $\partial/\partial\zeta$; thus **5D + time** (saved one dimension and removed high-frequency motion).

For a distribution of particles and $k_{\perp}\rho_i \ll 1$,
we may write the **drift kinetic equation**:

$$\frac{\partial F(\vec{X}, v_{\parallel}, \mu, t)}{\partial t} + \underbrace{v_{\parallel} \nabla_{\parallel} F}_{\text{parallel streaming}} + \underbrace{(\vec{V}_E + \vec{V}_d) \cdot \vec{\nabla} F}_{\text{guiding-center drifts}} + \underbrace{\frac{q}{m}(E_{\parallel} + \dots) \frac{\partial F}{\partial v_{\parallel}}}_{\text{parallel acceleration}} = 0. \quad (8)$$

- No $\partial/\partial\zeta$; thus **5D + time** (saved one dimension and removed high-frequency motion).
- No $\partial/\partial\mu$; μ is conserved and enters only as a parameter.

For a distribution of particles and $k_{\perp}\rho_i \ll 1$,
we may write the **drift kinetic equation**:

$$\frac{\partial F(\vec{X}, v_{\parallel}, \mu, t)}{\partial t} + \underbrace{v_{\parallel} \nabla_{\parallel} F}_{\text{parallel streaming}} + \underbrace{(\vec{V}_E + \vec{V}_d) \cdot \vec{\nabla} F}_{\text{guiding-center drifts}} + \underbrace{\frac{q}{m}(E_{\parallel} + \dots) \frac{\partial F}{\partial v_{\parallel}}}_{\text{parallel acceleration}} = 0. \quad (8)$$

● No $\partial/\partial\zeta$; thus **5D + time** (saved one dimension and removed high-frequency motion).

● No $\partial/\partial\mu$; μ is conserved and enters only as a parameter.


● For $k_{\perp}\rho \ll 1$, the particle position \vec{x} and the guiding-center position \vec{X} are essentially coincident.

For a distribution of particles and $k_{\perp}\rho_i \ll 1$,
we may write the **drift kinetic equation**:

$$\frac{\partial F(\vec{X}, v_{\parallel}, \mu, t)}{\partial t} + \underbrace{v_{\parallel} \nabla_{\parallel} F}_{\text{parallel streaming}} + \underbrace{(\vec{V}_E + \vec{V}_d) \cdot \vec{\nabla} F}_{\text{guiding-center drifts}} + \underbrace{\frac{q}{m}(E_{\parallel} + \dots) \frac{\partial F}{\partial v_{\parallel}}}_{\text{parallel acceleration}} = 0. \quad (8)$$

- No $\partial/\partial\zeta$; thus **5D + time** (saved one dimension and removed high-frequency motion).
- No $\partial/\partial\mu$; μ is conserved and enters only as a parameter.

- For $k_{\perp}\rho \ll 1$, the particle position \vec{x} and the guiding-center position \vec{X} are essentially coincident.
- But for $k_{\perp}\rho = O(1)$, one must introduce the notion of the **gyrocenter**, the *average* position of the particle. Gyrocenters feel *effective* potentials.



Gyrocenters feel **effective** electromagnetic fields. They move according to the **gyrokinetic equation**.

Let $\vec{E} = -\vec{\nabla}\varphi$ (electrostatics). Since $\varphi(\vec{x}) = (2\pi)^{-3} \int d\vec{k} \varphi_{\vec{k}} e^{i\vec{k}\cdot\vec{x}}$,

$$\langle \varphi(\vec{x}) \rangle_{\zeta} \equiv \langle \varphi \rangle(\vec{X}) = \frac{1}{2\pi} \int_0^{2\pi} d\zeta \int \frac{d\vec{k}}{(2\pi)^3} \varphi_{\vec{k}} e^{i\vec{k}\cdot[\vec{X} + \vec{\rho}(\zeta)]} \quad (9a)$$

$$= \int \frac{d\vec{k}}{(2\pi)^3} e^{i\vec{k}\cdot\vec{X}} \underbrace{J_0(k_{\perp}\rho) \varphi_{\vec{k}}}_{\langle \varphi \rangle_{\vec{k}}} \quad (9b)$$

If we appropriately introduce the effective fields, we can write the **gyrokinetic equation** immediately:

Gyrocenters feel **effective** electromagnetic fields. They move according to the **gyrokinetic equation**.

Let $\vec{E} = -\vec{\nabla}\varphi$ (electrostatics). Since $\varphi(\vec{x}) = (2\pi)^{-3} \int d\vec{k} \varphi_{\vec{k}} e^{i\vec{k}\cdot\vec{x}}$,

$$\langle \varphi(\vec{x}) \rangle_{\zeta} \equiv \langle \varphi \rangle(\vec{X}) = \frac{1}{2\pi} \int_0^{2\pi} d\zeta \int \frac{d\vec{k}}{(2\pi)^3} \varphi_{\vec{k}} e^{i\vec{k}\cdot[\vec{X} + \vec{\rho}(\zeta)]} \quad (9a)$$

$$= \int \frac{d\vec{k}}{(2\pi)^3} e^{i\vec{k}\cdot\vec{X}} \underbrace{J_0(k_{\perp}\rho)}_{\langle \varphi \rangle_{\vec{k}}} \varphi_{\vec{k}}. \quad (9b)$$

If we appropriately introduce the effective fields, we can write the **gyrokinetic equation** immediately:

$$\frac{\partial F(\vec{X}, v_{\parallel}, \mu, t)}{\partial t} + \underbrace{v_{\parallel} \nabla_{\parallel} F}_{\text{parallel streaming}} + \underbrace{(\langle \vec{V}_{\vec{E}} \rangle + \vec{V}_d) \cdot \vec{\nabla} F}_{\text{gyrocenter drifts}} + \underbrace{\frac{q}{m} (\langle \vec{E}_{\parallel} \rangle + \dots)}_{\text{parallel acceleration}} \frac{\partial F}{\partial v_{\parallel}} = 0. \quad (10)$$

The gyrokinetic equation is the workhorse of research on modern fusion microturbulence.

**In the GK formalism, the magnetic moment μ is assumed to be conserved.
It is just a parameter in the GKE.**

The structure of the GKE is

$$\frac{\partial F(\vec{X}, v_{\parallel}, \mu, t)}{\partial t} + \dot{\vec{X}} \cdot \vec{\nabla} F + v_{\parallel} \frac{\partial F}{\partial v_{\parallel}} + \underbrace{\dot{\mu}}_{0!} \frac{\partial F}{\partial \mu} + \underbrace{\dot{\zeta}}_{0!} \frac{\partial F}{\partial \zeta} = 0. \quad (11)$$

The coefficients of the gradients are the characteristic equations of motion:

- $\hat{\mathbf{b}} \cdot \dot{\vec{X}}$ — parallel streaming
- $\dot{\vec{X}}_{\perp}$ — perpendicular drifts
- \dot{v}_{\parallel} — parallel acceleration
- $\dot{\mu} = 0$ — μ is conserved.

To close the gyrokinetic equation, one needs the fields.
Thus we must consider the **GK Maxwell equations**.

For definiteness, consider electrostatics. Then one has $\vec{E} = -\vec{\nabla}\varphi$
and one must solve Poisson's equation

$$-\nabla^2\varphi(\vec{x}, t) = 4\pi \underbrace{\rho(\vec{x}, t)}_{\text{charge density}} . \quad (12)$$

To close the gyrokinetic equation, one needs the fields.
Thus we must consider the **GK Maxwell equations**.

For definiteness, consider electrostatics. Then one has $\vec{E} = -\vec{\nabla}\varphi$
and one must solve Poisson's equation

$$-\nabla^2\varphi(\vec{x}, t) = 4\pi \underbrace{\rho(\vec{x}, t)}_{\text{charge density}} . \quad (12)$$

Subtlety arises because:

- 🔴 In gyrokinetics, the particle position \vec{x} and the gyrocenter position \vec{X} are distinct.

To close the gyrokinetic equation, one needs the fields.
Thus we must consider the **GK Maxwell equations**.

For definiteness, consider electrostatics. Then one has $\vec{E} = -\vec{\nabla}\varphi$
and one must solve Poisson's equation

$$-\nabla^2\varphi(\vec{x}, t) = 4\pi \underbrace{\rho(\vec{x}, t)}_{\text{charge density}} . \quad (12)$$

Subtlety arises because:

- In gyrokinetics, the particle position \vec{x} and the gyrocenter position \vec{X} are distinct.
- One needs the particle charge density $\rho(\vec{x})$, but is given the *gyrocenter* (phase-space) density $F(\vec{Z})$.

To close the gyrokinetic equation, one needs the fields.
 Thus we must consider the **GK Maxwell equations**.

For definiteness, consider electrostatics. Then one has $\vec{E} = -\vec{\nabla}\varphi$ and one must solve Poisson's equation

$$-\nabla^2\varphi(\vec{x}, t) = 4\pi \underbrace{\rho(\vec{x}, t)}_{\text{charge density}}. \quad (12)$$

Subtlety arises because:

- In gyrokinetics, the particle position \vec{x} and the gyrocenter position \vec{X} are distinct.
- One needs the particle charge density $\rho(\vec{x})$, but is given the *gyrocenter* (phase-space) density $F(\vec{Z})$.
- $F(\vec{Z}) \xRightarrow{T} \rho(\vec{x}) \xRightarrow{\text{Poisson}} \varphi(\vec{x}) \xRightarrow{\vec{\nabla}} \vec{E}(\vec{x}) \xRightarrow{T^{-1}} \vec{E}(\vec{X}) \xRightarrow{\text{GKE}} F(\vec{Z})$.

The most important physical distinction between particles and gyrocenters lies in the **polarization drift** (of the ions).

Iterative solution of $m \frac{d\vec{u}}{dt} = q(\vec{E} + c^{-1}\vec{u} \times \vec{B}) + \dots$ for $B = \text{const}$ leads to $\vec{u} = \vec{u}_{\vec{E}} + \vec{u}^{\text{pol}} + \dots$, where the **polarization drift velocity** is

$$\vec{u}^{\text{pol}} \doteq \frac{1}{\omega_c} \frac{\partial}{\partial t} \left(\frac{c\vec{E}_{\perp}}{B} \right). \quad (13)$$

The polarization drift leads to a **polarization charge density** ρ^{pol} :

$$\partial_t \rho^{\text{pol}} = -\vec{\nabla} \cdot (nq\vec{u}^{\text{pol}}). \quad (14)$$




Fig. 7. $\vec{E} = E(t)\hat{y}$,
 $\vec{B} = B_0\hat{z}$. $\vec{V}_E \propto \hat{x}$,
 $\vec{V}^{\text{pol}} \propto \hat{y}$.

The most important physical distinction between particles and gyrocenters lies in the **polarization drift** (of the ions).

Iterative solution of $m \frac{d\vec{u}}{dt} = q(\vec{E} + c^{-1}\vec{u} \times \vec{B}) + \dots$ for $B = \text{const}$ leads to $\vec{u} = \vec{u}_{\vec{E}} + \vec{u}^{\text{pol}} + \dots$, where the **polarization drift velocity** is

$$\vec{u}^{\text{pol}} \doteq \frac{1}{\omega_c} \frac{\partial}{\partial t} \left(\frac{c\vec{E}_{\perp}}{B} \right). \quad (13)$$

The polarization drift leads to a **polarization charge density** ρ^{pol} :

$$\partial_t \rho^{\text{pol}} = -\vec{\nabla} \cdot (nq\vec{u}^{\text{pol}}). \quad (14)$$




Fig. 7. $\vec{E} = E(t)\hat{y}$,
 $\vec{B} = B_0\hat{z}$. $\vec{V}_E \propto \hat{x}$,
 $\vec{V}^{\text{pol}} \propto \hat{y}$.

**Gyrocenters move with $\vec{E} \times \vec{B}$;
 particles polarize.**

**The gyrokinetic equation contains no polarization drift!
 Polarization is accounted for in the GK Poisson equation.**

The most important physical distinction between particles and gyrocenters lies in the **polarization drift** (of the ions).

Iterative solution of $m \frac{d\vec{u}}{dt} = q(\vec{E} + c^{-1} \vec{u} \times \vec{B}) + \dots$ for $B = \text{const}$ leads to $\vec{u} = \vec{u}_{\vec{E}} + \vec{u}^{\text{pol}} + \dots$, where the **polarization drift velocity** is

$$\vec{u}^{\text{pol}} \doteq \frac{1}{\omega_c} \frac{\partial}{\partial t} \left(\frac{c \vec{E}_{\perp}}{B} \right). \quad (13)$$

The polarization drift leads to a **polarization charge density** ρ^{pol} :

$$\partial_t \rho^{\text{pol}} = -\vec{\nabla} \cdot (nq\vec{u}^{\text{pol}}). \quad (14)$$




Fig. 7. $\vec{E} = E(t)\hat{y}$,
 $\vec{B} = B_0\hat{z}$. $\vec{V}_E \propto \hat{x}$,
 $\vec{V}^{\text{pol}} \propto \hat{y}$.

**Gyrocenters move with $\vec{E} \times \vec{B}$;
 particles polarize.**

**The gyrokinetic equation contains no polarization drift!
 Polarization is accounted for in the GK Poisson equation.**

Ion polarization leads to an interesting form of the GK Poisson equation.

One finds (for $T_i = 0$) $\rho_i^{\text{pol}} = (nq)_i \rho_s^2 \nabla_{\perp}^2 \Phi$, where $\Phi \doteq e\varphi/T_e$ and

$$\rho_s \doteq c_s/\omega_{ci} \quad (\text{“sound radius”}; \rho_s = \rho_i \text{ for } T_i = T_e), \quad (15a)$$

$$c_s \doteq (T_e/m_i)^{1/2} \quad (\text{sound speed}). \quad (15b)$$

Ion polarization leads to an interesting form of the GK Poisson equation.

One finds (for $T_i = 0$) $\rho_i^{\text{pol}} = (nq)_i \rho_s^2 \nabla_{\perp}^2 \Phi$, where $\Phi \doteq e\varphi/T_e$ and

$$\rho_s \doteq c_s/\omega_{ci} \quad (\text{"sound radius"; } \rho_s = \rho_i \text{ for } T_i = T_e), \quad (15a)$$

$$c_s \doteq (T_e/m_i)^{1/2} \quad (\text{sound speed}). \quad (15b)$$

Then Poisson's equation becomes

$$-\nabla^2 \varphi = 4\pi \rho = 4\pi [(\rho_i^G + \rho_i^{\text{pol}}) + (\rho_e^G + \underbrace{\rho_e^{\text{pol}}}_{\text{negligible}})] \quad (16)$$

Ion polarization leads to an interesting form of the GK Poisson equation.

One finds (for $T_i = 0$) $\rho_i^{\text{pol}} = (nq)_i \rho_s^2 \nabla_{\perp}^2 \Phi$, where $\Phi \doteq e\varphi/T_e$ and

$$\rho_s \doteq c_s/\omega_{ci} \quad (\text{"sound radius"; } \rho_s = \rho_i \text{ for } T_i = T_e), \quad (15a)$$

$$c_s \doteq (T_e/m_i)^{1/2} \quad (\text{sound speed}). \quad (15b)$$

Then Poisson's equation becomes

$$-\nabla^2 \varphi = 4\pi \rho = 4\pi [(\rho_i^G + \rho_i^{\text{pol}}) + (\rho_e^G + \underbrace{\rho_e^{\text{pol}}}_{\text{negligible}})] \quad (16)$$

or

$$-\nabla^2 \Phi = k_{De}^2 \left(\frac{n_i^G}{\bar{n}_i} + \rho_s^2 \nabla_{\perp}^2 \Phi - \frac{n_e^G}{\bar{n}_e} \right) \quad (17)$$

Ion polarization leads to an interesting form of the GK Poisson equation.

One finds (for $T_i = 0$) $\rho_i^{\text{pol}} = (nq)_i \rho_s^2 \nabla_{\perp}^2 \Phi$, where $\Phi \doteq e\varphi/T_e$ and

$$\rho_s \doteq c_s/\omega_{ci} \quad (\text{"sound radius"; } \rho_s = \rho_i \text{ for } T_i = T_e), \quad (15a)$$

$$c_s \doteq (T_e/m_i)^{1/2} \quad (\text{sound speed}). \quad (15b)$$

Then Poisson's equation becomes

$$-\nabla^2 \varphi = 4\pi \rho = 4\pi [(\rho_i^G + \rho_i^{\text{pol}}) + (\rho_e^G + \underbrace{\rho_e^{\text{pol}}}_{\text{negligible}})] \quad (16)$$

or

$$-\nabla^2 \Phi = k_{De}^2 \left(\frac{n_i^G}{\bar{n}_i} + \rho_s^2 \nabla_{\perp}^2 \Phi - \frac{n_e^G}{\bar{n}_e} \right) \quad (17)$$

or

$$-\left[\underbrace{\nabla^2}_{\text{original Poisson}} + \underbrace{\left(\frac{\rho_s^2}{\lambda_{De}^2} \right) \nabla_{\perp}^2}_{\text{ion polarization}} \right] \Phi = k_{De}^2 \left(\underbrace{\frac{n_i^G}{\bar{n}_i} - \frac{n_e^G}{\bar{n}_e}}_{\text{Note: GKE contains no polarization drift}} \right). \quad (18)$$

Ion polarization is intimately related to the **vorticity** of the plasma flow.

Across \vec{B} , the plasma moves predominantly with the $\vec{E} \times \vec{B}$ flow $\vec{u}_{\vec{E}} \propto \hat{b} \times \vec{\nabla} \varphi$. The vorticity associated with that flow is

$$\vec{\omega} = \vec{\nabla} \times \vec{u}_{\vec{E}} \propto \nabla_{\perp}^2 \varphi \hat{b} \quad (\text{for constant } B). \quad (19)$$




Fig. 8. Illustration of the intimate relationship between vorticity and polarization charge for a strongly magnetized plasma. A concentration of charge leads to an electric field; the plasma then rotates due to $\vec{E} \times \vec{B}$.

Ion polarization is responsible for the “dielectric constant of the gyrokinetic vacuum.”

We had

$$-\left[\underbrace{\nabla^2}_{\text{original Poisson}} + \underbrace{\left(\frac{\rho_s^2}{\lambda_{De}^2} \right) \nabla_{\perp}^2}_{\text{ion polarization}} \right] \Phi = k_{De}^2 \left(\underbrace{\frac{n_i^G}{\bar{n}_i} - \frac{n_e^G}{\bar{n}_e}}_{\text{Note: GKE contains no polarization drift}} \right). \quad (20)$$

Ion polarization is responsible for the “dielectric constant of the gyrokinetic vacuum.”

We had

$$-\left[\underbrace{\nabla^2}_{\text{original Poisson}} + \underbrace{\left(\frac{\rho_s^2}{\lambda_{De}^2} \right) \nabla_{\perp}^2}_{\text{ion polarization}} \right] \Phi = k_{De}^2 \left(\underbrace{\frac{n_i^G}{\bar{n}_i} - \frac{n_e^G}{\bar{n}_e}}_{\text{Note: GKE contains no polarization drift}} \right). \quad (20)$$

Compare with $\epsilon \vec{\nabla} \cdot \vec{E} = 4\pi\rho$ for a dielectric medium. Define

$$\epsilon^G \doteq \frac{\rho_s^2}{\lambda_{De}^2} = \frac{\omega_{pi}^2}{\omega_{ci}^2} \left(\begin{array}{l} \text{dielectric constant of the “GK vacuum”,} \\ \text{analogous to permittivity } \epsilon_0 \text{ of free space} \end{array} \right). \quad (21)$$

Ion polarization is responsible for the “dielectric constant of the gyrokinetic vacuum.”

We had

$$-\left[\underbrace{\nabla^2}_{\text{original Poisson}} + \underbrace{\left(\frac{\rho_s^2}{\lambda_{De}^2} \right) \nabla_{\perp}^2}_{\text{ion polarization}} \right] \Phi = k_{De}^2 \left(\underbrace{\frac{n_i^G}{\bar{n}_i} - \frac{n_e^G}{\bar{n}_e}}_{\text{Note: GKE contains no polarization drift}} \right). \quad (20)$$

Compare with $\epsilon \vec{\nabla} \cdot \vec{E} = 4\pi\rho$ for a dielectric medium. Define

$$\epsilon^G \doteq \frac{\rho_s^2}{\lambda_{De}^2} = \frac{\omega_{pi}^2}{\omega_{ci}^2} \left(\text{dielectric constant of the “GK vacuum”,} \right. \\ \left. \text{analogous to permittivity } \epsilon_0 \text{ of free space} \right). \quad (21)$$

Then

$$-\lambda_{De}^2 \left(\underbrace{\nabla^2}_{\text{original Poisson}} + \underbrace{\epsilon^G \nabla_{\perp}^2}_{\text{ion polarization}} \right) \Phi = \underbrace{\frac{n_i^G}{\bar{n}_i} - \frac{n_e^G}{\bar{n}_e}}_{\text{gyrocenter (charge) density}}. \quad (22)$$

For fusion applications, $\epsilon^G \gg 1$, so we can neglect the original ∇^2 and deal with the **quasineutrality condition**:

The GK Poisson equation is a statement of quasineutrality.

The quasineutrality condition $n_i \approx n_e$ can be expressed as

$$-\underbrace{\rho_s^2 \nabla_{\perp}^2 \Phi}_{\text{ion polarization}} = \underbrace{\frac{n_i^G}{\bar{n}_i} - \frac{n_e^G}{\bar{n}_e}}_{\text{net gyrocenter density}}. \quad (23)$$

(This form is correct only for $T_i = 0$, but it can be generalized.)

The GK Poisson equation is a statement of quasineutrality.

The quasineutrality condition $n_i \approx n_e$ can be expressed as

$$-\underbrace{\rho_s^2 \nabla_{\perp}^2 \Phi}_{\text{ion polarization}} = \underbrace{\frac{n_i^G}{\bar{n}_i} - \frac{n_e^G}{\bar{n}_e}}_{\text{net gyrocenter density}}. \quad (23)$$

(This form is correct only for $T_i = 0$, but it can be generalized.)

We have now achieved closure: the **gyrokinetic–Poisson system**:
GKE:

$$\partial_t F = \underbrace{\cdots + \langle \vec{V}_{\vec{E}} \rangle [\Phi] \cdot \vec{\nabla} F + \cdots}_{\text{right-hand side depends on potential(s)}}; \quad (24)$$

The GK Poisson equation is a statement of quasineutrality.

The quasineutrality condition $n_i \approx n_e$ can be expressed as

$$-\underbrace{\rho_s^2 \nabla_{\perp}^2 \Phi}_{\text{ion polarization}} = \underbrace{\frac{n_i^G}{\bar{n}_i} - \frac{n_e^G}{\bar{n}_e}}_{\text{net gyrocenter density}}. \quad (23)$$

(This form is correct only for $T_i = 0$, but it can be generalized.)

We have now achieved closure: the **gyrokinetic–Poisson system**:
GKE:

$$\partial_t F = \underbrace{\dots + \langle \vec{V}_{\vec{E}} \rangle [\Phi] \cdot \vec{\nabla} F + \dots}_{\text{right-hand side depends on potential(s)}}; \quad (24)$$

GK Poisson eq'n:

$$-\rho_s^2 \hat{\nabla}_{\perp}^2 \Phi = \frac{n_i^G}{\bar{n}_i} - \frac{n_e^G}{\bar{n}_e}. \quad (25)$$

The GK Poisson equation is a statement of quasineutrality.

The quasineutrality condition $n_i \approx n_e$ can be expressed as

$$-\underbrace{\rho_s^2 \nabla_{\perp}^2 \Phi}_{\text{ion polarization}} = \underbrace{\frac{n_i^G}{\bar{n}_i} - \frac{n_e^G}{\bar{n}_e}}_{\text{net gyrocenter density}}. \quad (23)$$

(This form is correct only for $T_i = 0$, but it can be generalized.)

We have now achieved closure: the **gyrokinetic–Poisson system**:
GKE:

$$\partial_t F = \dots + \underbrace{\langle \vec{V}_{\vec{E}} \rangle [\Phi] \cdot \vec{\nabla} F}_{\text{right-hand side depends on potential(s)}} + \dots; \quad (24)$$

GK Poisson eq'n:

$$-\rho_s^2 \hat{\nabla}_{\perp}^2 \Phi = \frac{n_i^G}{\bar{n}_i} - \frac{n_e^G}{\bar{n}_e}. \quad (25)$$

The GK Poisson equation is a statement of quasineutrality.

The quasineutrality condition $n_i \approx n_e$ can be expressed as

$$-\underbrace{\rho_s^2 \nabla_{\perp}^2 \Phi}_{\text{ion polarization}} = \underbrace{\frac{n_i^G}{\bar{n}_i} - \frac{n_e^G}{\bar{n}_e}}_{\text{net gyrocenter density}}. \quad (23)$$

(This form is correct only for $T_i = 0$, but it can be generalized.)

We have now achieved closure: the **gyrokinetic–Poisson system**:
GKE:

$$\partial_t F = \dots + \underbrace{\langle \vec{V}_{\vec{E}} \rangle [\Phi] \cdot \vec{\nabla} F}_{\text{right-hand side depends on potential(s)}} + \dots; \quad (24)$$

GK Poisson eq'n:

$$-\rho_s^2 \hat{\nabla}_{\perp}^2 \Phi = \frac{n_i^G}{\bar{n}_i} - \frac{n_e^G}{\bar{n}_e}. \quad (25)$$

Knowledge of the gyrokinetic distribution function enables one to calculate turbulent transport.

Example — Particle transport:

$$\frac{\partial \bar{n}}{\partial t} + \frac{\partial \bar{\Gamma}}{\partial x} = 0, \quad (26)$$

where the turbulent particle flux is $\bar{\Gamma} = \overline{\delta V_{E,x} \delta n}$. Also, heat, momentum,




Fig. 9. A randomly selected simulation result showing the approach to saturated heat flux from arbitrary initial conditions. From Wang et al., Phys. Plasmas 13, 092505 (2006).

The GK formalism leads readily to the **drift wave**.

● Continuity equation for ion gyrocenters ($T_i = 0$):

$$\partial_t n_i^G + \vec{\nabla}_\perp \cdot (\vec{u}_{\vec{E}} n_i^G) + \underbrace{\nabla_\parallel (u_{\parallel i} n_i^G)}_{\text{neglect}} = 0. \quad (27)$$

Linearize assuming $L_n^{-1} \doteq -\partial_x \ln \bar{n}(x)$:

$$\partial_t (\delta n_i^G / \bar{n}) + \underbrace{\delta \vec{u}_{\vec{E}} \cdot \vec{\nabla} \ln \bar{n}}_{V_* \partial_y \delta \Phi} = 0, \text{ where } \underbrace{V_* \doteq \rho_s c_s / L_n}_{\text{"diamagnetic velocity"}}. \quad (28)$$

The GK formalism leads readily to the **drift wave**.

- Continuity equation for ion gyrocenters ($T_i = 0$):

$$\partial_t n_i^G + \vec{\nabla}_\perp \cdot (\vec{u}_E n_i^G) + \underbrace{\nabla_\parallel (u_{\parallel i} n_i^G)}_{\text{neglect}} = 0. \quad (27)$$

Linearize assuming $L_n^{-1} \doteq -\partial_x \ln \bar{n}(x)$:

$$\partial_t (\delta n_i^G / \bar{n}) + \underbrace{\delta \vec{u}_E \cdot \vec{\nabla} \ln \bar{n}}_{V_* \partial_y \delta \Phi} = 0, \text{ where } \underbrace{V_* \doteq \rho_s c_s / L_n}_{\text{“diamagnetic velocity”}}. \quad (28)$$

- “Adiabatic” electron response: $\delta n_e / \bar{n}_e \rightarrow e^{e\delta\varphi/T_e} - 1 \approx \delta\Phi$.

The GK formalism leads readily to the **drift wave**.

- Continuity equation for ion gyrocenters ($T_i = 0$):

$$\partial_t n_i^G + \vec{\nabla}_\perp \cdot (\vec{u}_{\vec{E}} n_i^G) + \underbrace{\nabla_\parallel (u_{\parallel i} n_i^G)}_{\text{neglect}} = 0. \quad (27)$$

Linearize assuming $L_n^{-1} \doteq -\partial_x \ln \bar{n}(x)$:

$$\partial_t (\delta n_i^G / \bar{n}) + \underbrace{\delta \vec{u}_{\vec{E}} \cdot \vec{\nabla} \ln \bar{n}}_{V_* \partial_y \delta \Phi} = 0, \text{ where } \underbrace{V_* \doteq \rho_s c_s / L_n}_{\text{"diamagnetic velocity"}}. \quad (28)$$

- “Adiabatic” electron response: $\delta n_e / \bar{n}_e \rightarrow e^{e\delta\varphi/T_e} - 1 \approx \delta\Phi$.

- Quasineutrality: $-\rho_s^2 \nabla_\perp^2 \delta\Phi = \frac{\delta n_i^G}{\bar{n}_i} - \underbrace{\frac{\delta n_e^G}{\bar{n}_e}}_{\delta\Phi}.$

The GK formalism leads readily to the **drift wave**.

- Continuity equation for ion gyrocenters ($T_i = 0$):

$$\partial_t n_i^G + \vec{\nabla}_\perp \cdot (\vec{u}_{\vec{E}} n_i^G) + \underbrace{\nabla_\parallel (u_{\parallel i} n_i^G)}_{\text{neglect}} = 0. \quad (27)$$

Linearize assuming $L_n^{-1} \doteq -\partial_x \ln \bar{n}(x)$:

$$\partial_t (\delta n_i^G / \bar{n}) + \underbrace{\delta \vec{u}_{\vec{E}} \cdot \vec{\nabla} \ln \bar{n}}_{V_* \partial_y \delta \Phi} = 0, \text{ where } \underbrace{V_* \doteq \rho_s c_s / L_n}_{\text{"diamagnetic velocity"}}. \quad (28)$$

- “Adiabatic” electron response: $\delta n_e / \bar{n}_e \rightarrow e^{e\delta\varphi/T_e} - 1 \approx \delta\Phi$.

- Quasineutrality: $-\rho_s^2 \nabla_\perp^2 \delta\Phi = \frac{\delta n_i^G}{\bar{n}_i} - \underbrace{\frac{\delta n_e^G}{\bar{n}_e}}_{\delta\Phi}$.

Thus $\partial_t [(1 - \rho_s^2 \nabla_\perp^2) \delta\Phi] + V_* \partial_y \delta\Phi = 0,$

The GK formalism leads readily to the **drift wave**.

- Continuity equation for ion gyrocenters ($T_i = 0$):

$$\partial_t n_i^G + \vec{\nabla}_\perp \cdot (\vec{u}_E n_i^G) + \underbrace{\nabla_\parallel (u_{\parallel i} n_i^G)}_{\text{neglect}} = 0. \quad (27)$$

Linearize assuming $L_n^{-1} \doteq -\partial_x \ln \bar{n}(x)$:

$$\partial_t (\delta n_i^G / \bar{n}) + \underbrace{\delta \vec{u}_E \cdot \vec{\nabla} \ln \bar{n}}_{V_* \partial_y \delta \Phi} = 0, \text{ where } \underbrace{V_* \doteq \rho_s c_s / L_n}_{\text{"diamagnetic velocity"}}. \quad (28)$$

- "Adiabatic" electron response: $\delta n_e / \bar{n}_e \rightarrow e^{e\delta\varphi/T_e} - 1 \approx \delta\Phi$.

- Quasineutrality: $-\rho_s^2 \nabla_\perp^2 \delta\Phi = \frac{\delta n_i^G}{\bar{n}_i} - \underbrace{\frac{\delta n_e^G}{\bar{n}_e}}_{\delta\Phi}$.

Thus $\partial_t [(1 - \rho_s^2 \nabla_\perp^2) \delta\Phi] + V_* \partial_y \delta\Phi = 0$, or, with $\omega_* \doteq k_y V_*$,

$$\Omega_{\vec{k}} = \frac{\omega_*(\vec{k})}{1 + k_\perp^2 \rho_s^2} \quad (\text{drift wave}). \quad (29)$$

The drift wave propagates in the poloidal direction.




Fig. 10. Essential physics of the drift wave (polarization ignored).

- Assume $\varphi \sim \cos(k_y y - \Omega t)$.
- $\vec{E} \times \vec{B} \propto \hat{x}$ creates ion density fluctuations by advecting the background density profile.
- Those must be neutralized by electron flow along the field lines.
- For consistency, the wave must propagate upwards.

Nonlinearly, the **Hasegawa–Mima equation** emerges.

- Continuity equation for ion gyrocenters ($T_i = 0$):

$$\partial_t n_i^G + \vec{\nabla}_\perp \cdot (\vec{u}_{\vec{E}} n_i^G) + \underbrace{\nabla_\parallel (u_{\parallel i} n_i^G)}_{\text{neglect}} = 0. \quad (30)$$

$$\frac{\partial(\delta n_i^G / \bar{n})}{\partial t} + \underbrace{\delta \vec{u}_{\vec{E}} \cdot \vec{\nabla} \ln \bar{n}}_{V_* \partial_y \delta \Phi} + \delta \vec{u}_{\vec{E}} \cdot \vec{\nabla} (\delta n_i^G / \bar{n}) - \langle \dots \rangle = 0. \quad (31)$$

- “Adiabatic” electron response: $\delta n_e / \bar{n}_e = \delta \Phi$.

- Quasineutrality: $-\rho_s^2 \nabla_\perp^2 \delta \Phi = \frac{\delta n_i^G}{\bar{n}_i} - \underbrace{\frac{\delta n_e^G}{\bar{n}_e}}_{\delta \Phi}.$

Thus we obtain the **Hasegawa–Mima equation**:

$$\left(\underbrace{1}_{\text{adiab. elect.}} - \underbrace{\rho_s^2 \nabla_\perp^2}_{\text{ion polarization}} \right) \frac{\partial \delta \Phi}{\partial t} + \underbrace{V_* \frac{\partial \delta \Phi}{\partial y}}_{\text{linear drift wave}} + \underbrace{\vec{u}_{\vec{E}} \cdot \vec{\nabla} (-\rho_s^2 \nabla_\perp^2 \delta \Phi)}_{\text{nonlinear } \vec{E} \times \vec{B} \text{ advection of vorticity}} = 0.$$

The Hasegawa–Mima equation is not suitable for the treatment of **zonal flows**!

Zonal flow: $\varphi(r, \theta, \phi) \rightarrow \varphi(r, \cancel{\theta}, \cancel{\phi}) \Rightarrow \vec{V}_E = V_{\vec{E}, \text{pol}}(r) \hat{\theta}.$

The Hasegawa–Mima equation is not suitable for the treatment of **zonal flows**!

Zonal flow: $\varphi(r, \theta, \phi) \rightarrow \varphi(r, \cancel{\theta}, \cancel{\phi}) \Rightarrow \vec{V}_E = V_{\vec{E}, \text{pol}}(r) \hat{\theta}$.

 Zonal flows are *sheared poloidal flows* (“zonal jets”).

The Hasegawa–Mima equation is not suitable for the treatment of **zonal flows**!

Zonal flow: $\varphi(r, \theta, \phi) \rightarrow \varphi(r, \cancel{\theta}, \cancel{\phi}) \Rightarrow \vec{V}_E = V_{\vec{E}, \text{pol}}(r) \hat{\theta}$.

- Zonal flows are *sheared poloidal flows* (“zonal jets”).
- Their shear can regulate the nonlinear saturation of drift waves.

The Hasegawa–Mima equation is not suitable for the treatment of **zonal flows**!

Zonal flow: $\varphi(r, \theta, \phi) \rightarrow \varphi(r, \cancel{\theta}, \cancel{\phi}) \Rightarrow \vec{V}_E = V_{\vec{E}, \text{pol}}(r) \hat{\theta}$.

- Zonal flows are *sheared poloidal flows* (“zonal jets”).
- Their shear can regulate the nonlinear saturation of drift waves.
- The Hasegawa–Mima equation must be *modified* in order to describe zonal flows (which do not permit adiabatic electron response because $k_{\parallel} = 0 \Rightarrow \omega/k_{\parallel} v_{te} \gg 1$, i.e., *fluid regime*). Let

$$\hat{\alpha} \doteq \begin{cases} 1 & (k_{\parallel} \neq 0) \\ 0 & (k_{\parallel} = 0). \end{cases} \quad (33)$$

The Hasegawa–Mima equation is not suitable for the treatment of **zonal flows**!

Zonal flow: $\varphi(r, \theta, \phi) \rightarrow \varphi(r, \cancel{\theta}, \cancel{\phi}) \Rightarrow \vec{V}_E = V_{\vec{E}, \text{pol}}(r) \hat{\theta}$.

- Zonal flows are *sheared poloidal flows* (“zonal jets”).
- Their shear can regulate the nonlinear saturation of drift waves.
- The Hasegawa–Mima equation must be *modified* in order to describe zonal flows (which do not permit adiabatic electron response because $k_{\parallel} = 0 \Rightarrow \omega/k_{\parallel} v_{te} \gg 1$, i.e., *fluid regime*). Let

$$\hat{\alpha} \doteq \begin{cases} 1 & (k_{\parallel} \neq 0) \\ 0 & (k_{\parallel} = 0). \end{cases} \quad (33)$$

Then we obtain the **modified Hasegawa–Mima equation**:

$$\left(\underbrace{\hat{\alpha}}_{\text{elect. response}} - \underbrace{\rho_s^2 \nabla_{\perp}^2}_{\text{ion polarization}} \right) \frac{\partial \delta \Phi}{\partial t} + \underbrace{V_* \frac{\partial \delta \Phi}{\partial y}}_{\text{linear drift wave}} + \underbrace{\vec{u}_{\vec{E}} \cdot \vec{\nabla} [(\hat{\alpha} - \rho_s^2 \nabla_{\perp}^2) \delta \Phi]}_{\text{nonlinear } \vec{E} \times \vec{B} \text{ advection of ion density}} = 0.$$

Zonal flows are driven by drift-wave Reynolds stresses.

For zonal flows, the dominant terms in the modified Hasegawa–Mima equation are

$$\frac{\partial(\nabla_{\perp}^2 \varphi^Z)}{\partial t} + \vec{\nabla} \cdot (\vec{V}_E^D \nabla_{\perp}^2 \varphi^D) = 0. \quad (35)$$

Average over (periodic) poloidal direction y :

$$\frac{\partial}{\partial t} \left(\frac{\partial^2 \overline{\varphi^Z}}{\partial x^2} \right) + \frac{\partial}{\partial x} \left[- \frac{\partial \varphi^D}{\partial y} \left(\frac{\partial^2 \varphi^D}{\partial x^2} + \frac{\partial^2 \varphi^D}{\partial y^2} \right) \right] = 0. \quad (36)$$

Note that $\partial \varphi / \partial x = u_y$. Integrate in x . Manipulate to find

$$\underbrace{\frac{\partial \overline{u}_y^Z(x)}{\partial t}}_{\text{sheared zonal flows}} + \frac{\partial}{\partial x} \left(\underbrace{\overline{\delta u_x^D \delta u_y^D}}_{\text{Reynolds stresses}} \right) = 0. \quad (37)$$

Self-consistent saturated turbulent states involve coupled **drift waves** and **zonal flows**.




Fig. 11. Drift waves and zonal flows form a self-regulating coupled system.




Fig. 12. Drift-wave (\vec{k}) sidebands (\vec{p}_+ and \vec{p}_-) couple to drive zonal flows (\vec{q}).



Fig. 13. Zonal-flow shearing in New York City. Zonal flows destroy drift-wave eddies \Rightarrow smaller turbulent transport.

Self-consistent saturated turbulent states involve coupled **drift waves** and **zonal flows**.




Fig. 11. Drift waves and zonal flows form a self-regulating coupled system.




Fig. 12. Drift-wave (\vec{k}) sidebands (\vec{p}_+ and \vec{p}_-) couple to drive zonal flows (\vec{q}).



Fig. 13. Zonal-flow shearing in New York City. Zonal flows destroy drift-wave eddies \Rightarrow smaller turbulent transport.

🔴 Statistical closure theory can be used to calculate the zonal flow generation rate.

Self-consistent saturated turbulent states involve coupled **drift waves** and **zonal flows**.




Fig. 11. Drift waves and zonal flows form a self-regulating coupled system.




Fig. 12. Drift-wave (\vec{k}) sidebands (\vec{p}_+ and \vec{p}_-) couple to drive zonal flows (\vec{q}).



Fig. 13. Zonal-flow shearing in New York City. Zonal flows destroy drift-wave eddies \Rightarrow smaller turbulent transport.

- Statistical closure theory can be used to calculate the zonal flow generation rate.
- Kraichnan's theory of 2D Navier–Stokes eddy viscosity can be recovered as a special limit.

Self-consistent saturated turbulent states involve coupled **drift waves** and **zonal flows**.




Fig. 11. Drift waves and zonal flows form a self-regulating coupled system.




Fig. 12. Drift-wave (\vec{k}) sidebands (\vec{p}_+ and \vec{p}_-) couple to drive zonal flows (\vec{q}).



Fig. 13. Zonal-flow shearing in New York City. Zonal flows destroy drift-wave eddies \Rightarrow smaller turbulent transport.

- Statistical closure theory can be used to calculate the zonal flow generation rate.
- Kraichnan's theory of 2D Navier–Stokes eddy viscosity can be recovered as a special limit.
- Connections to field-theoretic Hamiltonian theory and Casimir invariants can be demonstrated. . . . (But let's proceed with more discussion of GKs. . .)

Zonal flows can eat drift waves.




Fig. 14. Projection of a 10D center-manifold reduction of PDEs for ion-temperature-gradient-driven (ITG) turbulence onto two ITG amplitudes (x and y axes) and a zonal amplitude (z axis). Only the zonal mode survives for these parameters.

Zonal-flow suppression of drift waves can delay the onset of turbulence.




Fig. 15. (a) One might expect that linear instability would drive ITG turbulence and heat flux. (b) In actuality, zonal modes suppress turbulence onset for some region (the *Dimitis shift*) above linear threshold.

Summary of Heuristic Gyrokinetics

- **Gyrocenters** move (only) with effective $\vec{E} \times \vec{B}$ drifts and magnetic drifts.

Summary of Heuristic Gyrokinetics

- **Gyrocenters** move (only) with effective $\vec{E} \times \vec{B}$ drifts and magnetic drifts.
- Knowing those drifts, one can write the **gyrokinetic equation**.

Summary of Heuristic Gyrokinetics

- **Gyrocenters** move (only) with effective $\vec{E} \times \vec{B}$ drifts and magnetic drifts.
- Knowing those drifts, one can write the **gyrokinetic equation**.
- In time-varying fields, particles **polarize** with respect to the gyrocenter motion.

Summary of Heuristic Gyrokinetics

- **Gyrocenters** move (only) with effective $\vec{E} \times \vec{B}$ drifts and magnetic drifts.
- Knowing those drifts, one can write the **gyrokinetic equation**.
- In time-varying fields, particles **polarize** with respect to the gyrocenter motion.
- Polarization and **vorticity** are intimately related.

Summary of Heuristic Gyrokinetics

- **Gyrocenters** move (only) with effective $\vec{E} \times \vec{B}$ drifts and magnetic drifts.
- Knowing those drifts, one can write the **gyrokinetic equation**.
- In time-varying fields, particles **polarize** with respect to the gyrocenter motion.
- Polarization and **vorticity** are intimately related.
- Ion polarization endows the “**gyrokinetic vacuum**” with a large dielectric permittivity ϵ^G .

Summary of Heuristic Gyrokinetics

- **Gyrocenters** move (only) with effective $\vec{E} \times \vec{B}$ drifts and magnetic drifts.
- Knowing those drifts, one can write the **gyrokinetic equation**.
- In time-varying fields, particles **polarize** with respect to the gyrocenter motion.
- Polarization and **vorticity** are intimately related.
- Ion polarization endows the “**gyrokinetic vacuum**” with a large dielectric permittivity ϵ^G .
- For large ϵ^G , the GK Poisson equation reduces to **quasineutrality**:

$$(\text{net gyrocenter charge}) + (\text{ion polarization charge}) \approx 0. \quad (38)$$

Summary of Heuristic Gyrokinetics

- **Gyrocenters** move (only) with effective $\vec{E} \times \vec{B}$ drifts and magnetic drifts.
- Knowing those drifts, one can write the **gyrokinetic equation**.
- In time-varying fields, particles **polarize** with respect to the gyrocenter motion.
- Polarization and **vorticity** are intimately related.
- Ion polarization endows the “**gyrokinetic vacuum**” with a large dielectric permittivity ϵ^G .
- For large ϵ^G , the GK Poisson equation reduces to **quasineutrality**:

$$(\text{net gyrocenter charge}) + (\text{ion polarization charge}) \approx 0. \quad (38)$$

- Gyrokinetics predicts a turbulent “soup” of **interacting drift waves and zonal flows**.

Summary of Heuristic Gyrokinetics

- **Gyrocenters** move (only) with effective $\vec{E} \times \vec{B}$ drifts and magnetic drifts.
- Knowing those drifts, one can write the **gyrokinetic equation**.
- In time-varying fields, particles **polarize** with respect to the gyrocenter motion.
- Polarization and **vorticity** are intimately related.
- Ion polarization endows the “**gyrokinetic vacuum**” with a large dielectric permittivity ϵ^G .
- For large ϵ^G , the GK Poisson equation reduces to **quasineutrality**:

$$(\text{net gyrocenter charge}) + (\text{ion polarization charge}) \approx 0. \quad (38)$$

- Gyrokinetics predicts a turbulent “soup” of **interacting drift waves and zonal flows**.
- The GK–Poisson system can be simulated and used to calculate **turbulent transport fluxes**.

So far we have been doing handwaving. BUT: Various things have been swept under the rug!

- Most fundamentally, the formalism assumes that μ is conserved. But in reality, what most people call μ , namely $\mu \doteq \frac{1}{2}mv_{\perp}^2/\omega_c(\vec{x})$, is not exactly conserved. (It is not even Galilean invariant!) *What quantity $\bar{\mu}$ is really conserved?*

So far we have been doing handwaving. BUT: Various things have been swept under the rug!

- Most fundamentally, the formalism assumes that μ is conserved. But in reality, what most people call μ , namely $\mu \doteq \frac{1}{2}mv_{\perp}^2/\omega_c(\vec{x})$, is not exactly conserved. (It is not even Galilean invariant!) *What quantity $\bar{\mu}$ is really conserved?*
- Also, how does one generalize the notion of gyrokinetic polarization for $T_i \neq 0$ and $k_{\perp}\rho_i = O(1)$?

So far we have been doing handwaving. BUT: Various things have been swept under the rug!

- Most fundamentally, the formalism assumes that μ is conserved. But in reality, what most people call μ , namely $\mu \doteq \frac{1}{2}mv_{\perp}^2/\omega_c(\vec{x})$, is not exactly conserved. (It is not even Galilean invariant!) *What quantity $\bar{\mu}$ is really conserved?*
- Also, how does one generalize the notion of gyrokinetic polarization for $T_i \neq 0$ and $k_{\perp}\rho_i = O(1)$?
- When appropriately formulated, gyrokinetics is a Hamiltonian system (very useful to know both analytically and numerically). However, that is not apparent from the “derivation” given so far.

So far we have been doing handwaving. BUT: Various things have been swept under the rug!

- Most fundamentally, the formalism assumes that μ is conserved. But in reality, what most people call μ , namely $\mu \doteq \frac{1}{2}mv_{\perp}^2/\omega_c(\vec{x})$, is not exactly conserved. (It is not even Galilean invariant!) *What quantity $\bar{\mu}$ is really conserved?*
- Also, how does one generalize the notion of gyrokinetic polarization for $T_i \neq 0$ and $k_{\perp}\rho_i = O(1)$?
- When appropriately formulated, gyrokinetics is a Hamiltonian system (very useful to know both analytically and numerically). However, that is not apparent from the “derivation” given so far.
- Finally, are we sure that the gyrokinetic system conserves the appropriate things (e.g., energy or momentum)?

Gyrokinetics has an interesting intellectual history.

Catto (1978)

— Linearized gyrokinetics

Frieman & Chen (1982)

— First **nonlinear** gyrokinetic equation (GKE)

Lee (1983)

— Reformulated GKE for **particle simulation**

Gyrokinetics has an interesting intellectual history.

Catto (1978)

— Linearized gyrokinetics

Littlejohn (1979)

— “A guiding center Hamiltonian”

Frieman & Chen (1982)

— First nonlinear gyrokinetic equation (GKE)

Littlejohn (1982)

— Hamiltonian perturbation theory in noncanonical coordinates

Lee (1983)

— Reformulated GKE for particle simulation

Gyrokinetics has an interesting intellectual history.

Catto (1978)

— Linearized gyrokinetics

Littlejohn (1979)

— “A guiding center Hamiltonian”

Frieman & Chen (1982)

— First nonlinear gyrokinetic equation (GKE)

Littlejohn (1982)

— Hamiltonian perturbation theory in noncanonical coordinates

Lee (1983)

— Reformulated GKE for particle simulation

Dubin, Krommes, et al. (1983)

— Hamiltonian formulation of the nonlinear GK–Poisson system

Gyrokinetics has an interesting intellectual history.

Catto (1978)

— Linearized gyrokinetics

Littlejohn (1979)

— “A guiding center Hamiltonian”

Frieman & Chen (1982)

— First nonlinear gyrokinetic equation (GKE)

Littlejohn (1982)

— Hamiltonian perturbation theory in noncanonical coordinates

Lee (1983)

— Reformulated GKE for particle simulation

Dubin, Krommes, et al. (1983)

— Hamiltonian formulation of the nonlinear GK–Poisson system

Cary & Littlejohn (1983)

— Noncanonical Lagrangian mechanics

Gyrokinetics has an interesting intellectual history.

Catto (1978)

— Linearized gyrokinetics

Littlejohn (1979)

— “A guiding center Hamiltonian”

Frieman & Chen (1982)

— First nonlinear gyrokinetic equation (GKE)

Littlejohn (1982)

— Hamiltonian perturbation theory in noncanonical coordinates

Lee (1983)

— Reformulated GKE for particle simulation

Dubin, Krommes, et al. (1983)

— Hamiltonian formulation of the nonlinear GK–Poisson system

Cary & Littlejohn (1983)

— Noncanonical Lagrangian mechanics

Hahm (1988)

— Nonlinear gyrokinetics via the Lagrangian one-form method

Brizard (1989)

— More on the Lagrangian method

Qin (1998)

— Gyrokinetics and electromagnetics

Brizard & Hahm (2007)

— Review article: “Foundations of nonlinear gyrokinetic theory”

Gyrokinetics has an interesting intellectual history.

- | | |
|--------------------------------|---|
| Catto (1978) | — Linearized gyrokinetics |
| Littlejohn (1979) | — “A guiding center Hamiltonian” |
| Frieman & Chen (1982) | — First nonlinear gyrokinetic equation (GKE) |
| Littlejohn (1982) | — Hamiltonian perturbation theory in noncanonical coordinates |
| Lee (1983) | — Reformulated GKE for particle simulation |
| Dubin, Krommes, et al. (1983) | — Hamiltonian formulation of the nonlinear GK–Poisson system |
| Cary & Littlejohn (1983) | — Noncanonical Lagrangian mechanics |
| Hahm (1988) | — Nonlinear gyrokinetics via the Lagrangian one-form method |
| Brizard (1989) | — More on the Lagrangian method |
| Qin (1998) | — Gyrokinetics and electromagnetics |
| Brizard & Hahm (2007) | — Review article: “Foundations of nonlinear gyrokinetic theory” |
| Sugiyama; Parra & Catto (2008) | — Various <i>ATTACKS</i> on the foundations (more discussion later) |

Table of Contents

- ✓ ■ Introduction: Heuristic Gyrokinetics
- Systematic Gyrokinetics, Part I: The Gyrokinetic Equations of Motion
- Systematic Gyrokinetics, Part II: The Gyrokinetic Poisson Equation
- The Current Status of Gyrokinetics

Systematic Gyrokinetics, Part I: The Gyrokinetic Equations of Motion

Modern gyrokinetics via the Lagrangian one-form

Goals:

- Systematic derivation of gyrokinetics from first principles (Newton's laws and Maxwell's equations).
- Asymptotic construction of the “true” adiabatic invariant $\bar{\mu}$ (in complicated geometry with $\vec{\nabla} B \neq \vec{0}$ and also $\partial_t \vec{E} \neq \vec{0}$).

Modern gyrokinetics via the Lagrangian one-form

Goals:

- Systematic derivation of gyrokinetics from first principles (Newton's laws and Maxwell's equations).
- Asymptotic construction of the “true” adiabatic invariant $\bar{\mu}$ (in complicated geometry with $\vec{\nabla} B \neq \vec{0}$ and also $\partial_t \vec{E} \neq \vec{0}$).

Methodology (following Littlejohn):

- Lagrange's variational principle
- Differential forms
- Lie perturbation theory

Modern gyrokinetics via the Lagrangian one-form

Goals:

- Systematic derivation of gyrokinetics from first principles (Newton's laws and Maxwell's equations).
- Asymptotic construction of the “true” adiabatic invariant $\bar{\mu}$ (in complicated geometry with $\vec{\nabla} B \neq \vec{0}$ and also $\partial_t \vec{E} \neq \vec{0}$).

Methodology (following Littlejohn):

- Lagrange's variational principle
- Differential forms
- Lie perturbation theory

Remark: Much of analytical physics consists of finding the “best” variables. For example, contrast

- \vec{x} — particle position
- \vec{X} — *lowest-order* gyrocenter position (for circular motion)
- $\bar{\vec{X}}$ — “true” gyrocenter position

We'll transform $\{\vec{x}, \vec{v}\} \Rightarrow \{\vec{X}, U, \mu, \zeta\} \Rightarrow \{\bar{\vec{X}}, \bar{U}, \bar{\mu}, \bar{\zeta}\}$.

The dynamical equations of particle motion follow from **Lagrange's variational principle**.

$$0 = \delta \int_{t_0}^{t_1} L dt \quad (L \doteq \vec{p} \cdot \dot{\vec{q}} - H) \quad (39)$$

recovers the particle equations of motion.

The dynamical equations of particle motion follow from **Lagrange's variational principle**.

$$0 = \delta \int_{t_0}^{t_1} L dt \quad (L \doteq \vec{p} \cdot \dot{\vec{q}} - H) \quad (39)$$

recovers the particle equations of motion. In the language of differential forms,

$$0 = \delta \int_{t_0}^{t_1} \gamma, \quad (40)$$

where the **fundamental differential one-form** is

$$\gamma \doteq \vec{p} \cdot d\vec{q} - H dt. \quad (41)$$

The dynamical equations of particle motion follow from **Lagrange's variational principle**.

$$0 = \delta \int_{t_0}^{t_1} L dt \quad (L \doteq \vec{p} \cdot \dot{\vec{q}} - H) \quad (39)$$

recovers the particle equations of motion. In the language of differential forms,

$$0 = \delta \int_{t_0}^{t_1} \gamma, \quad (40)$$

where the **fundamental differential one-form** is

$$\gamma \doteq \vec{p} \cdot d\vec{q} - H dt. \quad (41)$$

- The variational principle does not care about the particular variables used to express γ .

The dynamical equations of particle motion follow from **Lagrange's variational principle**.

$$0 = \delta \int_{t_0}^{t_1} L dt \quad (L \doteq \vec{p} \cdot \dot{\vec{q}} - H) \quad (39)$$

recovers the particle equations of motion. In the language of differential forms,

$$0 = \delta \int_{t_0}^{t_1} \gamma, \quad (40)$$

where the **fundamental differential one-form** is

$$\gamma \doteq \vec{p} \cdot d\vec{q} - H dt. \quad (41)$$

- The variational principle does not care about the particular variables used to express γ .
- Therefore, we are free to use “**better**” **variables** (chosen to satisfy some criterion); then the variational principle gives us the equations of motion for those variables.

One can cast γ into a representation-free form.

Define

$$z^\nu \doteq \{t, \vec{x}, \vec{p}\}, \quad (42a)$$

$$\gamma_\nu \doteq \{-H, \vec{p}, \vec{0}\}. \quad (42b)$$

Then the one-form can be written **covariantly** as

$$\gamma = \gamma_\nu dz^\nu \quad (\text{summation convention}). \quad (43)$$

Obviously this can be transformed to any set of variables one pleases:

$$\gamma(z) = \overline{\gamma}(\overline{z}) = \overline{\gamma}_\nu d\overline{z}^\nu. \quad (44)$$

For any z^ν 's, the Euler–Lagrange equations of motion are

$$\omega_{\sigma\nu} \frac{dz^\nu}{dt} = 0, \quad (45)$$

where the **symplectic two-form** is

$$\omega_{\sigma\nu} \doteq \frac{\partial \gamma_\nu}{\partial z^\sigma} - \frac{\partial \gamma_\sigma}{\partial z^\nu}. \quad (46)$$

**Symmetries of the Lagrangian
are related to conservation laws.
In particular, Noether's Theorem is fundamental.**

Noether's Theorem:

If all of the γ_ν 's are independent
of a particular coordinate z^α ,
then γ_α is conserved.

I.e.,

$$\begin{aligned}\gamma = & \gamma_1(z^1, \dots, \cancel{z^\alpha}, \dots, z^n) dz^1 + \dots \\ & + \underbrace{\gamma_\alpha(z^1, \dots, \cancel{z^\alpha}, \dots, z^n)}_{\text{conserved!}} dz^\alpha + \dots \\ & + \gamma_n(z^1, \dots, \cancel{z^\alpha}, \dots, z^n) dz^n.\end{aligned}\tag{47}$$

Proof: *A simple exercise left to the reader.*

A special application of Noether's Theorem leads to the conserved magnetic moment $\bar{\mu}$.

Strategy: Exploit **gyrophase independence of gyrocenter motion** (i.e., gyrocenters possess a **gyrational symmetry**).

🔴 Consider gyrocenter coordinates $\bar{z}^\nu = \{\bar{\vec{X}}, \bar{U}, \bar{\mu}, \bar{\zeta}\}$ (to be defined).

A special application of Noether's Theorem leads to the conserved magnetic moment $\bar{\mu}$.

Strategy: Exploit **gyrophase independence of gyrocenter motion** (i.e., gyrocenters possess a **gyrational symmetry**).

- Consider gyrocenter coordinates $\bar{z}^\nu = \{\bar{\vec{X}}, \bar{U}, \bar{\mu}, \bar{\zeta}\}$ (to be defined).
- Choose $\bar{z}^\alpha = \bar{\zeta}$ (“true” gyrophase).

A special application of Noether's Theorem leads to the conserved magnetic moment $\bar{\mu}$.

Strategy: Exploit **gyrophase independence of gyrocenter motion** (i.e., gyrocenters possess a **gyrational symmetry**).

- Consider gyrocenter coordinates $\bar{z}^\nu = \{\bar{\vec{X}}, \bar{U}, \bar{\mu}, \bar{\zeta}\}$ (to be defined).
- Choose $\bar{z}^\alpha = \bar{\zeta}$ (“true” gyrophase).
- Arrange things such that $\bar{\gamma} = \bar{\gamma}_{(\nu)} d\bar{z}^{(\nu)} + \bar{\mu} d\bar{\zeta}$.

A special application of Noether's Theorem leads to the conserved magnetic moment $\bar{\mu}$.

Strategy: Exploit **gyrophase independence of gyrocenter motion** (i.e., gyrocenters possess a **gyrational symmetry**).

- Consider gyrocenter coordinates $\bar{z}^\nu = \{\bar{\vec{X}}, \bar{U}, \bar{\mu}, \bar{\zeta}\}$ (to be defined).
- Choose $\bar{z}^\alpha = \bar{\zeta}$ (“true” gyrophase).
- Arrange things such that $\bar{\gamma} = \bar{\gamma}_{(\nu)} d\bar{z}^{(\nu)} + \bar{\mu} d\bar{\zeta}$.
- Choose the $\bar{\gamma}_\nu$'s so that $\frac{\partial \bar{\gamma}_\nu}{\partial \bar{\zeta}} = 0 \quad (\forall \nu)$.

A special application of Noether's Theorem leads to the conserved magnetic moment $\bar{\mu}$.

Strategy: Exploit **gyrophase independence of gyrocenter motion** (i.e., gyrocenters possess a **gyrational symmetry**).

- Consider gyrocenter coordinates $\bar{z}^\nu = \{\bar{X}, \bar{U}, \bar{\mu}, \bar{\zeta}\}$ (to be defined).
- Choose $\bar{z}^\alpha = \bar{\zeta}$ (“true” gyrophase).
- Arrange things such that $\bar{\gamma} = \bar{\gamma}_{(\nu)} d\bar{z}^{(\nu)} + \bar{\mu} d\bar{\zeta}$.
- Choose the $\bar{\gamma}_\nu$'s so that $\frac{\partial \bar{\gamma}_\nu}{\partial \bar{\zeta}} = 0 \quad (\forall \nu)$.
- Then, according to Noether's theorem, **$\bar{\mu}$ will be conserved.**

A special application of Noether's Theorem leads to the conserved magnetic moment $\bar{\mu}$.

Strategy: Exploit **gyrophase independence of gyrocenter motion** (i.e., gyrocenters possess a **gyrational symmetry**).

- Consider gyrocenter coordinates $\bar{z}^\nu = \{\bar{\vec{X}}, \bar{U}, \bar{\mu}, \bar{\zeta}\}$ (to be defined).
- Choose $\bar{z}^\alpha = \bar{\zeta}$ (“true” gyrophase).
- Arrange things such that $\bar{\gamma} = \bar{\gamma}_{(\nu)} d\bar{z}^{(\nu)} + \bar{\mu} d\bar{\zeta}$.
- Choose the $\bar{\gamma}_\nu$'s so that $\frac{\partial \bar{\gamma}_\nu}{\partial \bar{\zeta}} = 0 \quad (\forall \nu)$.
- Then, according to Noether's theorem, **$\bar{\mu}$ will be conserved.**

Gyrokinetics is derived
by removing gyrophase dependence from $\bar{\gamma}$.
As a byproduct, one obtains
formulas for the conserved $\bar{\mu}$
and the other gyrocenter variables.

How does gyrophase dependence enter the one-form?

In canonical variables (\vec{x}, \vec{p}) , one has

$$\gamma = \vec{p} \cdot d\vec{x} - H dt, \quad (48)$$

with $\vec{p} \doteq m\vec{v} + (q/c)\vec{A}(\vec{x})$ and $H \doteq p^2/2m + q\varphi(\vec{x}, t)$. With this form, it is not obvious that the particle gyrates around a magnetic field line; the gyrophase ζ is not apparent. *It's better to use noncanonical variables...*




Fig. 16. Illustration of the lowest-order gyrokinetic variables. \hat{a} and \hat{c} rotate with the particle. Vectors can be decomposed with respect to either of the orthonormal sets $\{\hat{a}, \hat{b}, \hat{c}\}$ or $\{\hat{e}_1, \hat{e}_2, \hat{b}\}$.

Gyration is more apparent by using the lowest-order gyrocenter variables.

Instead of working with $\vec{z} \doteq \{\vec{x}, \vec{v}\}$, let's use the *lowest-order gyrocenter variables* $\vec{Z} \doteq \{\vec{x}, U, \mu, \zeta\}$. Then

$$\gamma = \underbrace{\frac{q}{c} \vec{A} \cdot d\vec{x}}_{O(\epsilon^{-1})} + \underbrace{m[U\hat{\mathbf{b}}(\vec{x}) + v_{\perp}\hat{\mathbf{c}}(\zeta, \vec{x})]}_{O(1)} \cdot d\vec{x} - \underbrace{\left(\frac{1}{2}mU^2 + \mu\omega_c(\vec{x})\right)}_{O(1)} dt + \underbrace{q\varphi(\vec{x}, t)dt}_{O(\epsilon)}. \quad (49)$$

gyrophase dependence

We used

$$\frac{mv_{\perp}}{qA/c} = \frac{v_{\perp}}{L_B[q(A/L_B)/mc]} = \frac{v_{\perp}/\omega_c}{L_B} = \frac{\rho}{L_B} = \epsilon_B. \quad (50)$$



The fundamental expansion parameter ϵ is

- the size of the magnetic inhomogeneity, $\epsilon = \epsilon_B \doteq \rho/L_B$;
- the size of the fluctuating fields, e.g., $\epsilon = \epsilon_{\varphi} \doteq e\delta\varphi/T_e$.

We are now set to remove ζ dependence (perturbatively) order by order.

We have

$$\gamma = \gamma^{(-1)} + \underbrace{\gamma^{(0)}}_{\zeta\text{-dependent}} + \gamma^{(1)}. \quad (51)$$

We are now set to remove ζ dependence (perturbatively) order by order.

We have

$$\gamma = \gamma^{(-1)} + \underbrace{\gamma^{(0)}}_{\zeta\text{-dependent}} + \gamma^{(1)}. \quad (51)$$

🔴 Change variables (perturbatively) from \vec{Z} to

$$\vec{\bar{Z}} = \underbrace{T(\epsilon)}_{\substack{\text{differential} \\ \text{transformation} \\ \text{operator}}} \vec{Z}. \quad (52)$$

We are now set to remove ζ dependence (perturbatively) order by order.

We have

$$\gamma = \gamma^{(-1)} + \underbrace{\gamma^{(0)}}_{\zeta\text{-dependent}} + \gamma^{(1)}. \quad (51)$$

🔴 Change variables (perturbatively) from \vec{Z} to

$$\vec{\bar{Z}} = \underbrace{T(\epsilon)}_{\substack{\text{differential} \\ \text{transformation} \\ \text{operator}}} \vec{Z}. \quad (52)$$

🔴 Determine $T(\epsilon)$ by demanding that the components of the new one-form $\bar{\gamma}$ are independent of $\bar{\zeta}$.

We are now set to remove ζ dependence (perturbatively) order by order.

We have

$$\gamma = \gamma^{(-1)} + \underbrace{\gamma^{(0)}}_{\zeta\text{-dependent}} + \gamma^{(1)}. \quad (51)$$

• Change variables (perturbatively) from \vec{Z} to

$$\vec{\bar{Z}} = \underbrace{T(\epsilon)}_{\substack{\text{differential} \\ \text{transformation} \\ \text{operator}}} \vec{Z}. \quad (52)$$

• Determine $T(\epsilon)$ by **demanding that the components of the new one-form $\bar{\gamma}$ are independent of $\bar{\zeta}$.**

• Then $\bar{\gamma}_{\bar{\zeta}} \equiv \bar{\mu}$ is conserved (by Noether's theorem).

How does the one-form γ transform?

The value of γ is independent of coordinate system: $\bar{\gamma}(\bar{z}) = \gamma(z)$.

Recall that $\bar{z} = Tz$. Then

$$\underbrace{T}_{\text{“pull-back”}} \bar{\gamma}(\bar{z}) = \gamma(\bar{z}), \quad (53a)$$

$$\bar{\gamma}(z) = \underbrace{T^{-1}}_{\text{“push-forward”}} \gamma(z). \quad (53b)$$

- Also, note that the differential of a scalar function S does not contribute to the equations of motion: $\delta \int_{t_0}^{t_1} dS = 0$.
- The “gauge scalar” S gives us extra freedom that helps us achieve the goal of removing gyrophase dependence.
- Thus the basic formula is

$$\cancel{\bar{\gamma}(\zeta)} = \underbrace{T^{-1}(\zeta)}_{\text{push-forward transformation}} \gamma(\zeta) + \underbrace{dS(\zeta)}_{d(\text{gauge scalar})}.$$

(54)

We will use a **Lie transformation** $T(\epsilon)$.




Fig. 17. **Evolution in time t :** The solution of $\partial_t \bar{z} = V(\bar{z})$ [$\bar{z}(0) = z$] can be thought of as the Lie transformation $\bar{z}(t; z) = U(t)z$.

We will use a **Lie transformation** $T(\epsilon)$.




Fig. 17. **Evolution in time t** : The solution of $\partial_t \bar{z} = V(\bar{z})$ [$\bar{z}(0) = z$] can be thought of as the Lie transformation $\bar{z}(t; z) = U(t)z$.




Fig. 18. **“Evolution” in perturbation strength ϵ** : The solution of $\partial_\epsilon \bar{z} = g(\bar{z})$ [$\bar{z}(0) = z$] can be thought of as the Lie transformation $\bar{z}(\epsilon; z) = T(\epsilon)z$.

By analogy with time-dependent evolution, we use a **Lie transformation** in the perturbation parameter ϵ : $\bar{z} = T(\epsilon)z$, where

$$T(\epsilon) = e^{\epsilon L_g} \approx 1 + \epsilon L_g + \dots \quad (55)$$

Here

$$L_g \doteq \underbrace{g(z)}_{\text{generating function}} \frac{\partial}{\partial z}. \quad (56)$$

For use in perturbative calculations, Lie transforms can be compounded.

Consider the set of generating functions $\{g_n \mid n = 1, 2, \dots\}$ $[g_n = O(\epsilon^n)]$, and define $L_n \equiv L_{g_n}$. Then

$$T = e^{L_1} e^{L_2} e^{L_3} e^{L_4} \dots, \quad (57a)$$

$$T^{-1} = \dots e^{-L_4} e^{-L_3} e^{-L_2} e^{-L_1} \quad (57b)$$

$$\begin{aligned} &= 1 - L_1 + \left(-L_2 + \frac{1}{2}L_1^2\right) + \left(-L_3 + L_2L_1 - \frac{1}{6}L_1^3\right) \\ &\quad + \left(-L_4 + L_3L_1 + \frac{1}{2}L_2^2 - \frac{1}{2}L_2L_1^2 + \frac{1}{24}L_1^4\right) + \dots \end{aligned} \quad (57c)$$

We can use these results and the formula^a

$$L_g \gamma = g^\sigma \omega_{\sigma\nu} dz^\nu \quad (58)$$

to expand the basic law $\bar{\gamma} = T^{-1}\gamma + dS$ order by order.

^aThe differential geometers will recognize that there is a fundamental swindle here; please see me after class!

From $\bar{\gamma} = T^{-1}\gamma + dS$, we are thus led to
the equations of **one-form perturbation theory**:

$$\bar{\gamma}^{(-1)} = \gamma^{(-1)} + dS^{(-1)}, \quad (59a)$$

$$\begin{aligned} \bar{\gamma}^{(0)} &= \gamma^{(0)} + dS^{(0)} \\ &\quad - L_1 \gamma^{(-1)}, \end{aligned} \quad (59b)$$

$$\begin{aligned} \bar{\gamma}^{(1)} &= \gamma^{(1)} + dS^{(1)} \\ &\quad - L_1 \gamma^{(0)} + \left(\frac{1}{2}L_1^2 - L_2\right)\gamma^{(-1)}, \end{aligned} \quad (59c)$$

$$\begin{aligned} \bar{\gamma}^{(2)} &= \gamma^{(2)} + dS^{(2)} \\ &\quad - L_1 \gamma^{(1)} + \left(\frac{1}{2}L_1^2 - L_2\right)\gamma^{(0)} + \left(-L_3 + L_2L_1 - \frac{1}{6}L_1^3\right)\gamma^{(-1)}, \end{aligned} \quad (59d)$$

$$\begin{aligned} \bar{\gamma}^{(3)} &= \gamma^{(3)} + dS^{(3)} \\ &\quad - L_1 \gamma^{(2)} + \left(\frac{1}{2}L_1^2 - L_2\right)\gamma^{(1)} + \left(-L_3 + L_2L_1 - \frac{1}{6}L_1^3\right)\gamma^{(0)} \\ &\quad + \left(-L_4 + L_3L_1 + \frac{1}{2}L_2^2 - \frac{1}{2}L_2L_1^2 + \frac{1}{24}L_1^4\right)\gamma^{(-1)}. \end{aligned} \quad (59e)$$

$$\bar{\gamma}^{(4)} = \dots . \quad (59f)$$

The algebra is now straightforward, if tedious.

- At n th order, one needs to determine g_n^σ and $S^{(n)}$.
- The goal is to remove ζ dependence from $\bar{\gamma}_\sigma^{(n)}$.
- There is more than enough freedom to do this.

An example taken from the midst of the algebra:

$$\begin{aligned}\bar{\gamma}^{(2)}(\vec{Z}) = & (\dots) \cdot d\vec{x} + (\dots)dt + (\dots)dU + (\dots)d\mu \\ & + \left(f_\zeta - g_1^\mu + \frac{\partial S^{(2)}}{\partial \zeta} \right) d\zeta.\end{aligned}\tag{60}$$

Write $S = \langle S \rangle + \delta S$, such that $\langle \delta S \rangle = 0$. (S must be periodic in ζ in order to avoid secularities.) δg_1^μ was determined at $O(\epsilon)$. From

$$\langle \bar{\gamma}_\zeta^{(2)} \rangle = \langle f_\zeta \rangle - \langle g_1^\mu \rangle,\tag{61}$$

choose $\langle g_1^\mu \rangle$ to eliminate $\langle \bar{\gamma}_\zeta^{(2)} \rangle$. Now the condition $\delta \bar{\gamma}_\zeta^{(2)} = 0$ determines $\delta S^{(2)} = \int d\zeta (-\delta f_\zeta + \delta g_1^\mu)$.

Everything is now determined (through some chosen order).

- The procedure gives us the new ($\bar{\zeta}$ -independent) one-form $\bar{\gamma}$.
Through second (relative) order,

$$\begin{aligned}\bar{\gamma} \approx & [(q/c)\vec{A} + m\bar{U}\hat{\mathbf{b}} - \bar{\mu}\vec{K}^*] \cdot d\vec{x} + \bar{\mu} d\bar{\zeta} \\ & - \underbrace{\left(\frac{1}{2}m\bar{U}^2 + \bar{\mu}\omega_c + q\langle\varphi\rangle\right)}_{\text{gyroaveraged Hamiltonian}} dt,\end{aligned}\tag{62}$$

where

$$\vec{K}^* \doteq \vec{K} + \frac{1}{2}(\underbrace{\hat{\mathbf{b}} \cdot \vec{\nabla} \times \hat{\mathbf{b}}}_{\text{torsion}})\hat{\mathbf{b}}, \quad \vec{K} \doteq \underbrace{(\vec{\nabla}\hat{e}_1) \cdot \hat{e}_2}_{\text{gyrogauge vector}}.\tag{63}$$

Everything is now determined (through some chosen order).

- The procedure gives us the new ($\bar{\zeta}$ -independent) one-form $\bar{\gamma}$. Through second (relative) order,

$$\begin{aligned} \bar{\gamma} \approx & [(q/c)\vec{A} + m\bar{U}\hat{\mathbf{b}} - \bar{\mu}\vec{K}^*] \cdot d\vec{x} + \bar{\mu} d\bar{\zeta} \\ & - \underbrace{\left(\frac{1}{2}m\bar{U}^2 + \bar{\mu}\omega_c + q\langle\varphi\rangle\right)}_{\text{gyroaveraged Hamiltonian}} dt, \end{aligned} \quad (62)$$

where

$$\vec{K}^* \doteq \vec{K} + \frac{1}{2} \underbrace{(\hat{\mathbf{b}} \cdot \vec{\nabla} \times \hat{\mathbf{b}})}_{\text{torsion}} \hat{\mathbf{b}}, \quad \vec{K} \doteq \underbrace{(\vec{\nabla} \hat{e}_1) \cdot \hat{e}_2}_{\text{gyrogauge vector}}. \quad (63)$$

- From $\bar{\gamma}$, we can calculate the symplectic two-form $\bar{\omega}$; thus we obtain the equations of motion for the gyrocenter:

$$\bar{\omega}_{\sigma\nu} \frac{d\bar{z}^\nu}{dt} = 0. \quad (64)$$

Everything is now determined (through some chosen order).

- The procedure gives us the new ($\bar{\zeta}$ -independent) one-form $\bar{\gamma}$.
Through second (relative) order,

$$\begin{aligned} \bar{\gamma} \approx & [(q/c)\vec{A} + m\bar{U}\hat{\mathbf{b}} - \bar{\mu}\vec{K}^*] \cdot d\vec{x} + \bar{\mu} d\bar{\zeta} \\ & - \underbrace{\left(\frac{1}{2}m\bar{U}^2 + \bar{\mu}\omega_c + q\langle\varphi\rangle\right)}_{\text{gyroaveraged Hamiltonian}} dt, \end{aligned} \quad (62)$$

where

$$\vec{K}^* \doteq \vec{K} + \frac{1}{2} \underbrace{(\hat{\mathbf{b}} \cdot \vec{\nabla} \times \hat{\mathbf{b}})}_{\text{torsion}} \hat{\mathbf{b}}, \quad \vec{K} \doteq \underbrace{(\vec{\nabla} \hat{e}_1) \cdot \hat{e}_2}_{\text{gyrogauge vector}}. \quad (63)$$

- From $\bar{\gamma}$, we can calculate the symplectic two-form $\bar{\omega}$; thus we obtain the equations of motion for the gyrocenter:

$$\bar{\omega}_{\sigma\nu} \frac{d\bar{z}^\nu}{dt} = 0. \quad (64)$$

- And the g 's give us $T(\epsilon)$, i.e., the asymptotic expansion of the proper ($\bar{\mu}$ -conserving) gyrocenter variables: $\bar{\vec{Z}} = T(\epsilon)\vec{Z}$.

From the equations of motion,
we obtain the (collisionless) **gyrokinetic equation**.



The Vlasov equation has the covariant form

$$\frac{\partial f(\vec{z}, t)}{\partial t} + \dot{z}^\nu \frac{\partial f}{\partial z^\nu} = 0. \quad (65)$$

From the equations of motion, we obtain the (collisionless) gyrokinetic equation.

- The Vlasov equation has the covariant form

$$\frac{\partial f(\vec{z}, t)}{\partial t} + \dot{z}^\nu \frac{\partial f}{\partial z^\nu} = 0. \quad (65)$$

- Thus it retains its form in any system of coordinates:

$$\frac{\partial \tilde{F}(\vec{Z}, t)}{\partial t} + \dot{Z}^\nu \frac{\partial \tilde{F}}{\partial Z^\nu} = 0, \quad (66)$$

$$\text{or } \frac{\partial \tilde{F}(\vec{X}, \bar{U}, \bar{\mu}, \bar{\zeta}, t)}{\partial t} + \dot{\vec{X}} \cdot \vec{\nabla} \tilde{F} + \dot{\bar{U}} \frac{\partial \tilde{F}}{\partial \bar{U}} + \underbrace{\dot{\bar{\mu}}}_0 \frac{\partial \tilde{F}}{\partial \bar{\mu}} + \dot{\bar{\zeta}} \frac{\partial \tilde{F}}{\partial \bar{\zeta}} = 0. \quad (67)$$

From the equations of motion, we obtain the (collisionless) gyrokinetic equation.

- The Vlasov equation has the covariant form

$$\frac{\partial f(\vec{z}, t)}{\partial t} + \dot{z}^\nu \frac{\partial f}{\partial z^\nu} = 0. \quad (65)$$

- Thus it retains its form in any system of coordinates:

$$\frac{\partial \tilde{F}(\vec{Z}, t)}{\partial t} + \dot{\vec{Z}}^\nu \frac{\partial \tilde{F}}{\partial \vec{Z}^\nu} = 0, \quad (66)$$

$$\text{or } \frac{\partial \tilde{F}(\vec{X}, \bar{U}, \bar{\mu}, \bar{\zeta}, t)}{\partial t} + \dot{\vec{X}} \cdot \bar{\nabla} \tilde{F} + \dot{\bar{U}} \frac{\partial \tilde{F}}{\partial \bar{U}} + \underbrace{\dot{\bar{\mu}}}_{0} \frac{\partial \tilde{F}}{\partial \bar{\mu}} + \dot{\bar{\zeta}} \frac{\partial \tilde{F}}{\partial \bar{\zeta}} = 0. \quad (67)$$

- We can trivially average this over $\bar{\zeta}$, defining $\bar{F} \doteq \langle \tilde{F} \rangle_{\bar{\zeta}}$:

$$\frac{\partial \bar{F}}{\partial t} + \dot{\vec{X}} \cdot \bar{\nabla} \bar{F} + \dot{\bar{U}} \frac{\partial \bar{F}}{\partial \bar{U}} = 0. \quad (68)$$

This is what is usually called the (collisionless) gyrokinetic equation.

Summary of Systematology, Part I:

The gyrokinetic equations of motion

● Lagrange's variational principle: $\delta \int \gamma = 0$.

Summary of Systematology, Part I:

The gyrokinetic equations of motion

- Lagrange's variational principle: $\delta \int \gamma = 0$.
- Write the one-form γ in any convenient set of variables.

Summary of Systematology, Part I:

The gyrokinetic equations of motion

- **Lagrange's variational principle:** $\delta \int \gamma = 0$.
- Write the one-form γ in any convenient set of variables.
- Isolate gyrophase (ζ) dependence by introducing the **noncanonical variables** $\vec{Z} \doteq \{\vec{x}, U, \mu, \zeta\}$.

Summary of Systematology, Part I:

The gyrokinetic equations of motion

- **Lagrange's variational principle:** $\delta \int \gamma = 0$.
- Write the one-form γ in any convenient set of variables.
- Isolate gyrophase (ζ) dependence by introducing the **noncanonical variables** $\vec{Z} \doteq \{\vec{x}, U, \mu, \zeta\}$.
- Use **Lie perturbation theory** to remove ζ dependence from all components of γ by changing variables to $\vec{\bar{Z}} \doteq \{\vec{\bar{x}}, \bar{U}, \bar{\mu}, \bar{\zeta}\}$ and demanding that $\bar{\gamma}(\vec{\bar{Z}}) = \bar{\gamma}(\dots, \cancel{\zeta}, \dots)$.

Summary of Systematology, Part I:

The gyrokinetic equations of motion

- **Lagrange's variational principle:** $\delta \int \gamma = 0$.
- Write the one-form γ in any convenient set of variables.
- Isolate gyrophase (ζ) dependence by introducing the **noncanonical variables** $\vec{Z} \doteq \{\vec{x}, U, \mu, \zeta\}$.
- Use **Lie perturbation theory** to remove ζ dependence from all components of γ by changing variables to $\vec{\bar{Z}} \doteq \{\vec{\bar{x}}, \bar{U}, \bar{\mu}, \bar{\zeta}\}$ and demanding that $\bar{\gamma}(\vec{\bar{Z}}) = \bar{\gamma}(\dots, \cancel{\zeta}, \dots)$.
- By **Noether's theorem**, $\bar{\mu}$ is conserved.

Summary of Systematology, Part I:

The gyrokinetic equations of motion

- **Lagrange's variational principle:** $\delta \int \gamma = 0$.
- Write the one-form γ in any convenient set of variables.
- Isolate gyrophase (ζ) dependence by introducing the **noncanonical variables** $\vec{Z} \doteq \{\vec{x}, U, \mu, \zeta\}$.
- Use **Lie perturbation theory** to remove ζ dependence from all components of γ by changing variables to $\vec{\bar{Z}} \doteq \{\vec{\bar{x}}, \bar{U}, \bar{\mu}, \bar{\zeta}\}$ and demanding that $\bar{\gamma}(\vec{\bar{Z}}) = \bar{\gamma}(\dots, \cancel{\zeta}, \dots)$.
- By **Noether's theorem**, $\bar{\mu}$ is conserved.
- Use the characteristic gyrocenter equations of motion to write the **gyrokinetic equation** for the gyrocenter distribution \bar{F} .

Table of Contents

- ✓ ■ Introduction: Heuristic Gyrokinetics
- ✓ ■ Systematic Gyrokinetics, Part I: The Gyrokinetic Equations of Motion
- Systematic Gyrokinetics, Part II: The Gyrokinetic Poisson Equation
- The Current Status of Gyrokinetics

Systematic Gyrokinetics, Part II: The Gyrokinetic Poisson Equation

**To obtain the gyrokinetic Poisson equation,
we must express the charge density of the particles
in terms of the gyrocenter distribution.**

Following Dubin, Krommes, Oberman, and Lee (1983), we must consider various distribution functions:

$f(\vec{z}, t)$ — particle PDF (69a)

$\tilde{F}(\vec{Z}, t)$ — particle PDF in lowest-order gyrocenter coordinates (69b)

$\tilde{\overline{F}}(\vec{Z}, t)$ — particle PDF in “true” gyrocenter coordinates (69c)

$\overline{F}(\vec{Z}, t)$ — gyroaveraged PDF $\doteq \langle \tilde{\overline{F}} \rangle_{\vec{\zeta}}$. (69d)

**To obtain the gyrokinetic Poisson equation,
we must express the charge density of the particles
in terms of the gyrocenter distribution.**

Following Dubin, Krommes, Oberman, and Lee (1983), we must consider various distribution functions:

$$f(\vec{z}, t) \text{ — particle PDF} \quad (69a)$$

$$\tilde{F}(\vec{Z}, t) \text{ — particle PDF in lowest-order gyrocenter coordinates} \quad (69b)$$

$$\tilde{\overline{F}}(\vec{Z}, t) \text{ — particle PDF in “true” gyrocenter coordinates} \quad (69c)$$

$$\overline{F}(\vec{Z}, t) \text{ — gyroaveraged PDF} \doteq \langle \tilde{\overline{F}} \rangle_{\vec{\zeta}}. \quad (69d)$$

Poisson’s equation is naturally written in the particle (laboratory) coordinate system:

$$-\nabla^2 \varphi(\vec{x}, t) = 4\pi \sum_s (\overline{n}q)_s \int d\vec{v}' \underbrace{f_s(\vec{x}, \vec{v}', t)}_{\substack{\text{must obtain} \\ \text{from } \tilde{\overline{F}}}}. \quad (70)$$

Now

$$I \doteq \int d\vec{v}' f(\vec{x}, \vec{v}', t) = \underbrace{\int d\vec{z}'}_{\text{integral over complete phase space}} \delta(\vec{x} - \vec{x}') f(\vec{z}', t) \quad (71a)$$

$$= \int J(\vec{Z}) d\vec{Z} \delta(\vec{x} - \vec{x}'(\vec{Z})) \tilde{F}(\vec{Z}, t), \quad (71b)$$

where the Jacobian is

$$J \doteq \frac{\partial(\vec{z}')}{\partial(\vec{Z})}. \quad (72)$$

Now recall the pull-back transformation

$$\tilde{F} = T\tilde{\tilde{F}}. \quad (73)$$

Therefore

$$I = \int \bar{J} d\bar{\vec{Z}} \delta(\vec{x} - \vec{x}'(\bar{\vec{Z}})) \textcolor{red}{T} \underbrace{\tilde{\tilde{F}}(\bar{\vec{Z}}, t)}_{\text{still depends on } \zeta}. \quad (74)$$

[This is still formally exact (assuming $\bar{\mu}$ is conserved).]

Use the pull-back transformation $T(\epsilon)$
to close Poisson's equation
in terms of the gyrocenter distribution \tilde{F} .

$$\frac{\partial \tilde{F}}{\partial t} + \dot{\vec{X}} \cdot \vec{\nabla} \tilde{F} + \dot{v}_{\parallel} \frac{\partial \tilde{F}}{\partial v_{\parallel}} + \underbrace{\dot{\mu}}_0 \frac{\partial \tilde{F}}{\partial \mu} + \dot{\zeta} \frac{\partial \tilde{F}}{\partial \zeta} = -C[\tilde{F}], \quad (75)$$

$$-\nabla^2 \varphi = 4\pi \sum_s (\bar{n}q)_s \int \bar{J} d\bar{\vec{Z}} \delta(\vec{x} - \vec{x}'(\bar{\vec{Z}})) \underbrace{T}_{\text{pull-back}} \tilde{F}(\bar{\vec{Z}}, t). \quad (76)$$

Important insight (Dubin, Krommes, Oberman, and Lee, 1983):

- By construction, $\dot{\vec{X}}$, \dot{v}_{\parallel} , and $\dot{\zeta}$ are all independent of $\bar{\zeta}$.
- Therefore, $\bar{\zeta}$ dependence enters only from
 - initial conditions,
 - collisional effects.
- For *collisionless* theory, the kinetic equation does not couple the evolution of $\bar{F} \doteq \langle \tilde{F} \rangle_{\bar{\zeta}}$ and $\delta \bar{F} \doteq \tilde{F} - \langle \tilde{F} \rangle_{\bar{\zeta}}$.

For collisionless theory, we make the **GK closure**:

$$\tilde{\bar{F}}(\bar{\vec{z}}, t) \approx \underbrace{\bar{F}(\bar{\vec{Z}}, t)}_{\text{obeys GKE}}. \quad (77)$$

$$\partial_t \bar{F} + \dots = 0.$$

If we write $T = 1 + \delta T$, then

$$I \approx \int \bar{J} d\bar{\vec{Z}} \delta(\vec{x} - \vec{x}'(\bar{\vec{Z}})) \left(\underbrace{1}_{\text{gyrocenter density}} + \underbrace{\delta T}_{\text{polarization density}} \right) \bar{F}(\bar{\vec{Z}}, t). \quad (78)$$

For collisionless theory, we make the **GK closure**:

$$\tilde{\overline{F}}(\overline{\vec{z}}, t) \approx \underbrace{\overline{F}(\overline{\vec{Z}}, t)}_{\text{obeys GKE}}. \quad (77)$$

$$\partial_t \overline{F} + \dots = 0.$$

If we write $T = 1 + \delta T$, then

$$I \approx \int \overline{J} d\overline{\vec{Z}} \delta(\vec{x} - \vec{x}'(\overline{\vec{Z}})) \left(\underbrace{1}_{\text{gyrocenter density}} + \underbrace{\delta T}_{\text{polarization density}} \right) \overline{F}(\overline{\vec{Z}}, t). \quad (78)$$

- The polarization density represents the difference between the particle motion and the gyrocenter motion.

For collisionless theory, we make the **GK closure**:

$$\tilde{\overline{F}}(\overline{\vec{z}}, t) \approx \underbrace{\overline{F}(\overline{\vec{Z}}, t)}_{\text{obeys GKE}}. \quad (77)$$

$$\partial_t \overline{F} + \dots = 0.$$

If we write $T = 1 + \delta T$, then

$$I \approx \int \overline{J} d\overline{\vec{Z}} \delta(\vec{x} - \vec{x}'(\overline{\vec{Z}})) \left(\underbrace{1}_{\text{gyrocenter density}} + \underbrace{\delta T}_{\text{polarization density}} \right) \overline{F}(\overline{\vec{Z}}, t). \quad (78)$$

- The polarization density represents the difference between the particle motion and the gyrocenter motion.
- It leads to a generalization of the GK dielectric constant ϵ^G for $T_i \neq 0$ and $k_\perp \rho_i = O(1)$.

For collisionless theory, we make the **GK closure**:

$$\tilde{\bar{F}}(\bar{\vec{z}}, t) \approx \underbrace{\bar{F}(\bar{\vec{Z}}, t)}_{\text{obeys GKE}}. \quad (77)$$

$$\partial_t \bar{F} + \dots = 0.$$

If we write $T = 1 + \delta T$, then

$$I \approx \int \bar{J} d\bar{\vec{Z}} \delta(\vec{x} - \vec{x}'(\bar{\vec{Z}})) \left(\underbrace{1}_{\text{gyrocenter density}} + \underbrace{\delta T}_{\text{polarization density}} \right) \bar{F}(\bar{\vec{Z}}, t). \quad (78)$$

- The polarization density represents the difference between the particle motion and the gyrocenter motion.
- It leads to a generalization of the GK dielectric constant ϵ^G for $T_i \neq 0$ and $k_\perp \rho_i = O(1)$.
- δT is complicated; it must be truncated at some order in ϵ !

The coupled gyrokinetic-Poisson equations comprise
a new nonlinear dynamical system
with unique properties.

The gyrokinetic nonlinear dynamical system
describes an ensemble of gyrocenters
placed into a *gyrokinetic vacuum*
possessing large dielectric permittivity ϵ^G .
They move with the $\vec{E} \times \vec{B}$ and magnetic drifts.

The coupled gyrokinetic-Poisson equations comprise
a new nonlinear dynamical system
with unique properties.

The gyrokinetic nonlinear dynamical system
describes an ensemble of gyrocenters
placed into a *gyrokinetic vacuum*
possessing large dielectric permittivity ϵ^G .
They move with the $\vec{E} \times \vec{B}$ and magnetic drifts.

This dynamical system . . .

The coupled gyrokinetic-Poisson equations comprise a new nonlinear dynamical system with unique properties.

The gyrokinetic nonlinear dynamical system describes an ensemble of gyrocenters placed into a *gyrokinetic vacuum* possessing large dielectric permittivity ϵ^G . They move with the $\vec{E} \times \vec{B}$ and magnetic drifts.

This dynamical system . . .

- has a particular **energy constant** (for first-order truncation; can be reduced in a particular limit to the energy invariant of the Charney–Hasegawa–Mima equation);

The coupled gyrokinetic-Poisson equations comprise a new nonlinear dynamical system with unique properties.

The gyrokinetic nonlinear dynamical system describes an ensemble of gyrocenters placed into a *gyrokinetic vacuum* possessing large dielectric permittivity ϵ^G . They move with the $\vec{E} \times \vec{B}$ and magnetic drifts.

This dynamical system . . .

- has a particular **energy constant** (for first-order truncation; can be reduced in a particular limit to the energy invariant of the Charney–Hasegawa–Mima equation);
- has a **lower level of thermal discreteness noise** (because of shielding due to ion polarization).

The coupled gyrokinetic-Poisson equations comprise a new nonlinear dynamical system with unique properties.

The gyrokinetic nonlinear dynamical system describes an ensemble of gyrocenters placed into a *gyrokinetic vacuum* possessing large dielectric permittivity ϵ^G . They move with the $\vec{E} \times \vec{B}$ and magnetic drifts.

This dynamical system . . .

- has a particular **energy constant** (for first-order truncation; can be reduced in a particular limit to the energy invariant of the Charney–Hasegawa–Mima equation);
- has a **lower level of thermal discreteness noise** (because of shielding due to ion polarization).
- is **gyrogauge-invariant** (physical quantities like $\bar{\mu}$ are independent of the choice of perpendicular unit vectors \hat{e}_1 and \hat{e}_2).

Energetic consistency is best achieved by working with a Lagrangian field theory.

To get a workable theory, the transformation operator $T(\epsilon)$ must be truncated at some order in ϵ . Truncations are used in two places:

Energetic consistency is best achieved by working with a Lagrangian field theory.

To get a workable theory, the transformation operator $T(\epsilon)$ must be truncated at some order in ϵ . Truncations are used in two places:

● GK equation (\vec{X} and \vec{v}_{\parallel});

Energetic consistency is best achieved by working with a **Lagrangian field theory**.

To get a workable theory, the transformation operator $T(\epsilon)$ must be truncated at some order in ϵ . Truncations are used in two places:

- GK equation (\vec{X} and \vec{v}_{\parallel});
- GK Poisson equation (form of the polarization density).

This must be done consistently.

Energetic consistency is best achieved by working with a **Lagrangian field theory**.

To get a workable theory, the transformation operator $T(\epsilon)$ must be truncated at some order in ϵ . Truncations are used in two places:

- GK equation (\vec{X} and \vec{v}_{\parallel});
- GK Poisson equation (form of the polarization density).

This must be done consistently. It is best to derive both equations from one **field-theoretic variational principle** (Sugama, 2000; Brizard, 2000; Scott, unpublished):

$$S[\overline{F}, \varphi]$$

GK equation


GK Poisson equation

Energetic consistency is best achieved by working with a **Lagrangian field theory**.

To get a workable theory, the transformation operator $T(\epsilon)$ must be truncated at some order in ϵ . Truncations are used in two places:

- GK equation (\vec{X} and \vec{v}_{\parallel});
- GK Poisson equation (form of the polarization density).

This must be done consistently. It is best to derive both equations from one **field-theoretic variational principle** (Sugama, 2000; Brizard, 2000; Scott, unpublished):




Energetic consistency is best achieved by working with a **Lagrangian field theory**.

To get a workable theory, the transformation operator $T(\epsilon)$ must be truncated at some order in ϵ . Truncations are used in two places:

- GK equation (\vec{X} and $\vec{v}_{||}$);
- GK Poisson equation (form of the polarization density).

This must be done consistently. It is best to derive both equations from one **field-theoretic variational principle** (Sugama, 2000; Brizard, 2000; Scott, unpublished):




Energetic consistency is best achieved by working with a **Lagrangian field theory**.

To get a workable theory, the transformation operator $T(\epsilon)$ must be truncated at some order in ϵ . Truncations are used in two places:

- GK equation ($\vec{\bar{X}}$ and $\vec{v}_{||}$);
- GK Poisson equation (form of the polarization density).

This must be done consistently. It is best to derive both equations from one **field-theoretic variational principle** (Sugama, 2000; Brizard, 2000; Scott, unpublished):



**A form of Noether's theorem guarantees that
any approximation to S
leads to an energetically consistent theory!**

Summary of Systematology, Part II:

The gyrokinetic Poisson equation

- The particle PDF can be transformed to various variables.

Summary of Systematology, Part II:

The gyrokinetic Poisson equation

- The particle PDF can be transformed to various variables.
- In particular, $\tilde{F} = T\tilde{\tilde{F}}$ (pull-back transformation).

Summary of Systematology, Part II:

The gyrokinetic Poisson equation

- The particle PDF can be transformed to various variables.
- In particular, $\tilde{F} = T\tilde{\tilde{F}}$ (pull-back transformation).
- Make the gyrokinetic closure $\tilde{F} \approx \overline{F}$, where $\overline{F} \doteq \langle \tilde{\tilde{F}} \rangle_{\zeta}$.

Summary of Systematology, Part II:

The gyrokinetic Poisson equation

- The particle PDF can be transformed to various variables.
- In particular, $\tilde{F} = T\tilde{\tilde{F}}$ (pull-back transformation).
- Make the gyrokinetic closure $\tilde{F} \approx \bar{F}$, where $\bar{F} \doteq \langle \tilde{\tilde{F}} \rangle_{\zeta}$.
- Truncate T to some order in ϵ [usually $O(\epsilon)$] (being sure to not lose energy conservation).

Summary of Systematology, Part II:

The gyrokinetic Poisson equation

- The particle PDF can be transformed to various variables.
- In particular, $\tilde{F} = T\tilde{\tilde{F}}$ (pull-back transformation).
- Make the **gyrokinetic closure** $\tilde{\tilde{F}} \approx \overline{F}$, where $\overline{F} \doteq \langle \tilde{\tilde{F}} \rangle_{\zeta}$.
- Truncate T to some order in ϵ [usually $O(\epsilon)$] (being sure to not lose energy conservation).
- Alternatively, derive both the GKE and the GK Poisson equation from a **single variational principle** employing an approximate Lagrangian (thereby *guaranteeing* energetic consistency).

Table of Contents

- ✓ ■ Introduction: Heuristic Gyrokinetics
- ✓ ■ Systematic Gyrokinetics, Part I: The Gyrokinetic Equations of Motion
- ✓ ■ Systematic Gyrokinetics, Part II: The Gyrokinetic Poisson Equation
- The Current Status of Gyrokinetics

The Current Status of Gyrokinetics

Gyrokinetics is used extensively for studies of low-frequency microturbulence in both fusion and astrophysical contexts.

- Analytical theory of drift waves and related modes:
 - linear theory in general magnetic geometry is well developed;
 - wave (gyro)kinetic equations for weak turbulence are also studied.

Gyrokinetics is used extensively for studies of low-frequency microturbulence in both fusion and astrophysical contexts.

- Analytical theory of drift waves and related modes:
 - linear theory in general magnetic geometry is well developed;
 - wave (gyro)kinetic equations for weak turbulence are also studied.
- Derive nonlinear **gyrofluid equations** (e.g., HME) by taking moments of the GKE (plus implementing important *Landau-fluid closure* to capture effects of wave–particle resonances):
 - works well linearly;
 - has problems with nonlinear wave–wave–particle interactions.

Gyrokinetics is used extensively for studies of low-frequency microturbulence in both fusion and astrophysical contexts.

- Analytical theory of drift waves and related modes:
 - linear theory in general magnetic geometry is well developed;
 - wave (gyro)kinetic equations for weak turbulence are also studied.
- Derive nonlinear **gyrofluid equations** (e.g., HME) by taking moments of the GKE (plus implementing important *Landau-fluid closure* to capture effects of wave–particle resonances):
 - works well linearly;
 - has problems with nonlinear wave–wave–particle interactions.
- Simulation techniques:
 - particle-in-cell (PIC) / Monte Carlo sampling
 - “continuum” or “Vlasov” approach (direct solution of the 5D GKE; see, for example, the GYRO code of Candy & Waltz)

There remain some serious concerns.

- In the PIC approach to solution of the GKE, **Monte Carlo sampling noise** may sometimes be an issue [see review article of Krommes (2007)].

There remain some serious concerns.

- In the PIC approach to solution of the GKE, **Monte Carlo sampling noise** may sometimes be an issue [see review article of Krommes (2007)].
- $\bar{\mu}$ is not always conserved.
 - There is an analog to KAM theory (particle motion is Hamiltonian).
 - Nonlinear interaction between gyration and slower degrees of freedom gives rise to stochastic regions ($\Rightarrow \bar{\mu}$ is merely an “**adiabatic invariant**”).

There remain some serious concerns.

- In the PIC approach to solution of the GKE, **Monte Carlo sampling noise** may sometimes be an issue [see review article of Krommes (2007)].
- $\bar{\mu}$ is not always conserved.
 - There is an analog to KAM theory (particle motion is Hamiltonian).
 - Nonlinear interaction between gyration and slower degrees of freedom gives rise to stochastic regions ($\Rightarrow \bar{\mu}$ is merely an “adiabatic invariant”).
- **Is gyrokinetics ill-posed** for general 3D magnetic fields (torsional or stochastic), as has recently been asserted? **NO** (but subtle: anholonomic frame fields, global coordinate systems, etc.).

There remain some serious concerns.

- In the PIC approach to solution of the GKE, **Monte Carlo sampling noise** may sometimes be an issue [see review article of Krommes (2007)].
- $\bar{\mu}$ is not always conserved.
 - There is an analog to KAM theory (particle motion is Hamiltonian).
 - Nonlinear interaction between gyration and slower degrees of freedom gives rise to stochastic regions ($\Rightarrow \bar{\mu}$ is merely an “adiabatic invariant”).
- **Is gyrokinetics ill-posed** for general 3D magnetic fields (torsional or stochastic), as has recently been asserted? **NO** (but subtle: anholonomic frame fields, global coordinate systems, etc.).
- Is the conventional gyrokinetic closure adequate for the study of **long-wavelength phenomena** such as
 - macroscopic electric fields and plasma rotation,
 - zonal flows?

Perhaps not (Parra & Catto, 2008, 2009)...

Physics on the transport time scale: GK momentum conservation, rotation, etc.

- Gyrokinetics is good for studies of low-frequency microturbulence on the nonlinear saturation time scale (\ll plasma confinement time).
- What about transport-time-scale physics?
 - profile relaxation
 - radial diffusion of momentum
- There might be interchanges of limits ($\epsilon \rightarrow 0$; $t \rightarrow \infty$), inconsistent truncations [GKE vs GK Poisson; $k_{\perp} \rho_i \rightarrow 0$ although originally derived for $k_{\perp} \rho_i = O(1)$], etc.

Following F. Parra (2009, unpublished), consider a slab limit with constant \vec{B} (as did Dubin et al., 1983):

x — “radial” direction (of profile gradients)

y — “poloidal” direction

z — direction of \vec{B} .

Standard gyrokinetic truncation may introduce a **spurious momentum source**.

Let $\overline{Q}(x) \doteq (L_y L_z)^{-1} \int_0^{L_y} dy \int_0^{L_z} dz Q(x, y, z)$ denote the “flux-surface average.” Then an *exact* moment of the Vlasov equation leads to

$$\frac{\partial(n_i m_i \overline{u}_{i,y})}{\partial t} = -\frac{\partial}{\partial x} \left(\underbrace{\overline{\pi}_{xy}}_{\substack{\text{off-diagonal} \\ \text{component of} \\ \text{stress tensor}}} \right). \quad (79)$$

Here

$$\overline{\pi}_{xy} \sim n_i m_i \overline{V_{E,x} V_{E,y}} \quad (\text{Reynolds stresses}). \quad (80)$$

Parra’s result is that when slab gyrokinetics is truncated to $O(\epsilon^2)$,

$$\frac{\partial(n_i m_i \overline{u}_{i,y})}{\partial t} = -\frac{\partial \overline{\pi}_{xy}}{\partial x} + \underbrace{(\text{spurious momentum source})}_{O(\epsilon^3), \text{ but nonconservative!}}. \quad (81)$$

(There are also further problems with collisions and recovery of neoclassical theory.) **Final resolutions still pending...**

Summary

- Gyrokinetics is the key formalism for the study of low-frequency microturbulence in magnetized plasmas.
- It has an elegant and systematic development in terms of Hamiltonian / Lagrangian formalism and Lie perturbation theory.
- It has enjoyed many practical successes (see next-week's talk by Ron Waltz).
- There are some interesting outstanding problems.
 - The magnetic moment $\bar{\mu}$ is not always conserved.
 - Currently, there is some confusion about the role and proper expression of momentum conservation in gyrokinetics.

Gyrokinetics has had an interesting intellectual development; we should pay tribute to the pioneers.

Some key players on gyrokinetic *theory*:

- Catto
- Frieman & Chen
- Lee
- Dubin et al.
- Hahm
- Brizard
- Qin

More recently,

- Sugama
- Scott

...and especially **Robert Littlejohn**.

**Companion talk on
the application of nonlinear gyrokinetics
to magnetic fusion:**

**“Gyrokinetic Simulation of
Turbulent Transport in Fusion Plasmas”**

R. E. Waltz

Tuesday morning, Aug. 4