A. De Pasquale<sup>1,2</sup>, P. Facchi<sup>3,4</sup>, V. Giovannetti<sup>2</sup>, G. Parisi<sup>6,7,8</sup>, S. Pascazio<sup>1,4</sup>, A. Scardicchio<sup>9,10</sup>

<sup>1</sup>Dipartimento di Fisica and MECENAS, Università di Bari, I-70126 Bari, Italy <sup>2</sup>NEST, Scuola Normale Superiore and Istituto Nanoscienze-CNR, Piazza dei Cavalieri 7, I-56126 Pisa, Italy

<sup>3</sup>Dipartimento di Matematica and MECENAS, Università di Bari, I-70125 Bari, Italy <sup>4</sup>INFN, Sezione di Bari, I-70126 Bari, Italy

<sup>6</sup>Dipartimento di Fisica, Università di Roma "La Sapienza", 00185 Roma, Italy <sup>7</sup>Centre for Statistical Mechanics and Complexity (SMC), CNR-INFM, 00185 Roma, Italy

<sup>8</sup>INFN, Sezione di Roma, 00185 Roma, Italy

<sup>9</sup>Abdus Salam International Center for Theoretical Physics, I-34014 Trieste, Italy <sup>10</sup>INFN, Sezione di Trieste, I-34014 Trieste, Italy

**Abstract.** The local purity of large many-body quantum systems can be studied by following a statistical mechanical approach based on a random matrix model. Restricting the analysis to the case of global pure states, this method proved to be successful and a full characterization of the statistical properties of the local purity was obtained by computing the partition function of the problem. Here we generalize these techniques to the case of global mixed states. Since the computation of the partition function is far more challenging than in the pure case, we focus on the computation of the first moments of the local purity. Finally, we establish a connection with the theory of twirling maps in quantum channels.

PACS numbers: 03.67.Mn, 03.65.Ud, 89.75.-k, 03.67.-a

In quantum mechanics the purity of a quantum state measures the amount of indeterminacy which originates either from lack of information on the process that led to its preparation, or from the correlations that have been established with some external party (environment). Mathematically speaking the two mechanisms are strongly interconnected, as the former can always be represented as an instance of the latter via purification [1, 2]. For composite quantum system one can distinguish between global purity (i.e. the purity of the system as a whole), and the local (or "marginal") purities (i.e the purities associated with the subsystems that compose it). Clarifying the connection between the global and local purities of a quantum state is an important problem of quantum information theory [3, 4, 5] which is closely related to the characterization of bipartite entanglement [6, 7]. In particular, for pure global states, the local purity of a subsystem provides a direct measure of the bipartite entanglement between the two parts: the smaller the purity, the larger the bipartite entanglement. For mixed (non pure) global states instead, the connection between local purity and entanglement is more subtle: no direct relation between the two quantities exists and bipartite entanglement measures for the global system can be obtained only by taking proper convex roof extensions of the local purity (the average being computed over all convex decompositions of the initial global state). Still, studying how the global indeterminacy of a composite system affects the indeterminacy of its constituents raises important theoretical questions which call for a thoughtful investigation [3, 4, 5].

The distribution of the local purity of pure global states of large composite systems, was studied in Refs. [8, 9] by using statistical mechanical methods. In particular a partition function for a suitable canonical ensemble was introduced and the problem was translated in terms of the distribution of the eigenvalues of random matrices. Here we endeavor to generalize the above procedure to a system in a global mixed state. In such a case, however, the direct computation of the partition function is far more complicated, since the average over the unitary group does not factorize. On the contrary, a direct characterization of the first moments associated with the local purity distributions, appears to be feasible. Therefore, in an attempt to collect some preliminary insight on the statistical properties of the problem, we focus on this latter issue.

This paper is organized as follows. In Sec. 2 we introduce the notation and set the basis of the statistical mechanical approach to the problem, starting from the simpler case of pure states and generalizing it to the case of generic mixed states. In Sec. 3 we consider the high temperature expansion of the partition function and compute the first moments of the purity, by making use of Zuber's solution of some basic integrals over the unitary group [10]. In Sec. 4 we establish an interesting connection between our problem and the theory of quantum channels. In particular, we present an alternative approach for computing the moments, that exploits the properties of twirling maps. We conclude in Sec. 5 by summarizing our findings and discussing them in terms of future perspectives.

# 2. The statistical approach: partition function

In this section we review the statistical approach introduced in Ref. [8] for studying the local mixedness properties of pure states and discuss its generalization to the case of mixed states.

Consider a bipartite system X = AB made up of two subsystems A and B, described by the Hilbert space  $\mathcal{H}_X = \mathcal{H}_A \otimes \mathcal{H}_B$ , with  $\dim \mathcal{H}_A = N_A$ ,  $\dim \mathcal{H}_B = N_B$ and  $\dim \mathcal{H}_X = N = N_A N_B$ . Without loss of generality we will assume that  $N_A \leq N_B$ . The states of X are represented by the set  $\mathfrak{S}(\mathcal{H}_X)$  of nonnegative unit-trace operators (density matrices) on the Hilbert space  $\mathcal{H}_X$ . The purity of such states, defined as

$$\pi_{AB}(\rho) := \operatorname{Tr} \rho^2 \in [1/N, 1],$$
(1)

for each  $\rho \in \mathfrak{S}(\mathcal{H}_X)$ , provides a characterization of the global mixing of the system and induces a partition of  $\mathfrak{S}(\mathcal{H}_X)$  into a collection of distinct subsets  $\mathfrak{S}_x(\mathcal{H}_X) := \{\rho \in \mathfrak{S}(\mathcal{H}_X) : \operatorname{Tr} \rho^2 = x\}$ . The minimum value of x = 1/N is attained when X is in the completely mixed state I/N, whereas the maximum x = 1 is attained over the set  $\mathfrak{S}_1(\mathcal{H}_X)$  consisting of all pure states  $|\psi\rangle_X$ . For each  $\rho \in \mathfrak{S}(\mathcal{H}_X)$  we can also define its *A-local* and *B-local* purity functions as

$$\pi_A(\rho) := \operatorname{Tr} \rho_A^2 , \qquad \pi_B(\rho) := \operatorname{Tr} \rho_B^2 , \qquad (2)$$

with  $\rho_A = \text{Tr}_B \rho$  and  $\rho_B = \text{Tr}_A \rho$  being the reduced density matrices of the subsystem A and B, respectively.

#### 2.1. Total system in a pure state

On the special set  $\mathfrak{S}_1(\mathcal{H}_X)$  of pure states  $\rho = |\psi\rangle_X \langle \psi|$  of X, the A and the B-local purities coincide

$$\pi(\psi) := \pi_A(|\psi\rangle_X\langle\psi|) = \pi_B(|\psi\rangle_X\langle\psi|) , \qquad (3)$$

and provide a measure of the bipartite entanglement between A and B: the smaller  $\pi(\psi)$ , the larger the entanglement contained in  $|\psi\rangle_X\langle\psi|$ . The statistical distribution of  $\pi(\psi)$  on  $\mathfrak{S}_1(\mathcal{H}_X)$  has been studied in Refs. [8, 9]. This was done by introducing the partition function

$$\mathcal{Z}(\beta) = \int d\mu(\psi) \, e^{-\beta \, \pi(\psi)} \,, \tag{4}$$

where the local purity  $\pi(\psi)$  of  $|\psi\rangle_X$  plays the role of an effective energy of the system,  $\beta$ is a Lagrange multiplier that fixes the value of the purity/energy and selects an isopurity manifold [11], and  $d\mu(\psi)$  is a (normalized) measure on the space of pure states  $\mathfrak{S}_1(\mathcal{H}_X)$ . The natural choice for the latter is induced by the Haar (probability) measure  $d\mu_H(U)$ on the unitary group  $\mathcal{U}(\mathcal{H}_X) \simeq \mathcal{U}(N)$ , through the mapping

$$|\psi\rangle_X := U_X |\psi_0\rangle_X , \qquad (5)$$

with  $|\psi_0\rangle_X$  an arbitrary reference unit vector of  $\mathcal{H}_X$ . Thus the partition function becomes

$$\mathcal{Z}(\beta) = \int \mathrm{d}\mu_H(U_X) \exp\left(-\beta \operatorname{Tr}(\operatorname{Tr}_B(U_X|\psi_0\rangle_X\langle\psi_0|U_X^{\dagger})^2)\right).$$
(6)

Then, by noticing that for every  $U_X \in \mathcal{U}(N)$  the reduced density matrix  $\rho_A = \text{Tr}_B(|\psi\rangle_X\langle\psi|)$  can be written as  $\rho_A = U_A \Lambda_A U_A^{\dagger}$ , with  $U_A \in \mathcal{U}(N_A)$  and  $\Lambda_A = \text{diag}\{\lambda_{A,1}, \lambda_{A,2}, \ldots, \lambda_{A,N_A}\}$ , the expression (6) becomes

$$\mathcal{Z}(\beta) = \int d\mu_H(U_A) \int d\sigma(\Lambda_A) \, e^{-\beta \operatorname{Tr} \Lambda_A^2} = \int d\sigma(\Lambda_A) \, e^{-\beta \operatorname{Tr} \Lambda_A^2},\tag{7}$$

where we exploited the fact that the measure induced over the density matrices  $\rho_A$ by  $d\mu_H(U_X)$  factorizes into the product of a measure over the unitary group  $d\mu_H(U_A)$ (related to the eigenvectors of  $\rho_A$ ) and a measure  $d\sigma(\Lambda_A)$  over the  $(N_A - 1)$ -dimensional simplex of its eigenvalues  $\lambda_{A,j}$  [12, 13]. In particular, it can be shown that [15, 16, 17]

$$d\sigma(\Lambda_A) = C_{N_A,N_B} \delta\left(1 - \sum_{1 \le i \le N_A} \lambda_{A,i}\right) \prod_{1 \le j \le N_A} \theta(\lambda_{A,j}) \lambda_{A,j}^{N_B - N_A} \times \prod_{1 \le l < m \le N_A} (\lambda_{A,l} - \lambda_{A,m})^2 d^{N_A} \lambda_A ,$$
(8)

with

$$C_{N_A,N_B} = \frac{\Gamma(N_A N_B)}{\prod_{0 \le j \le N_A - 1} \Gamma(N_B - j) \Gamma(N_A - j + 1)},\tag{9}$$

and  $\theta(x)$  and  $\Gamma$  being the unit step and the Euler gamma function, respectively. The derivatives of  $\mathcal{Z}(\beta)$ , evaluated for  $\beta = 0$ , yield the moments of  $\pi(\psi)$  with respect to the measure  $d\mu(\psi)$ , i.e.

$$\mathcal{M}_{n} := \left\langle \left( \mathrm{Tr} \rho_{A}^{2} \right)^{n} \right\rangle_{\mathfrak{S}_{1}(\mathcal{H}_{X})} = \int \mathrm{d}\mu(\psi) \ \pi(\psi)^{n} = (-1)^{n} \left. \frac{\partial^{n} \mathcal{Z}(\beta)}{\partial \beta^{n}} \right|_{\beta=0} \ . \tag{10}$$

These functions fully determine the statistical distribution of  $\pi(\psi)$  on  $\mathfrak{S}_1(\mathcal{H}_X)$  and, in the high temperature regime, provide an expansion of  $\mathcal{Z}(\beta)$ . More generally, in analogy to what is commonly done in statistical physics, one can also define the moments of  $\pi(\psi)$  for  $\beta \neq 0$  as

$$\mathcal{M}_{n}(\beta) := \left\langle \left( \mathrm{Tr} \rho_{A}^{2} \right)^{n} \right\rangle_{\beta,\mathfrak{S}_{1}(\mathcal{H}_{X})} = \int \mathrm{d}\mu_{\beta}(\psi) \ \pi(\psi)^{n} = \frac{(-1)^{n}}{\mathcal{Z}(\beta)} \ \frac{\partial^{n} \mathcal{Z}(\beta)}{\partial \beta^{n}} \ , \ (11)$$

with  $d\mu_{\beta}(\psi)$  being the canonical measure

$$d\mu_{\beta}(\psi) := d\mu(\psi) \frac{e^{-\beta \pi(\psi)}}{\mathcal{Z}(\beta)} .$$
(12)

The latter is a deformation of the Haar measure  $d\mu(\psi)$  obtained by including a nonuniform weight which explicitly depends upon the local purity, through  $\beta$ . In particular, as  $\beta$  increases  $d\mu_{\beta}(\psi)$  enhances the role of the states with lower values of  $\pi(\psi)$  (i.e. larger values of bipartite entanglement), to the extent that for  $\beta \to +\infty$  only the maximally entangled elements of  $\mathfrak{S}_1(\mathcal{H}_X)$  contribute to the values (11). Since  $\pi(\psi)$  is bounded above, one can also consider negative temperatures, where the role of less entangled states is enhanced, and when  $\beta \to -\infty$  only the separable elements of  $\mathfrak{S}_1(\mathcal{H}_X)$ contributes to the values (11). Consequently, across different ranges of temperatures, the moments in Eq. (11) characterize the statistical distribution of the local purity of X computed with respect to a *canonical* ensemble whose constituents are selected according to an effective thermal distribution characterized by the parameter  $\beta$ .

In the limit of large N, the  $\beta$ -dependence of the statistics of the local purity  $\pi(\psi)$ (and hence of the bipartite entanglement of the system) was characterized in Ref. [8, 9] by identifying the class of states which maximize the distributions (12), i.e. typical states with respect to the canonical measure (12). In this context it was shown that the system undergoes two main phase transitions, related to different distributions of the eigenvalues  $\Lambda_A$  of the typical states: a second-order phase transition, mentioned above, associated to a  $\mathbb{Z}_2$  symmetry breaking, and related to the vanishing of some eigenvalues of  $\rho_A$ , followed by a first-order phase transition, associated to the evaporation of the largest eigenvalue from the sea of the others.

# 2.2. Total system in a mixed state

A natural question is what happens when the global system X is in a *mixed* state  $\rho$  of purity x < 1, rather than in a pure state. A generalization of Eq. (4) is obtained by replacing  $\pi(\psi)$  with (say) the A-local purity  $\pi_A(\rho)$  of Eq. (2) and the measure  $d\mu(\psi)$ with a proper measure  $d\mu_x(\rho)$  on the set  $\mathfrak{S}_x(\mathcal{H}_X)$ . This yields the following definition of the partition function of the A-local purity

$$\mathcal{Z}_A(x,\beta) = \int \mathrm{d}\mu_x(\rho) \,\mathrm{e}^{-\beta \,\pi_A(\rho)} = C_x \int \mathrm{d}\mu(\rho) \,\delta(\mathrm{Tr}\,\rho^2 - x) \,\mathrm{e}^{-\beta \,\pi_A(\rho)} \,, \quad (13)$$

where  $d\mu(\rho)$  is a probability measure on the set of mixed states (see below), and  $C_x := \left[\int d\mu(\rho) \,\delta(\operatorname{Tr} \rho^2 - x)\right]^{-1}$  a normalization factor. An analogous expression for the *B*-local purity partition function  $\mathcal{Z}_B(x,\beta)$  is obtained by replacing  $\pi_A(\rho)$  with  $\pi_B(\rho)$  in Eq. (13): notice, however, that, at variance with the case analyzed in Ref. [8, 9], for x < 1 the partition function  $\mathcal{Z}_B(x,\beta)$  will in general differ form  $\mathcal{Z}_A(x,\beta)$ .

It is worth stressing that the function  $\mathcal{Z}_A(x,\beta)$  provides only statistical information on the local mixedness of X, but not directly on its bipartite entanglement properties: this is due to the fact that for generic mixed states  $\rho$  of X the local purities  $\pi_A(\rho)$ and  $\pi_B(\rho)$  are not entanglement measures. A generalization of Eq. (4) that retains the ability of characterizing the statistical properties of the bipartite entanglement of X for x < 1 could in principle be constructed by replacing  $\pi(\psi)$  with the convex-roof counterpart of  $\pi_A(\rho)$ , namely  $\bar{\pi}_A(\rho) = \max_{\mathcal{E}} \sum_j p_j \pi_A(\psi_j)$ , where the maximum is taken over all ensembles  $\mathcal{E} := \{p_j, |\psi_j\rangle_X\}_j$  which yield a convex decomposition of the mixed state  $\rho$  (see e.g. Refs. [19, 21]). The quantity  $\bar{\pi}_A(\rho)$  is a proper measure of the bipartite entanglement, but the resulting partition function does not allow for a simple analytic treatment and will not be discussed in the present paper. Finally, since there is no unique measure on mixed states [22], we need to properly specify the choice of  $d\mu(\rho)$  which enters in Eq. (13) – the consistency requirement that for pure states Eq. (13) should reduce back to Eq. (4) (i.e.  $\mathcal{Z}_A(1,\beta) = \mathcal{Z}(\beta)$ ) does not eliminate such ambiguity. Indeed, as previously discussed for  $\rho_A$ , the Hermitian matrix  $\rho$  can always be diagonalized by a unitary operation, and as consequence we can write the measure  $d\mu(\rho)$  as the product of a measure on the (N-1)-dimensional symplex of the eigenvalues and a measure on the unitary group  $\mathcal{U}(N)$  related to the eigenvectors. However, if on the one hand it would be natural to take the Haar measure on  $\mathcal{U}(N)$ so that  $d\mu(\rho) = d\mu(U_X\rho U_X^{\dagger})$ , on the other hand the measure on the eigenvalues can be chosen in different possible ways [23]. In order to overcome this ambiguity we will use a balanced purification strategy. Let us introduce hence the composite Hilbert space  $\mathcal{H}_{XX'} = \mathcal{H}_X \otimes \mathcal{H}_{X'}$ , where  $\mathcal{H}_X \simeq \mathcal{H}_{X'}$  are isomorphic. In this  $N^2$ -dimensional Hilbert space, each  $\rho$  of X can be represented by those pure states  $|\Psi\rangle_{XX'}$  which provide a purification for such density matrix, i.e. which satisfy the identity

$$\rho = \operatorname{Tr}_{X'}(|\Psi\rangle_{XX'}\langle\Psi|) . \tag{14}$$

Thanks to this identification we can now induce a measure on  $\mathfrak{S}(\mathcal{H}_X)$  by sampling the pure states on  $\mathcal{H}_{XX'}$  according to the unique, unitarily invariant Haar measure which, as usual, is induced by the Haar measure on the unitary group  $\mathcal{U}(N^2)$  through the mapping  $|\Psi\rangle_{XX'} := U_{XX'}|\Psi_0\rangle_{XX'}$ , where  $|\Psi_0\rangle_{XX'}$  is an arbitrary reference vector and  $U_{XX'} \in \mathcal{U}(N^2)$ . With this choice the partition function becomes

$$\mathcal{Z}_A(x,\beta) = C_x \int d\mu_H(U_{XX'}) \,\delta\left(x - \operatorname{Tr}(\operatorname{Tr}_{X'}(|\Psi\rangle_{XX'}\langle\Psi|)^2) \,\mathrm{e}^{-\beta \operatorname{Tr}\left((\operatorname{Tr}_B(\operatorname{Tr}_{X'}|\Psi\rangle_{XX'}\langle\Psi|))^2\right)},\tag{15}$$

where we used the fact that  $\rho_A = \text{Tr}_B \rho = \text{Tr}_B(\text{Tr}_{X'}|\Psi\rangle_{XX'}\langle\Psi|)$ . Analogously to what we have seen for the pure case, x = 1, by writing  $\rho = U_X \Lambda_X U_X^{\dagger}$  with  $\Lambda_X = \text{diag}(\lambda_1, \lambda_2, ..., \lambda_N)$ , we get

$$\mathcal{Z}_A(x,\beta) = C_x \int \mathrm{d}\mu_H(U_X) \int \mathrm{d}\sigma(\Lambda_X) \,\delta\left(x - \operatorname{Tr}\Lambda_X^2\right) \,\mathrm{e}^{-\beta \operatorname{Tr}\left((\operatorname{Tr}_B(U_X\Lambda_X U_X^{\dagger}))^2\right)}, \quad (16)$$

where  $d\mu_H(U_X)$  is the Haar measure on  $\mathcal{U}(N)$  and

$$d\sigma(\Lambda_X) = C_N \,\delta\left(1 - \sum_{1 \le i \le N} \lambda_i\right) \prod_{1 \le i \le N} \theta(\lambda_i) \prod_{1 \le i < j \le N} (\lambda_i - \lambda_j)^2 d^N \lambda, \quad (17)$$

with

$$C_N = \frac{\Gamma(N^2)}{\Gamma(N+1) \prod_{1 \le k \le N} \Gamma(k)^2}.$$
(18)

Therefore, we have identified the measure  $d\mu_x(\rho)$  of Eq. (13) with

$$d\mu_x(\rho) = C_x d\mu_H(U_X) d\sigma(\Lambda_X) \delta\left(x - \operatorname{Tr} \Lambda_X^2\right).$$
(19)

Notice that in the case of pure states, i.e. x = 1, the density operator of the system reduces to  $\rho = |\psi\rangle_X \langle \psi|$ , where  $|\psi\rangle_X = U_X |\psi_0\rangle_X$ ,  $|\psi_0\rangle_X$  being an arbitrary reference state (see Eq. (5)), and the matrix  $\Lambda_X$  becomes a rank one projection. Thus the expression (16) reduces to (6), namely

$$\mathcal{Z}_A(1,\beta) = \mathcal{Z}(\beta) . \tag{20}$$

#### 2.3. Asymptotic behavior and analysis of moments

For x < 1, the integration over the unitary group  $\mathcal{U}(N)$  in Eq. (16) does not factorize, making the computation of the partition function far more complicated than for the case of a pure state (7). The only notable exception is the case of maximally mixed states (i.e. x = 1/N), when the Dirac delta in Eq. (16) selects a unique diagonal matrix  $\Lambda_X$  (the totally mixed state of X). This makes the exponent equal to  $e^{-\beta/N_A}$  for all  $U_X$ and yields the following exact expression

$$\mathcal{Z}_A(1/N,\beta) = \mathrm{e}^{-\beta/N_A} \,. \tag{21}$$

Otherwise, for intermediate values of the purity, 1/N < x < 1, the situation is much more complicated. Still, as we will show in the following, at small  $\beta$  the evaluation of the moments  $\mathcal{M}_n^A(x,\beta)$  associated with  $\mathcal{Z}_A(x,\beta)$  admits an exact analytical treatment. The latter are formally defined as

$$\mathcal{M}_{n}^{A}(x,\beta) := \int \mathrm{d}\mu_{x,\beta}(\rho) \ \pi_{A}^{n}(\rho) = \frac{(-1)^{n}}{\mathcal{Z}_{A}(x,\beta)} \ \frac{\partial^{n} \mathcal{Z}_{A}(x,\beta)}{\partial \beta^{n}} \ , \tag{22}$$

and represent the average value of  $\pi_A^n(\rho)$  with the canonical measure

$$d\mu_{x,\beta}(\rho) := d\mu_x(\rho) \; \frac{e^{-\beta \; \pi_A(\rho)}}{\mathcal{Z}_A(x,\beta)} \;, \tag{23}$$

with  $d\mu_x(\rho)$  given by Eq. (19). For pure states (x = 1) the  $\mathcal{M}_n^A(x,\beta)$  coincide with the moments  $\mathcal{M}_n(\beta)$  defined in Eq. (11): at  $\beta = 0$ , in the large N limit the expression for such quantities has been computed in Ref. [8], while the exact expressions for first five of them can be found in Ref. [24]. In the case of a totally mixed state (x = 1/N)Eq. (21) yields instead values which are independent of the temperature  $\beta$ , namely

$$\mathcal{M}_{n}^{A}(1/N,\beta) = \mathcal{M}_{n}^{A}(1/N,0) = N_{A}^{-n} .$$
(24)

For intermediate values of x, by expanding Eq. (22) up to the first order in  $\beta$ , we get

$$\mathcal{M}_n^A(x,\beta) \sim \mathcal{M}_n^A(x,0) - \beta \left[ \mathcal{M}_{n+1}^A(x,0) - \mathcal{M}_1^A(x,0) \mathcal{M}_n^A(x,0) \right], \qquad \beta \to 0.$$
(25)

Incidentally, notice that in agreement with Eq. (24), the  $\beta$ -corrections of Eq. (25) vanish when x = 1/N. The above expression shows that, at least in the high temperature regime, we can focus on the unbiased moments  $\mathcal{M}_n^A(x, 0)$ .

#### **3.** Moments of the purity at $\beta = 0$

#### 3.1. First moment

In this section we compute the exact first moment of the purity  $\mathcal{M}_1^A(x,0)$ , by making use of Zuber's solution of some basic integrals over the unitary group [10]. In particular, we will show that the only dependence on the spectrum of the density matrix of the global system is in terms of its purity x, whose value is fixed in the partition function (13).

Let us fix the spectrum of the global density matrix  $\rho$  of the system:

$$\Lambda_X = \operatorname{diag}(\lambda_{N_B(\alpha-1)+\beta}),\tag{26}$$

with  $\alpha = 1, \ldots, N_A$  and  $\beta = 1, \ldots, N_B$ . A purification of  $\Lambda_X$  in the space  $\mathcal{H}_{XX'}$ , with X = AB and X' = A'B' is

$$|\Psi\rangle_{XX'} = \sum_{1 \le \alpha \le N_A} \sum_{1 \le \beta \le N_B} \sqrt{\lambda_{\alpha\beta}} \ |\alpha\beta\rangle_{AB} \otimes |\alpha\beta\rangle_{A'B'}, \tag{27}$$

where we have set, for simplicity,  $\lambda_{\alpha\beta} = \lambda_{N_B(\alpha-1)+\beta}$  and  $|\alpha\beta\rangle_{AB} = |\alpha\rangle_A \otimes |\beta\rangle_B$ ,  $\{|\alpha\rangle_A\}$ and  $\{|\beta\rangle_B\}$  ( $\{|\alpha\rangle_{A'}\}$  and  $\{|\beta\rangle_{B'}\}$ ) being the reference basis in  $\mathcal{H}_A$  and  $\mathcal{H}_B$  ( $\mathcal{H}_{A'}$  and  $\mathcal{H}_{B'}$ ), respectively. The set of vectors in  $\mathcal{H}_{XX'}$  with the same Schmidt coefficients is given by  $U_{XX'}|\Psi\rangle_{XX'}$ , where  $U_{XX'} = U_X \otimes U_{X'}$ , with  $U_X, U_{X'} \in \mathcal{U}(N)$ , and yields the set of density matrices with the same spectrum  $\Lambda_X$ , namely  $\rho = U_X \Lambda_X U_X^{\dagger}$ . By partial tracing over subsystem B one obtains the set of reduced density matrices  $\rho_A(U) = \operatorname{Tr}_B(\operatorname{Tr}_{X'}(U_{XX'}|\Psi\rangle_{XX'}\langle\Psi|U_{XX'}^{\dagger}))$ . Notice that this expression does not depend on  $U_{X'} \in \mathcal{U}(N)$ :

$$\rho_{A} = \sum_{1 \le \alpha \le N_{A}} \sum_{1 \le \beta \le N_{B}} \lambda_{\alpha\beta} \operatorname{Tr}_{B}(U_{AB} | \alpha\beta\rangle_{AB} \langle \alpha\beta | U_{AB}^{\dagger}) \\
= \sum_{1 \le \alpha \le N_{A}} \sum_{1 \le \beta, j \le N_{B}} \lambda_{\alpha\beta} {}_{B} \langle j | U_{AB} | \alpha\beta\rangle_{AB} \langle \alpha\beta | U_{AB}^{\dagger} | j\rangle_{B} ,$$
(28)

The purity is given by

$$\pi_{A}(U_{X}\Lambda_{X}U_{X}^{\dagger}) = \operatorname{Tr} \rho_{A}^{2} = \sum_{1 \leq \alpha_{1}, \alpha_{2} \leq N_{A}} \sum_{1 \leq \beta_{1}, \beta_{2} \leq N_{B}} \sum_{1 \leq j_{1}, j_{2} \leq N_{B}} \lambda_{\alpha_{1}\beta_{1}}\lambda_{\alpha_{2}\beta_{2}} \times_{AB} \langle \alpha_{2}\beta_{2} | U_{AB}^{\dagger} | j_{2} \rangle_{B} \langle j_{1} | U_{AB} | \alpha_{1}\beta_{1} \rangle_{AB} \times_{AB} \langle \alpha_{1}\beta_{1} | U_{AB}^{\dagger} | j_{1} \rangle_{B} \langle j_{2} | U_{AB} | \alpha_{2}\beta_{2} \rangle_{AB} , \qquad (29)$$

which, by the completeness relation for subsystem A, becomes

$$\pi_{A}(U_{X}\Lambda_{X}U_{X}^{\dagger}) = \sum_{1 \leq \alpha_{1}, \alpha_{2} \leq N_{A}} \sum_{1 \leq \beta_{1}, \beta_{2} \leq N_{B}} \sum_{1 \leq i_{1}, i_{2} \leq N_{A}} \sum_{1 \leq j_{1}, j_{2} \leq N_{B}} \lambda_{\alpha_{1}\beta_{1}}\lambda_{\alpha_{2}\beta_{2}}$$
$$\times_{AB} \langle \alpha_{2}\beta_{2} | U_{AB}^{\dagger} | i_{1}j_{2} \rangle_{AB} \langle i_{1}j_{1} | U_{AB} | \alpha_{1}\beta_{1} \rangle_{AB}$$
$$\times_{AB} \langle \alpha_{1}\beta_{1} | U_{AB}^{\dagger} | i_{2}j_{1} \rangle_{AB} \langle i_{2}j_{2} | U_{AB} | \alpha_{2}\beta_{2} \rangle_{AB}.$$
(30)

Let us now compute the first moment of the purity (22) at  $\beta = 0$ . By recalling that  $\mathcal{Z}_A(x,0) = 1$ ,  $d\mu_{x,0}(\rho) = d\mu_x(\rho)$ , and  $d\mu_x(\rho) = C_x d\mu_H(U_X) d\sigma(\Lambda_X) \delta\left(x - \operatorname{Tr} \Lambda_X^2\right)$ , we get

$$\mathcal{M}_{n}^{A}(x,0) = \int \mathrm{d}\mu_{x}(\rho) \ \pi_{A}^{n}(\rho)$$
$$= C_{x} \int \mathrm{d}\sigma(\Lambda_{X})\delta(x - \operatorname{Tr}\Lambda_{X}^{2}) \ \mathcal{M}_{n}^{A}(\Lambda_{X}) =: \langle \mathcal{M}_{n}^{A}(\Lambda_{X}) \rangle_{x}, \tag{31}$$

where

$$\mathcal{M}_{n}^{A}(\Lambda_{X}) := \int \mathrm{d}\mu_{H}(U_{X}) \ \pi_{A}^{n}(U_{X}\Lambda_{X}U_{X}^{\dagger}).$$
(32)

From (30) the average over the unitary group of the first moment particularizes to

$$\mathcal{M}_{1}^{A}(\Lambda_{X}) = \sum_{1 \leq \alpha_{1}, \alpha_{2} \leq N_{A}} \sum_{1 \leq \beta_{1}, \beta_{2} \leq N_{B}} \lambda_{\alpha_{1}\beta_{1}}\lambda_{\alpha_{2}\beta_{2}} \\ \times \sum_{1 \leq i_{1}, i_{2} \leq N_{A}} \sum_{1 \leq j_{1}, j_{2} \leq N_{B}} \int d\mu_{H}(U_{X})U_{i_{1}j_{1},\alpha_{1}\beta_{1}}U_{i_{2}j_{2},\alpha_{2}\beta_{2}}U_{\alpha_{2}\beta_{2},i_{1}j_{2}}^{\dagger}U_{\alpha_{1}\beta_{1},i_{2}j_{1}}^{\dagger}, \quad (33)$$

where  $U_{ij,\alpha\beta} = {}_{AB} \langle ij | U_{AB} | \alpha\beta \rangle_{AB}$ . This integral can be explicitly computed by using Zuber's solution [10]:

$$\int \mathrm{d}\mu_H(U) U_{i_1 j_1} \dots U_{i_n j_n}(U) U_{k_1 l_1}^{\dagger} \dots U_{k_n l_n}^{\dagger} = \sum_{\tau, \sigma \in S_n} C[\sigma] \prod_{1 \le a \le n} \delta(i_a, \ell_{\tau(a)}) \delta(j_a, k_{\tau\sigma(a)}), (34)$$

with

$$C[\sigma] = \sum_{|Y|=n} \frac{(\chi^{(k)}(1))^2 \chi^{(k)}([\sigma])}{n!^2 s_k(I)},$$
(35)

where  $C[\sigma]$  is the sum over the Young diagrams Y of the character  $\chi^{(k)}([\sigma])$  of the symmetric group  $S_n$  associated to Y, depending on the conjugacy class  $[\sigma]$  of the permutation  $\sigma$ ,  $s_k(I)$  is the dimension of the representation, and  $\delta(a, b)$  is the Kronecker delta. Applying this solution to (33) we get

$$\mathcal{M}_{1}^{A}(\Lambda_{X}) = \sum_{1 \leq \alpha_{1}, \alpha_{2} \leq N_{A}} \sum_{1 \leq \beta_{1}, \beta_{2} \leq N_{B}} \lambda_{\alpha_{1}\beta_{1}}\lambda_{\alpha_{2}\beta_{2}} \\ \times \sum_{\tau, \sigma \in S_{2}} C[\sigma]f_{1}(\tau)\delta(\alpha_{1}\beta_{1}, \alpha_{\tau\sigma(2)}\beta_{\tau\sigma(2)})\delta(\alpha_{2}\beta_{2}, \alpha_{\tau\sigma(1)}\beta_{\tau\sigma(1)}) \\ = \sum_{1 \leq \alpha_{1}, \alpha_{2} \leq N_{A}} \sum_{1 \leq \beta_{1}, \beta_{2} \leq N_{B}} \sum_{\tau, \sigma \in S_{2}} \sum_{c \in \mathcal{C}(S_{2})} C[\sigma]f_{1}(\tau)\delta([\tau\sigma s], c)\lambda_{\alpha_{c(1)}\beta_{c(1)}}\lambda_{\alpha_{c(2)}\beta_{c(2)}},$$

$$(36)$$

where  $f_1(\pi)$  depends on the permutation  $\pi \in S_2$ 

$$f_1(\pi) := \sum_{1 \le i_1, i_2 \le N_A} \delta(i_1, i_{\pi(1)}) \delta(i_2, i_{\pi(2)}) \sum_{1 \le j_1, j_2 \le N_B} \delta(j_1, j_{\pi(2)}) \delta(j_2, j_{\pi(1)}), \quad (37)$$

 $s \in S_2$  is the transposition (swapping) of pairs of nearby indices ([s] = [2])

$$i_{s(1)} = i_2$$
 and  $i_{s(2)} = i_1$ , (38)

and  $\mathcal{C}(S_2) = \{[1^2], [2]\}\)$  is the set of the conjugacy classes of the symmetric group  $S_2$ . From (36) it can be easily inferred that the only possible contributions of the spectrum are related to the conjugacy classes of the symmetric group  $S_2$ :

$$[\tau\sigma s] = [1^2] \Rightarrow \left(\sum_{1 \le \alpha \le N_A} \sum_{1 \le \beta \le N_B} \lambda_{\alpha\beta}\right)^2 = 1,$$
  
$$[\tau\sigma s] = [2] \Rightarrow \left(\sum_{1 \le \alpha \le N_A} \sum_{1 \le \beta \le N_B} \lambda_{\alpha\beta}^2\right) = \operatorname{Tr} \Lambda_X^2.$$
(39)

By summing and by using the explicit expressions of the coefficients (35) [10]

$$C[1^2] = \frac{1}{(N-1)(N+1)}, \qquad C[2] = -\frac{1}{(N-1)N(N+1)}, \tag{40}$$

we get

$$\mathcal{M}_{1}^{A}(\Lambda_{X}) = \frac{N_{A}(N_{B}^{2}-1)}{N_{A}^{2}N_{B}^{2}-1} + \frac{N_{B}(N_{A}^{2}-1)}{N_{A}^{2}N_{B}^{2}-1} \operatorname{Tr} \Lambda_{X}^{2}.$$
(41)

The first moment of the purity of subsystem A, is the average of (41) over the spectrum of the system. By plugging (41) into (31), we finally get

$$\mathcal{M}_{1}^{A}(x,0) = \frac{N_{A}(N_{B}^{2}-1)}{N_{A}^{2}N_{B}^{2}-1} + \frac{N_{B}(N_{A}^{2}-1)}{N_{A}^{2}N_{B}^{2}-1}x.$$
(42)

Some special cases:- It is worth noticing that for a balanced bipartition  $N_A = N_B = \sqrt{N} \gg 1$  Eq. (42) yields

$$\mathcal{M}_{1}^{A}(x,0) = \frac{\sqrt{N}(1+x)}{N+1} \sim \frac{1+x}{\sqrt{N}} \,. \tag{43}$$

At x = 1 (i.e. pure global states), Eq. (43) coincides with that obtained in Refs. [8, 24]. Finally, consider the case in which  $\rho$  is maximally mixed, i.e. is the density matrix I/N. In this case x = 1/N and Eq. (42) gives

$$\mathcal{M}_{1}^{A}(1/N,0) = \frac{1}{N_{A}}, \qquad (44)$$

in agreement with the general result (24).

# 3.2. k-th moment

The technique shown in the previous section can be easily generalized in order to compute from (32) higher moments at  $\beta = 0$ . We get

$$\mathcal{M}_{k}^{A}(\Lambda_{X}) = \int d\mu_{H}(U_{X}) \pi_{A}^{k}(U_{X}\Lambda_{X}U_{X}^{\dagger})$$

$$= \sum_{1 \leq \alpha_{1}, \dots, \alpha_{2k} \leq N_{A}} \sum_{1 \leq \beta_{1}, \dots, \beta_{2k} \leq N_{B}} \prod_{1 \leq i \leq 2k} \lambda_{\alpha_{i}\beta_{i}}$$

$$\times \sum_{1 \leq i_{1}, \dots, i_{2k} \leq N_{A}} \sum_{1 \leq j_{1}, \dots, j_{2k} \leq N_{B}} \int d\mu_{H}(U_{X}) \prod_{1 \leq \ell \leq 2k} U_{i_{\ell}j_{\ell}, \alpha_{\ell}\beta_{\ell}}$$

$$\times \prod_{1 \leq m \leq k} U_{\alpha_{2m}\beta_{2m}, i_{2m-1}j_{2m}}^{\dagger} U_{\alpha_{2m-1}\beta_{2m-1}, i_{2m}j_{2m-1}}^{\dagger}.$$
(45)

Equation (34) for n = 2k gives

$$\mathcal{M}_{k}^{A}(\Lambda_{X}) = \sum_{1 \leq \alpha_{1}, \dots, \alpha_{2k} \leq N_{A}} \sum_{1 \leq \beta_{1}, \dots, \beta_{2k} \leq N_{B}} \sum_{\tau, \sigma \in S_{2k}} \sum_{c \in \mathcal{C}(S_{2k})} C[\sigma] f_{k}(\tau) \delta([\tau \sigma s], c) \prod_{1 \leq i \leq 2k} \lambda_{\alpha_{c(i)}\beta_{c(i)}},$$

$$(46)$$

where 
$$f_k(\pi)$$
 depends on the permutation  $\pi \in S_{2k}$   
 $f_k(\pi) := \sum_{1 \le i_1, \dots, i_{2k} \le N_A} \sum_{1 \le j_1, \dots, j_{2k} \le N_B} \prod_{1 \le \ell \le 2k} \delta(i_\ell, i_{\pi(\ell)}) \prod_{1 \le m \le k} \delta(j_{2m-1}, j_{\pi(2m)}) \delta(j_{2m}, j_{\pi(2m-1)})$ 
(47)

and, analogously to Eq. (38), s is the swapping of pairs of nearby indices

$$i_{s(2\ell-1)} = i_{2\ell}$$
 and  $i_{s(2\ell)} = i_{2\ell-1} \quad \forall \ell = 1, \dots k.$  (48)

Observe that when k = 1 we retrieve  $\mathcal{M}_1^A(\Lambda_X)$  (see Eq. (36)). The different contributions of the spectrum can be classified in terms of the the conjugacy classes of the symmetric group, as shown in Eq. (39). However, for k > 1, they do not depend only upon the purity  $x = \operatorname{Tr} \Lambda_X^2$ , but exhibit a more complex dependence on the spectrum, through its higher order invariants  $\operatorname{Tr} \Lambda_X^k$ , with k > 2. Thus the integral on the spectrum (31) is in general non trivial.

#### 3.3. Second moment

Now let us fully compute the second moment of the purity for arbitrary bipartite states, with purity  $x \in [1/N, 1]$ , generalizing some results found for pure states, x = 1, [8, 14, 24]. The second moment can be directly computed by setting k = 2 in Eq. (45). The expression for the coefficients  $C[\pi]$  in (34), when  $\pi \in \mathcal{C}(S_4)$  is [10]

$$C \left[1^{4}\right] = \frac{N^{4} - 8N^{2} + 6}{(N-3)(N-2)(N-1)N^{2}(N+1)(N+2)(N+3)}$$

$$C \left[2, 1^{2}\right] = -\frac{1}{(N-3)(N-1)N(N+1)(N+3)}$$

$$C \left[2^{2}\right] = \frac{N^{2} + 6}{(N-3)(N-2)(N-1)N^{2}(N+1)(N+2)(N+3)}$$

$$C \left[3, 1\right] = \frac{2N^{2} - 3}{(N-3)(N-2)(N-1)N^{2}(N+1)(N+2)(N+3)}$$

$$C \left[4\right] = -\frac{5}{(N-3)(N-2)(N-1)N(N+1)(N+2)(N+3)}.$$
(49)

The symmetric group  $S_4$ , consists of five conjugacy classes, that yield the following contributions to the integral (46) in terms of the spectrum of  $\rho$ :

$$\begin{aligned} [\tau\sigma s] &= \begin{bmatrix} 1^4 \end{bmatrix} \quad \Rightarrow \left( \sum_{1 \le \alpha \le N_A} \sum_{1 \le \beta \le N_B} \lambda_{\alpha\beta} \right)^4 = 1 \\ [\tau\sigma s] &= \begin{bmatrix} 2, 1^2 \end{bmatrix} \Rightarrow \left( \sum_{1 \le \alpha_1 \le N_A} \sum_{1 \le \beta_1 \le N_B} \lambda_{\alpha_1\beta_1}^2 \right) \left( \sum_{1 \le \alpha_2 \le N_A} \sum_{1 \le \beta_2 \le N_B} \lambda_{\alpha_2\beta_2} \right)^2 = \operatorname{Tr} \Lambda_X^2 \\ [\tau\sigma s] &= \begin{bmatrix} 2^2 \end{bmatrix} \quad \Rightarrow \left( \sum_{1 \le \alpha \le N_A} \sum_{1 \le \beta \le N_B} \lambda_{\alpha\beta}^2 \right)^2 = (\operatorname{Tr} \Lambda_X^2)^2 \\ [\tau\sigma s] &= \begin{bmatrix} 3, 1 \end{bmatrix} \quad \Rightarrow \left( \sum_{1 \le \alpha_1 \le N_A} \sum_{1 \le \beta_1 \le N_B} \lambda_{\alpha_1\beta_1}^3 \right) \left( \sum_{1 \le \alpha_2 \le N_A} \sum_{1 \le \beta_2 \le N_B} \lambda_{\alpha_2\beta_2} \right) = \operatorname{Tr} \Lambda_X^3 \end{aligned}$$

$$[\tau\sigma s] = [4] \qquad \Rightarrow \sum_{1 \le \alpha \le N_A} \sum_{1 \le \beta \le N_B} \lambda_{\alpha\beta}^4 = \operatorname{Tr} \Lambda_X^4, \tag{50}$$

with  $\tau, \sigma \in S_4$  and  $s \in S_2$  defined in (48) being k = 2. By gathering all we get

$$\begin{split} \mathcal{M}_{2}^{A}(\Lambda_{X}) &= \left\{ \begin{array}{c} \left[ C[1^{4}]N_{A}^{2}N_{B}^{4} + C[2,1^{2}]2N_{A}\left(N_{A}^{2}+2\right)N_{B}^{3} \\ &+ C[2^{2}]N_{A}^{2}\left(N_{A}^{2}+2\right)N_{B}^{2} + C[3,1]8N_{A}^{2}N_{B}^{2} \\ &+ C[4]2N_{A}\left(2N_{A}^{2}+1\right)N_{B} \right] \\ &+ \operatorname{Tr}\Lambda_{X}^{2} \left[ C[1^{4}]2N_{A}\left(N_{A}^{2}+2\right)N_{B}^{3} \\ &+ C[2,1^{2}]2N_{A}N_{B}\left(N_{A}^{2}\left(N_{B}^{2}+4\right)+2\left(N_{B}^{2}+1\right)\right) \\ &+ C[3,1]8N_{A}N_{B}\left(N_{A}^{2}\left(N_{B}^{2}+2\right)+2N_{B}^{2}+1\right) \\ &+ C[4]4N_{A}^{2}\left(N_{A}^{2}+8\right)N_{B}^{2} \right] \\ &+ (\operatorname{Tr}\Lambda_{X}^{2})^{2} \left[ C[1^{4}]N_{A}^{2}\left(N_{A}^{2}+2\right)N_{B}^{3} \\ &+ C[2,1^{2}]2N_{A}N_{B}\left(N_{A}^{2}\left(N_{B}^{2}+4\right)+2\left(N_{B}^{2}+1\right)\right) \\ &+ C[2^{2}]N_{A}^{2}N_{B}^{2}\left(2N_{A}^{2}+3N_{B}^{2}+4\right) \\ &+ C[2,1^{2}]2N_{A}N_{B}\left(N_{A}^{2}\left(N_{B}^{2}+4\right)+2\left(N_{B}^{2}+1\right)\right) \\ &+ C[3,1]24N_{A}^{2}N_{B}^{2} \\ &+ C[4]2N_{A}N_{B}\left(2N_{A}^{2}\left(N_{B}^{2}+1\right)+4N_{B}^{2}+1\right) \right] \\ &+ \operatorname{Tr}\Lambda_{X}^{3} \left[ C[1^{4}]8N_{A}^{2}N_{B}^{2} \\ &+ C[3,1]8N_{A}N_{B}\left(N_{A}^{2}\left(N_{B}^{2}+2\right)+2N_{B}^{2}+1\right) \\ &+ C[2^{2}]24N_{A}^{2}N_{B}^{2}\left(N_{A}^{2}+N_{B}^{2}+6\right) \\ &+ C[4]8N_{A}N_{B}\left(N_{A}^{2}\left(N_{B}^{2}+2\right)+2N_{B}^{2}+1\right) \right] \\ &+ \operatorname{Tr}\Lambda_{X}^{4} \left[ C[1^{4}]2N_{A}\left(2N_{A}^{2}+1\right)N_{B}+C[2,1^{2}]4N_{A}^{2}\left(N_{A}^{2}+8\right)N_{B}^{2} \\ &+ C[2^{2}]2N_{A}N_{B}\left(2N_{A}^{2}\left(N_{B}^{2}+2\right)+2N_{B}^{2}+1\right) \\ &+ C[3,1]8N_{A}N_{B}\left(N_{A}^{2}\left(N_{B}^{2}+2\right)+2N_{B}^{2}+1\right) \\ &+ C[4]2N_{A}^{2}N_{B}^{2}\left(N_{A}^{2}+3N_{B}^{2}+14\right) \right] \right\},$$
(51)

from which it follows  $\hfill \Gamma$ 

$$\mathcal{M}_{2}^{A}(\Lambda_{X}) = c_{N_{A},N_{B}} \left[ (N_{B}^{2} - 1)(N_{A}^{4}N_{B}^{2}(N_{B}^{2} - 1) - 2N_{A}^{2}(6N_{B}^{2} - 7) + 22) \right. \\ \left. + \operatorname{Tr} \Lambda_{X}^{2} (2N_{A}N_{B}(N_{A}^{2} - 1)(N_{B}^{2} - 1)(N_{A}^{2}N_{B}^{2} - 14)) \right. \\ \left. + (\operatorname{Tr} \Lambda_{X}^{2})^{2} (N_{A}^{2} - 1)(N_{B}^{4}N_{A}^{4} + N_{B}^{4}N_{A}^{2} - 14N_{A}^{2}N_{B}^{2} + 6N_{B}^{2} + 30) \right. \\ \left. + \operatorname{Tr} \Lambda_{X}^{3} 40(N_{A}^{2} - 1)(N_{B}^{2} - 1) \right. \\ \left. + \operatorname{Tr} \Lambda_{X}^{4} (-10N_{A}N_{B})(N_{A}^{2} - 1)(N_{B}^{2} - 1) \right],$$
(52)

12

where

$$c_{N_A,N_B} = \frac{1}{N_A^2 N_B^2 (N_A^2 N_B^2 - 7)^2 - 36}.$$
(53)

This expression generalizes the already known result for the pure case, when  $\operatorname{Tr} \Lambda_X^k = 1$  for all k [17, 18]. In particular if  $N_B = N_A = \sqrt{N}$  we get

$$\mathcal{M}_{2}^{4}(\Lambda_{X}) = \left\{ \begin{bmatrix} C[1^{4}]N^{3} + C[2, 1^{2}]2N^{2}(N+2) \\ + C[2^{2}]N^{2}(N+2) + C[3, 1]8N^{2} + C[4]2N(2N+1) \end{bmatrix} \\ + \operatorname{Tr} \Lambda_{X}^{2} \left[ C[1^{4}]2N^{2}(N+2) + C[2, 1^{2}]4N^{2}(2N+7) \\ + C[2^{2}]2N(N(N+6)+2) \\ + C[3, 1]8N(N(N+4)+1) + C[4]4N^{2}(N+8) \end{bmatrix} \\ + (\operatorname{Tr} \Lambda_{X}^{2})^{2} \left[ C[1^{4}]N^{2}(N+2) + C[2, 1^{2}]2N(N(N+6)+2) \\ + C[2^{2}]N^{2}(5N+4) + C[3, 1]24N^{2} + C[4]2N(2N(N+3)+1) \right] \\ + \operatorname{Tr} \Lambda_{X}^{3} \left[ C[1^{4}]8N^{2} + C[2, 1^{2}]8N(N(N+4)+1) \\ + C[2^{2}]24N^{2} + C[3, 1]16N^{2}(N+3) + C[4]8N(N(N+4)+1) \right] \\ + \operatorname{Tr} \Lambda_{X}^{4} \left[ C[1^{4}]2N(2N+1) + C[2, 1^{2}]4N^{2}(N+8) \\ + C[2^{2}]2N(2N(N+3)+1) + C[3, 1]8N(N(N+4)+1) \\ + C[4]4N^{2}(2N+7) \right] \right\},$$
(54)

that is

$$\mathcal{M}_{2}^{A}(\Lambda_{X}) = c_{N} \bigg[ (N^{5} - 2N^{4} - 11N^{3} + 26N^{2} + 8N - 22) + \operatorname{Tr} \Lambda_{X}^{2} (2N^{5} - 4N^{4} - 26N^{3} + 56N^{2} - 28N) + (\operatorname{Tr} \Lambda_{X}^{2})^{2} (N^{5} - 15N^{3} + 20N^{2} + 24N - 30) + \operatorname{Tr} \Lambda_{X}^{3} 40(N - 1)^{2} + \operatorname{Tr} \Lambda_{X}^{4} (-10N)(N - 1)^{2} \bigg],$$
(55)

with

$$c_N = \frac{1}{N^2 (N^2 - 7)^2 - 36}.$$
(56)

In the thermodynamical limit, N >> 1, we find

$$\mathcal{M}_{2}^{A} = \frac{1}{N}(1+x)^{2} + \mathcal{O}\left(\frac{1}{N^{2}}\right).$$
 (57)

From Eqs. (42), (52) and (53) we can now compute the exact expression for the second cumulant of the purity at  $\beta = 0$ :

$$\mathcal{K}_2^A(x,0) = \mathcal{M}_2^A(x,0) - (\mathcal{M}_1^A(x,0))^2$$



Figure 1. (Color online) First moment of the purity as a function of  $\beta$ . The red line refers to the set of totally mixed states, with  $\mathcal{M}_1^A(1/N,\beta) = 1/N_A$ . The black line refers to pure states for  $\beta > 0$ . Finally the blue line corresponds to the first moment of the local purity at  $\beta = 0$ , for arbitrary mixed states, while the light-blue region refers to high temperatures.

$$= + \frac{2(N_A^2 - 1)(N_B^2 - 1)(N_A^2 N_B^2 + 11)}{(N_A^2 N_B^2 - 1)^2 (N_A^4 N_B^4 - 13N_A^2 N_B^2 + 36)} + x \frac{2(N_A^2 - 1)(N_B^2 - 1)(-2N_A N_B)(N_A^2 N_B^2 + 11)}{(N_A^2 N_B^2 - 1)^2 (N_A^4 N_B^4 - 13N_A^2 N_B^2 + 36)} + x^2 \frac{2(N_A^2 - 1)(N_B^2 - 1)(N_A^4 N_B^4 - 4N_A^2 N_B^2 + 15)}{(N_A^2 N_B^2 - 1)^2 (N_A^4 N_B^4 - 13N_A^2 N_B^2 + 36)} + \langle \operatorname{Tr} \Lambda_X^3 \rangle_x \frac{40(N_A^2 - 1)(N_B^2 - 1)}{N_A^2 N_B^2 (N_A^2 N_B^2 - 7)^2 - 36} + \langle \operatorname{Tr} \Lambda_X^4 \rangle_x \frac{(-10N_A N_B)(N_A^2 - 1)(N_B^2 - 1)}{N_A^2 N_B^2 (N_A^2 N_B^2 - 7)^2 - 36}.$$
(58)

See Ref. [14, 24] for the case x = 1, when all the traces are 1.

# 3.4. High temperature expansion of the first moment of the purity

We can now compute the approximate expression for the second moment of the purity for small  $\beta$ , by plugging Eqs. (41) and (58) in Eq. (25), with n = 2:

$$\begin{split} \mathcal{M}_{1}^{A}(x,\beta) &\sim \mathcal{M}_{1}^{A}(x,0) + \beta \left[ (\mathcal{M}_{1}^{A}(x,0))^{2} - \mathcal{M}_{2}^{A}(x,0) \right] \\ &= \frac{N_{B}(N_{A}^{2}-1)}{N_{A}^{2}N_{B}^{2}-1} \; x + \frac{N_{A}(N_{B}^{2}-1)}{N_{A}^{2}N_{B}^{2}-1} \\ &+ \beta \left[ -\frac{2(N_{A}^{2}-1)(N_{B}^{2}-1)(N_{A}^{2}N_{B}^{2}+11)}{(N_{A}^{2}N_{B}^{2}-1)^{2}(N_{A}^{4}N_{B}^{4}-13N_{A}^{2}N_{B}^{2}+36)} \\ &- x \frac{2(N_{A}^{2}-1)(N_{B}^{2}-1)(-2N_{A}N_{B})(N_{A}^{2}N_{B}^{2}+11)}{(N_{A}^{2}N_{B}^{2}-1)^{2}(N_{A}^{4}N_{B}^{4}-13N_{A}^{2}N_{B}^{2}+36)} \\ &- x^{2} \; \frac{2(N_{A}^{2}-1)(N_{B}^{2}-1)(N_{A}^{2}N_{B}^{4}-4N_{A}^{2}N_{B}^{2}+15)}{(N_{A}^{2}N_{B}^{2}-1)^{2}(N_{A}^{4}N_{B}^{4}-13N_{A}^{2}N_{B}^{2}+36)} \end{split}$$

$$- \langle \operatorname{Tr} \Lambda_X^3 \rangle_x \frac{40(N_A^2 - 1)(N_B^2 - 1)}{N_A^2 N_B^2 (N_A^2 N_B^2 - 7)^2 - 36} - \langle \operatorname{Tr} \Lambda_X^4 \rangle_x \frac{(-10N_A N_B)(N_A^2 - 1)(N_B^2 - 1)}{N_A^2 N_B^2 (N_A^2 N_B^2 - 7)^2 - 36} \right].$$
(59)

For a balanced bipartition,  $N_B = N_A = \sqrt{N}$ , we get

$$\mathcal{M}_{1}^{A}(x,\beta) \sim \mathcal{M}_{1}^{A}(x,0) + \beta \left[ (\mathcal{M}_{1}^{A}(x,0))^{2} - \mathcal{M}_{2}^{A}(x,0) \right] \\= \frac{\sqrt{N}(1+x)}{1+N} \\+ \beta \left[ -\frac{2(N^{2}+11)}{(N+1)^{2}(N^{4}-13N^{2}+36)} \\+ x \frac{4N(N^{2}+11)}{(N+1)^{2}(N^{4}-13N^{2}+36)} \\- x^{2} \frac{2(N^{4}-4N^{2}+15)}{(N+1)^{2}(N^{4}-13N^{2}+36)} \\- \langle \operatorname{Tr} \Lambda_{X}^{3} \rangle_{x} \frac{40(N-1)^{2}}{N^{2}(N^{2}-7)^{2}-36} \\- \langle \operatorname{Tr} \Lambda_{X}^{4} \rangle_{x} \frac{(-10N)(N-1)^{2}}{N^{2}(N^{2}-7)^{2}-36} \right],$$
(60)

and in the thermodynamical limit

$$\mathcal{M}_1^A(x,\beta) \sim \frac{1+x}{\sqrt{N}} - \frac{2\beta}{N^2} x^2 + \mathcal{O}\left(\frac{1}{N^{3/2}}\right).$$
(61)

One might wonder whether higher order cumulants follow a pattern similar to (61). Notice that (61) suggests a convergence radius for the high temperature expansion  $\beta_c \sim N^{3/2}(1+x)/2x^2$ , which grows indefinitely when  $x \to 0$ , in accordance with Eq. (21). See also Fig. 1.

Equation (61) at x = 1 can be compared with the results of Ref. [8] where  $\beta$  was replaced by the scaled quantity  $\beta = \beta' N^{3/2}$ . With this choice our expression yields

$$\mathcal{M}_{1}^{A}(1,\beta'N^{3/2}) \sim \mathcal{M}_{1}^{A}(1,0) + \beta'N^{3/2} \left[ (\mathcal{M}_{1}^{A}(1,0))^{2} - \mathcal{M}_{2}^{A}(1,0) \right] \\ \sim (1-\beta') \frac{2}{\sqrt{N}} , \qquad (62)$$

in perfect agreement with the behavior reported in Fig. 2 of Ref. [8]. Figure 1 yields an overview and summarizes our results.

# 4. An alternative approach based on Twirling

In this section we will establish an interesting connection between our problem and the theory of quantum channels. In particular, we present an alternative approach for computing the moments  $\mathcal{M}_n^A(x,0)$  of Eq. (22) which exploits the properties of *twirling transformations* [25, 26, 27, 19, 20].

For explanatory purposes we start in Sec. 4.1 by deriving a general expressions for the moments  $\mathcal{M}_n^A(x = 1, 0)$  associated with the case in which  $\rho$  is pure (i.e. the quantities  $\mathcal{M}_n$  of Eq. (10)) and verify that it yields the exact value given by Lubkin [14] for n = 1. The case of mixed states is then addressed in Sec. 4.2 showing that for n = 1 it reproduces the results of Sec. 3.1.

# 4.1. Pure initial states

Let us consider a fixed (normalized) pure state of the global system X = AB,  $|\psi_0\rangle_X$ , and parametrize the pure states of  $\mathcal{H}_X$  as in (5), i.e.  $|\psi\rangle_X := U_X |\psi_0\rangle_X$ , with  $U_X \in \mathcal{U}(\mathcal{H}_X) \simeq \mathcal{U}(N)$  distributed according to the Haar measure  $d\mu_H(U)$ . Its local purity (3) can be expressed as

$$\pi(\psi) = \sum_{1 \le \ell, \ell' \le N_B} \operatorname{Tr} \left( {}_{B} \langle \ell | U_{AB} | \psi_0 \rangle_{AB} \langle \psi_0 | U_{AB}^{\dagger} | \ell \rangle_{B} {}_{B} \langle \ell' | U_{AB} | \psi_0 \rangle_{AB} \langle \psi_0 | U_{AB}^{\dagger} | \ell' \rangle_{B} \right)$$
$$= \sum_{1 \le \ell, \ell' \le N_B} {}_{AB} \langle \psi_0 | \left( U_{AB}^{\dagger} | \ell \rangle_B \langle \ell' | U_{AB} \right) | \psi_0 \rangle_{AB} {}_{AB} \langle \psi_0 | \left( U_{AB}^{\dagger} | \ell' \rangle_B \langle \ell | U_{AB} \right) | \psi_0 \rangle_{AB} , \quad (63)$$

where  $\{|\ell\rangle_B\}$  is an orthonormal basis of  $\mathcal{H}_B$ , and the cyclicity of the trace was used. We can recast this expression into a more compact form by doubling the Hilbert space, i.e. adding two auxiliary copies A' and B' of A and B, respectively. We get

$$\pi(\psi) = \operatorname{Tr}\left[ \left( U_{AB} \otimes U_{A'B'} \right) \left( |\psi_0\rangle_{AB} \langle \psi_0| \otimes |\psi_0\rangle_{A'B'} \langle \psi_0| \right) \left( U_{AB}^{\dagger} \otimes U_{A'B'}^{\dagger} \right) \times \left( S_{B|B'} \otimes I_{AA'} \right) \right],$$
(64)

where the trace is over all degree of freedom (i.e. AA'BB'),  $I_{AA'}$  is the identity operator on AA', and

$$S_{B|B'} := \sum_{1 \le \ell, \ell' \le N_B} |\ell\rangle_B \langle \ell'| \otimes |\ell'\rangle_{B'} \langle \ell| , \qquad (65)$$

is the SWAP operator on BB' – this is the unitary, self-adjoint transformation which, for all operators  $\Theta_B$  and  $\Upsilon_{B'}$ , gives

$$S_{B|B'}(\Theta_B \otimes \Upsilon_{B'}) S_{B|B'} = \Upsilon_B \otimes \Theta_{B'}.$$
(66)

Remembering that first moment  $\mathcal{M}_1^A(x = 1, 0) = \mathcal{M}_1$  of Eq. (10) is obtained by averaging over all possible  $U_X$ , we can then write

$$\mathcal{M}_1 = \operatorname{Tr}\left(\mathcal{T}^{(2)}(|\psi_0\rangle_X \langle \psi_0| \otimes |\psi_0\rangle_{X'} \langle \psi_0|) \left(S_{B|B'} \otimes I_{AA'}\right)\right), \tag{67}$$

where X = AB, X' = A'B', and where  $\mathcal{T}^{(2)}$  is the Completely Positive, Trace Preserving (CPTP) twirling channel [25, 26, 27, 28] which transforms the operators  $\Theta_{XX'}$  of XX' into

$$\mathcal{T}^{(2)}(\Theta_{XX'}) = \int d\mu_H(U) \ (U_X \otimes U_{X'}) \ \Theta_{XX'} \ (U_X^{\dagger} \otimes U_{X'}^{\dagger}).$$
(68)

This map plays an important role in quantum information theory where it was first introduced as a tool for characterizing the distillability of bipartite entanglement [20, 19]. It has several properties which allows us to simplify the calculation. For instance it is known that  $\mathcal{T}^{(2)}$  maps all the states of the system into (generalized) Werner

$$\mathcal{M}_1 = \left({}_X \langle \psi_0 | \otimes_{X'} \langle \psi_0 | \right) \mathcal{T}^{(2)} \left( S_{B|B'} \otimes I_{AA'} \right) \left( |\psi_0\rangle_X \otimes |\psi_0\rangle_{X'} \right).$$
(69)

Explicit expressions for the action of  $\mathcal{T}^{(2)}$  can be obtained by exploiting the symmetry of  $d\mu_H(U)$ . In particular it is possible to show that  $\mathcal{T}^{(2)}(\Theta_{XX'})$  can be decomposed as a linear combination of the projections on the symmetric and anti-symmetric subspaces of XX' = ABA'B' (with respect to the bipartition AB|A'B'). Introducing then the SWAP operator which exchanges X with X' this can then be written as

$$\mathcal{T}^{(2)}(\Theta_{XX'}) = \frac{NI_{XX'} - S_{X|X'}}{N(N^2 - 1)} \operatorname{Tr} \Theta_{XX'} + \frac{NS_{X|X'} - I_{XX'}}{N(N^2 - 1)} \operatorname{Tr}(S_{X|X'}\Theta_{XX'})$$
(70)  
$$- \frac{N\operatorname{Tr} \Theta_{XX'} - \operatorname{Tr}(S_{X|X'}\Theta_{XX'})}{N(N^2 - 1)} I_{XX'} + \frac{N\operatorname{Tr}(S_{X|X'}\Theta_{XX'}) - \operatorname{Tr} \Theta_{XX'}}{N(N^2 - 1)} S_{XX'}$$
(70)

$$= \frac{(1,1,2,...,N)}{N(N^2 - 1)} I_{XX'} + \frac{(1,1,2,...,N)}{N(N^2 - 1)} S_{X|X'},$$
(71)

where  $N = N_A N_B$  is the dimension of  $\mathcal{H}_{AB}$ . (Here  $I_{XX'}$  is the identity operator while  $S_{X|X'} = S_{AB|A'B'} = S_{B|B'} \otimes S_{A|A'}$  is the SWAP which exchanges AB with A'B').

Thanks to this expression we can now easily compute the value of  $\mathcal{M}_1$ , either using Eq. (63) or Eq. (69). Consider for instance the first approach. We have first to compute the quantities  $\operatorname{Tr} \Theta_{XX'}$  and  $\operatorname{Tr}(S_{X|X'}\Theta_{XX'})$  with  $\Theta_{XX'}$  being the operator  $|\psi_0\rangle_X\langle\psi_0|\otimes|\psi_0\rangle_{X'}\langle\psi_0|$ . This is

$$\operatorname{Tr}(|\psi_0\rangle_X \langle \psi_0| \otimes |\psi_0\rangle_{X'} \langle \psi_0|) = 1 ,$$
  
$$\operatorname{Tr}\left((S_{B|B'} \otimes S_{A|A'})|\psi_0\rangle_{AB} \langle \psi_0| \otimes |\psi_0\rangle_{A'B'} \langle \psi_0|\right) = 1 ,$$
(72)

where in the second expression we used the fact that  $|\psi\rangle_{AB} \otimes |\psi\rangle_{A'B'}$  is invariant uder  $S_{X|X'}$ , i.e.  $(S_{B|B'} \otimes S_{A|A'})(|\psi\rangle_{AB} \otimes |\psi\rangle_{A'B'}) = |\psi\rangle_{AB} \otimes |\psi\rangle_{A'B'}$ . Replacing all this in Eq. (70) we get

$$\mathcal{T}^{(2)}(|\psi_0\rangle_X\langle\psi_0|\otimes|\psi_0\rangle_{X'}\langle\psi_0|) = \frac{I_{XX'} + S_{X|X'}}{N(N+1)},$$
(73)

and thus

$$\mathcal{M}_{1} = \frac{1}{N(N+1)} \operatorname{Tr} \left( (I_{ABA'B'} + S_{B|B'} \otimes S_{A|A'}) \left( S_{B|B'} \otimes I_{AA'} \right) \right)$$
$$= \frac{1}{N(N+1)} \left( \operatorname{Tr} \left( S_{B|B'} \otimes I_{AA'} \right) + \operatorname{Tr} \left( I_{BB'} \otimes S_{A|A'} \right) \right)$$
(74)

(here we exploited the fact that  $S^2_{B|B'} = I_{BB'}$ ). Now we can use  $N = N_A N_B$  and the identities

$$\operatorname{Tr}\left(S_{B|B'}\otimes I_{AA'}\right) = N_B N_A^2 , \quad \operatorname{Tr}\left(I_{BB'}\otimes S_{A|A'}\right) = N_A N_B^2 , \tag{75}$$

to get

$$\mathcal{M}_1 = \frac{N_A + N_B}{N_A N_B + 1} , \qquad (76)$$

which coincides with the correct value [14].

We mention that the same techniques can also be applied to higher moments  $\mathcal{M}_n$ . The extension of Eq. (67) for  $n \ge 2$  is obtained by introducing 2n copies of AB organized in the *n* pairs, i.e.  $A_1B_1A'_1B'_1$ ,  $A_2B_2A'_2B'_2$ ,  $\cdots$ ,  $A_nB_nA'_nB'_n$ . We then introduce the following generalized twirling transformation acting on  $XX' = A_1B_1A'_1B'_1$ ,  $A_2B_2A'_2B'_2$ ,  $\cdots$ ,  $A_nB_nA'_nB'_n$ , i.e.

$$\mathcal{T}^{(2n)}(\Theta_{XX'}) = \int d\mu_H(U) \; (\underbrace{U \otimes U \otimes \cdots \otimes U}_{2n}) \; \Theta_{XX'} \; (\underbrace{U^{\dagger} \otimes U^{\dagger} \otimes \cdots \otimes U^{\dagger}}_{2n}), \qquad (77)$$

with  $\Theta_{XX'}$  being a generic operator on  $\mathcal{H}_{XX'} := \mathcal{H}_{AB}^{\otimes 2n}$ . This channel is a proper generalization of the map  $\mathcal{T}^{(2)}$  whose properties can be established along the lines of Ref. [10]. With this choice Eq. (10) then can be expressed as

$$\mathcal{M}_{n}^{A}(x=1,0) = \mathcal{M}_{n} = \operatorname{Tr}\left(\mathcal{T}^{(2n)}\left(|\Psi^{\otimes 2}\rangle\langle\Psi^{\otimes 2}|^{\otimes n}\right)(S_{\mathcal{B}}^{(2n)}\otimes I_{\mathcal{A}}^{(2n)})\right), \quad (78)$$

where  $|\Psi^{\otimes 2}\rangle^{\otimes n} := \bigotimes_{j=1}^{n} (|\psi\rangle_{A_{j}B_{j}} \otimes |\psi\rangle_{A'_{j}B'_{j}})$ ,  $I_{\mathcal{A}}$  being the identity on the 2*n* copies of A, i.e.  $\mathcal{A} = A_{1}A'_{1}\cdots A_{n}A'_{n}$ , and with  $S_{\mathcal{B}}^{(2n)}$  being the SWAP operator which exchanges  $B_{1}B_{2}\cdots B_{n}$  with  $B'_{1}B'_{2}\cdots B'_{n}$  pairwise, i.e.  $S_{\mathcal{B}}^{(2n)} = \bigotimes_{j=1}^{n} S_{B_{j}|B'_{j}}$ .

#### 4.2. Mixed initial states

Consider now the case with x < 1. Following the parameterization introduced in Secs. 2 and 3, we split the average over the set  $\mathfrak{S}_x(\mathcal{H}_X)$  of the density matrixes of global purity x, as an average over the unitary rotations of acting on  $\mathcal{H}_X$  followed by an average over the space of the eigenvalues  $\rho$  – see Eq. (16). Specifically this is accomplished by writing  $\rho(U) = U_{AB} \Lambda_{AB} U_{AB}^{\dagger}$ , with  $U_{AB}$  being a generic unitary transformation on  $\mathcal{H}_{AB}$ , while  $\Lambda_{AB}$  represents a given arbitrary choice of the system spectrum, see Eq. (26). For convenience, let us rewrite the purification (27) of the density matrix  $\rho$  as

$$|\Psi\rangle_{ABab} = \sum_{1 \le \alpha \le N_A} \sum_{1 \le \beta \le N_B} \sqrt{\lambda_{\alpha\beta}} \ |\alpha\rangle_A \otimes |\beta\rangle_B \otimes |\alpha\rangle_a \otimes |\beta\rangle_b , \qquad (79)$$

where now the ancillary systems, isomorphic to A and B, are labelled by a and b, respectively. The reduced density matrix  $\rho_A(U) = \operatorname{Tr}_B \rho(U)$  can thus be written as

$$\rho_A(U) = \operatorname{Tr}_{Bab}(U_{AB}|\Psi\rangle_{ABab}\langle\Psi|U_{AB}^{\dagger}) = \sum_{1 \le q \le N_A N_B^2} {}_{Bab}\langle q|U_{AB}|\Psi\rangle_{ABab}\langle\Psi|U_{AB}^{\dagger}|q\rangle_{Bab} , \qquad (80)$$

with  $\{|q\rangle_{Bab}\}$  being an orthonormal basis of *Bab*. Similarly the local A-purity of  $\rho_A(U)$  becomes

$$\pi_{A}(\rho) = \operatorname{Tr} \rho_{A}^{2}(U) = \sum_{1 \leq q, q' \leq N_{A}N_{B}^{2}} \operatorname{Tr} \left( Bab \langle q | U_{AB} | \Psi \rangle_{ABab} \langle \Psi | U_{AB}^{\dagger} | q \rangle_{Bab} \right)$$
$$\times Bab \langle q' | U_{AB} | \Psi \rangle_{ABab} \langle \Psi | U_{AB}^{\dagger} | q' \rangle_{Bab} \left)$$
$$= \sum_{1 \leq q, q' \leq N_{A}N_{B}^{2}} ABab \langle \Psi | U_{AB}^{\dagger} | q \rangle_{Bab} \langle q' | U_{AB} | \Psi \rangle_{ABab}$$
$$\times ABab \langle \Psi | U_{AB}^{\dagger} | q' \rangle_{Bab} \langle q | U_{AB} | \Psi \rangle_{ABab}$$
(81)

which, once more, can be casted as an expectation value on  $|\Psi\rangle^{\otimes 2}$  by doubling the space (see Eq. (29)). Therefore by integrating over  $U_{AB}$  we get,

$$\mathcal{M}_{1}^{A}(\Lambda_{X}) = \operatorname{Tr}\left(\mathcal{T}^{(2)}\left(|\Psi\rangle_{Xx}\langle\Psi|\otimes|\Psi\rangle_{X'x'}\langle\Psi|\right) \left(S_{Bab|B'a'b'}\otimes I_{AA'}\right)\right)$$
$$= \left(_{Xx}\langle\Psi|\otimes_{X'x'}\langle\Psi|\right)\mathcal{T}^{(2)}\left(S_{Bab|B'a'b'}\otimes I_{AA'}\right)\left(|\Psi\rangle_{Xx}\otimes|\Psi\rangle_{X'x'}\right), \quad (82)$$

where  $\mathcal{T}^{(2)}$  being the twirling transformation on XX' of Eq. (68) with X = AB, X' = A'B', x = ab, and x' = a'b' (here A', a', B', b' are the auxiliary copies of A, a, B, and b respectively). The above expression is the average purity of the subsystem Acomputed for states  $\rho$  having the same spectra  $\Lambda_X$ .

To compute the above quantity this time we use the last identity of Eq. (82). According to Eq. (71) we have to compute  $\operatorname{Tr} \Theta_{XX'}$  and  $\operatorname{Tr}(S_{X|X'}\Theta_{XX'})$  with  $\Theta_{XX'}$ being the operator  $S_{Bab|B'a'b'} \otimes I_{AA'} = S_{B|B'} \otimes S_{b|b'} \otimes S_{a|a'} \otimes I_{AA'}$ . That is

$$\operatorname{Tr}_{ABA'B'}(S_{Bab|B'a'b'} \otimes I_{AA'}) = \operatorname{Tr}_{ABA'B'}(S_{B|B'} \otimes S_{b|b'} \otimes S_{a|a'} \otimes I_{AA'})$$
$$= \operatorname{Tr}(S_{B|B'}) \operatorname{Tr}(I_{AA'}) S_{b|b'} \otimes S_{a|a'} = N_A^2 N_B S_{b|b'} \otimes S_{a|a'} , \qquad (83)$$

$$\operatorname{Tr}_{ABA'B'}\left(S_{AB|A'B'}(S_{Bab|B'a'b'}\otimes I_{AA'})\right) = \operatorname{Tr}_{ABA'B'}\left(I_{BB'}\otimes S_{A|A'}\otimes S_{b|b'}\otimes S_{a|a'}\right)$$
$$= \operatorname{Tr}(I_{BB'}) \operatorname{Tr}(S_{A|A'}) S_{b|b'}\otimes S_{a|a'} = N_B^2 N_A S_{b|b'}\otimes S_{a|a'} .$$
(84)

Thus from Eq. (71) we get

$$\mathcal{T}^{(2)}\Big(S_{Bab|B'a'b'} \otimes I_{AA'}\Big) = \frac{N_B(N_A^2 - 1)}{N_A^2 N_B^2 - 1} I_{BB'} \otimes I_{AA'} \otimes S_{b|b'} \otimes S_{a|a'} + \frac{N_A(N_B^2 - 1)}{N_A^2 N_B^2 - 1} S_{B|B'} \otimes S_{A|A'} \otimes S_{b|b'} \otimes S_{a|a'} , \qquad (85)$$

where we used  $N = N_A N_B$ . Replace now this into Eq. (82) and employ the identities

$$\begin{pmatrix} X_x \langle \Psi | \otimes_{X'x'} \langle \Psi | \end{pmatrix} (I_{XX'} \otimes S_{x|x'}) \left( |\Psi \rangle_{Xx} \otimes |\Psi \rangle_{X'x'} \right) = \operatorname{Tr} \Lambda_X^2 , \begin{pmatrix} X_x \langle \Psi | \otimes_{X'x'} \langle \Psi | \end{pmatrix} (S_{X|X'} \otimes S_{x|x'}) \left( |\Psi \rangle_{Xx} \otimes |\Psi \rangle_{X'x'} \right) = 1 .$$

$$(86)$$

The final result is thus

$$\mathcal{M}_{1}^{A}(\Lambda_{X}) = \frac{N_{B}(N_{A}^{2}-1)}{N_{A}^{2}N_{B}^{2}-1} \operatorname{Tr} \Lambda_{X}^{2} + \frac{N_{A}(N_{B}^{2}-1)}{N_{A}^{2}N_{B}^{2}-1}$$
(87)

which is Eq.(41) and depends upon the spectrum  $\Lambda_X$  only through its purity. By averaging upon on  $\Lambda_X$  while keeping fix x gives us the same result (42).

# 5. Conclusions

From the results obtained in the previous sections one can infer that the same phenomenon of concentration of measure that occurs for the eigenvalues of the reduced density matrices of pure states [8, 9] occurs in the present case as well. Indeed, we observe that for large N the leading order of all k-moments equals the k-th power of the first moment (43), that, for a balanced bipartition  $N_A = N_B = \sqrt{N}$ , reads

$$\mathcal{M}_1^A(x,0) = \frac{(1+x)}{\sqrt{N}} + \mathcal{O}\left(\frac{1}{N^{3/2}}\right).$$
(88)

This observation spurs from the calculation of the second moment (55):

$$\mathcal{M}_{2}^{A}(x,0) = \frac{(1+x)^{2}}{N} + \mathcal{O}\left(\frac{1}{N^{2}}\right).$$
(89)

Therefore,

$$\mathcal{M}_2^A(x,0) = \mathcal{M}_1^A(x,0)^2 + \mathcal{O}\left(\frac{1}{N^2}\right).$$
(90)

By observing that the contributions of  $\mathcal{O}(1/N)$  to the second moment come from the coefficients proportional to  $C[1^4]$  in (46), i.e. the identity permutation  $\sigma = 0$ , and conjecturing that the leading contribution for all the k-th moments comes only from  $C[1^k] \sim 1/N^k$  one gets

$$\mathcal{M}_{n}^{A}(x,0) = \left(\frac{(1+x)}{\sqrt{N}}\right)^{k} + \mathcal{O}\left(\frac{1}{N^{(k+2)/2}}\right).$$
(91)

Another check of the validity of (91) derives by the interpolation between maximally mixed and pure global states. See Fig. 1. The scaling with N is preserved, therefore allowing to interchange the  $x \to 1$  and  $N \to \infty$  limits.

This uncovers the issue of computing the subdominant terms in the k-th moments expansion that become instead *leading* order terms in the k-th *cumulants*. This could be possible if we had an appropriate asymptotic expansion of the combinatorics coefficients C, which we do not know. We leave this as a challenge for future work.

# References

- M.A. Nielsen and I.L. Chuang, Quantum Computation and Quantum Information (Cambridge University Press, Cambridge, 2000).
- [2] G. Benenti, G. Casati and G. Strini, Principles of quantum computation and information, Volume I: Basic concepts (World Scientific, Singapore, 2004).
- [3] S. Bose and V. Vedral, Phys. Rev. A 61, 040101(R) (2000).
- [4] G. Adesso, A. Serafini, F. Illuminati, Phys. Rev. A 70 (2004).
- [5] L. Mazzola, B. Bellomo, R. Lo Franco, and G. Compagno, Phys. Rev. A 81, 052116 (2010)
- [6] O. Gühne and G. Tóth, Phys. Rep. 474, 1 (2009).
- [7] D. Bruss, J. Math. Phys. 43, 4237 (2002).
- [8] P. Facchi, U. Marzolino, G. Parisi, S. Pascazio and A. Scardicchio, Phys. Rev. Lett. 101 (2008) 050502.
- [9] A. De Pasquale, P. Facchi, G. Parisi, S. Pascazio and A. Scardicchio, Phys. Rev. A 81 (2010) 052324.
- [10] J. B. Zuber, J. Phys. A: Math. Theor. **41** (2008) 382001.
- [11] M. M. Sinolecka, K. Zyczkowski, M. Kus, Acta Physica Polonica B 33, 2081 (2001).
- [12] K. Zyczkowski, P. Horodecki, A. Sanpera and M. Lewenstein, Phys. Rev. A 58 (1998) 833.
- [13] K. Zyczkowski, Phys. Rev. A 60 (1999) 3496.
- [14] E. Lubkin, J. Math. Phys. 19, 1028 (1978).
- [15] S. Lloyd and H. Pagels, Ann. Phys., NY, **188**, 186 (1988).
- [16] D.N. Page, Phys. Rev. Lett. **71**, 1291 (1993).
- [17] K Zyczkowski and H.-J. Sommers, J. Phys. A 34, 7111 (2001)
- [18] A. J. Scott and C. M. Caves, J. Phys. A: Math. Gen. 36, 9553 (2003).
- [19] C. H. Bennett, D. P. DiVincenzo, J. A. Smolin, and W. K. Wootters, Phys. Rev. A 54, 3824 (1996).

- [20] C. H. Bennett, G. Brassard, S. Popescu, B. Schumacher, J. A. Smolin, and W. K. Wootters, Phys. Rev. Lett. 76, 722 (1996).
- [21] M. F. Cornelio and M. C. de Oliveira, Phys. Rev. A 81, 032332 (2010).
- [22] K. Zyczkowski and H. Sommers, J. Phys. A: Math. Gen 75 (2007) 7111.
- [23] P. B. Slater, J. Phys. A: Math. Gen **32** (1999) 5261.
- [24] O. Giraud, J. Phys. A: Math. Theor. 40 (2007) 2793.
- [25] K. G. H. Vollebrecht and R. F. Werner, Phys. Rev. A 64, 062307 (2001).
- [26] S. Lee, et al., Phys. Rev. A 68, 062304 (2003).
- [27] M. Hayashi, et al., Phys. Rev. A 77, 012104 (2008).
- [28] Bennett C H, DiVincenzo D P, Mor T, Shor P W, Smolin J A, and Terhal B M 1999 Phys. Rev. Lett. 82, 5385.