Unravelling the Mysteries of Matter with the CERN Large Hadron Collider An Introduction/Overview of Particle Physics Introductory Lecture August 3rd 2014

Bobby Samir Acharya International Centre for Theoretical Physics and King's College London

African School of Fundamental Physics and Applications 2014, Dakar, Senegal



KING'S College LONDON



Welcome Everyone to the 3rd African School in Fundamental Physics and its Applications

Especially welcome to the students who have travelled from all over Africa and beyond and for whom this school has been developed. Also welcome to the lecturers and the LOC and UCAD for making this possible

Finally: thank you to the numerous international institutes and universities for their support.





Particles, the Universe and the LHC









Large Hadron Collider

All known elementary particles

An introduction and overview of particle physics. Also: why and what the LHC is for. Two Mysteries: Dark Matter and the Higgs Boson



ICTP/KCL

Atoms and the Periodic Table



Particles, Matter and the Universe

We, and the matter which makes up all of the Earth is made of atoms

The planets and stars also ALL seem to be made of atoms

The galaxy contains hundreds of billions (i.e. $\sim 10^{11}$) of stars which are basically alike: so much of the galaxy is also be made of atoms.



Remarkably, all of the galaxies (and there are $\sim 10^{11}$ of them) also look quite similar!

Hence, we learn that much of the matter of the entire Universe is made of atoms

We will ADD some very important details to this picture later.



Abundance of Elements in the Universe



KING'S College LONDON

ICTP/KCL

ASP2014, UCAD, Dakar, Senegal

The Structure of Atoms

The structure and regularity of the periodic table suggests that atoms are made of even simpler objects

Electrons were discovered by Thompson in so-called cathode ray tubes around 1897.A wire filament with a current passing through was seen to emit particles with negative charge. This showed that atoms contained "electrons". α -particles (now known to be Helium nuclei) were discovered as radiation emitted by various Rutherford used beams of α -particles scattered off Gold to prove that atoms "must" contain a very dense nucleus

The α -particles were deflected at large angles, proving that atoms had "structure" - dense nucleus.

Shortly after this, the Bohr model of the atom (protons and neutrons in a dense nucleus, surrounded by a cloud of electrons) was "established".



The Atom



- The atom consists of a very dense nucleus surrounded by a "cloud of electrons"
- Atoms have sizes of order 10^{-11} to 10^{-10} m.
- Nucleus is 10^{-4} times smaller!
- Protons are positively charged. Neutrons are neutral.
- Electrons have *minus* the charge of protons but are *much* lighter ($m_e \sim m_p/2000$)
- But are protons, neutrons Kings and electrons fundamental

ICTP/KCL

Charged particles in Electric and Magnetic Fields



- Electric current is just moving charge
- You can create a magnetic field by coiling a current carrying wire
- By the same token, a magnetic field exerts a force on charged particles
- Strong electric and magnetic fields can be used to accelerate charged particles to high energies!
- This is how particle accelerators work: you will learn this in week 3!



The galaxy is a particle collider!



- The galaxy has a strong magnetic field and charged particles!
- A high energy particle is created in a star and brought to Earth by the galactic magnetic field
- It strikes the upper atmosphere of Earth
- Multiple interactions occur between the particle and the atoms in the atmosphere
- Creates a cosmic ray "showe KING'S LONDON



The galaxy as a particle collider!



- This is a particular example when the original particle is a proton
- This example produces pions, photons, muons, electrons and neutrinos!
- A muon from a cosmic ray like this just passed through your body!
- In fact, muons and pions were discovered in cosmic rays



The Particle Zoo

- As a result of cosmic ray and ground-based particle collider experiments we now know that protons and neutrons are actually made of more fundamental particles called quarks and gluons
- We also know that protons and neutrons have many 'cousins' which are also made of quarks and gluons. Hundreds of these have been discovered over the past six decades.
- Neutrons, when not bound inside atoms, actually decay into a proton, an electron and a neutral (almost massless) particle called a neutrino.



• The most remarkable fact, though, is that *all of these hundreds of particles and their properties are* **precisely** *described by a very simple model.*



The Standard Model of Particle Physics



The entire periodic table can be explained by just the first and last columns! Last column are the force carrying particles There are three "families" of quarks and leptons. Why not just one? Why these particular masses? Are there more than just these particles?



ICTP/KCL

Bobby Samir Acharya

ASP2014, UCAD, Dakar, Senegal

Mass and The Higgs Boson. Original Slide from 2011. See next slide for update

- There is one particle predicted by the Standard Model which has not yet been discovered
- This is the so-called Higgs boson, after Peter Higgs who made significant contributions to this part of the Standard Model
- According to the model, it is the Higgs boson which is responsible for giving the quarks and leptons their mass
- The heavier the particle, the stronger its interactions with the Higgs particle, so the top quark has the strongest interaction with the Higgs, the electron the weakest one.
- One of the main reasons for the LHC physics programme is to find the Higgs boson, or prove that it doesn't exist!
- It's discovery would shed enormous light on the nature and origin of mass and "complete" the story of the Standard



Bobby Samir Acharya ASP2014, UCAD, Dakar, Senegal

Mass and The Higgs Boson. Slide updated 2013

- There is one additional particle predicted by the Standard Model which was not yet mentioned.
- This is the so-called Higgs boson, after Peter Higgs who made significant contributions to this part of the Standard Model in 1964
- According to the model, it is the Higgs boson which is responsible for giving the quarks and leptons their mass
- The heavier the particle, the stronger its interactions with the Higgs particle, so the top quark has the strongest interaction with the Higgs, the electron the weakest one.
- One of the main reasons for the LHC physics programme was to find the Higgs boson, or prove that it doesn't exist!



• It's discovery would shed enormous light on the nature and origin of mass and "complete" the story of the Standard



ICTP/KCL

The LHC and The Higgs Boson

- Much data from the LHC collected in 2011 and 2012
- The first evidence for the existence of the Higgs boson has emerged
- A new particle has been discovered
- So far its properties are remarkably consistent with the Higgs boson
- More data is being collected and analysed to determine if the Standard Model is correct
- These are very exciting times. More about the LHC and the Higgs boson later in the course.



Dark Matter

- Einstein's theory of gravity has been tested with remarkable accuracy in various systems within the galaxy
- But, if you estimate the mass of galaxies by the number of atoms that they contain, there's a huge problem
- Galaxies of that mass would not move in the way that the galaxies have been observed to be moving
- There is now very strong evidence that there is additional, "dark" matter in the Universe, beyond protons, neutrons, electrons, photons and neutrinos
- In fact there seems to be about five times more dark matter than "atomic matter"!



 There are some plausible arguments that particles made of dark matter could be produced directly at the LHC (more on K this later)

Cosmic History



or the se in d Mode



ICTP/KCL

Bobby Samir Acharya

ASP2014, UCAD, Dakar, Senegal

The Large Hadron Collider

• The basic idea is to :

- accelerate beams of protons to as high energy as possible
- smash them into each other as often as possible
- Create new particles like the Higgs boson or dark matter
- Detect these particles in "particle detectors" which surround the "collisions"
- The production of "new particles" is rare, so the more collisions we have the better
- The LHC ran successfully in 2011 and 2012. Will restart in 2015 at even higher energy!





The LHC Accelerator complex



Bobby Samir Acharya ASP2014, UCAD, Dakar, Senegal

The LHC Accelerator complex

About 1600 superconducting magnets (most weigh more than 20 tonnes!) used to accelerate the proton beams About 100 tonnes of liquid helium required to keep the magnets at operating temperature o minus 271 Centigrade The LHC is the coolest place in the Universe! The magnetic field is about 8 Tesla (about 10^5 times that of the Earth!)

Bobby Samir Acharya ASP2014, UCAD, Dakar, Senegal

(CT

The LHC beam and Energy

- Each proton beam is made of bunches of protons. Up to 2808 bunches with 10¹¹ protons each. Energy of each proton 7000 GeV.
- Note: One GeV = about 1.6×10^{-10} Joules and 1 J = 1 Watt second.
- Energy per beam = $2808 \times 10^{11} \times 7000 \text{ GeV}$ = $2808 \times 10^{11} \times 7000 \times 1.6 \times 10^{-10} \text{J}$ = 362 MJ
- Equivalent to 87 Kilograms of TNT, which has about 10^{16} times as many protons as the LHC beam!
- Equivalent to the Kinetic Energy of a small aircraft carrier moving at 40 km per hour!
- The energy in the magnets is about 10¹⁰ Joules. (A typical house in Europe consumes about 2000 Joules per second on average).



A Particle Detector



Bobby Samir Acharya ASP2014, UCAD, Dakar, Senegal

C

ICTP/KCL

'ING'S

LONDON

A Particle Detector





Particle Detection in ATLAS





Bobby Samir Acharya ASP2014, UCAD, Dakar, Senegal

ÍCTP



Processes and Production Rates at the LHC





Bobby Samir Acharya ASP2014, UCAD, Dakar, Senegal

Producing Higgs Bosons with the LHC



Bobby Samir Acharya ASP2014, UCAD, Dakar, Senegal

Many ideas to be explored.....

- Theoretical physicists have come up with many ideas for physics beyond the Standard Model
- These predict new particles and phenomena for the LHC:
- Supersymmetry predicts a new particle for every Standard Model particle eg a super-electron (or selectron).
- The LHC may discover extra dimensions. These are predicted by string theory and could lead to the discovery of tiny black holes!
- So far, these ideas have not been discovered at the LHC, so limits on such models have been obtained!



KING'S College LONDON