

Chela-Flores, J. (1995).

Is the Salam phase transition relevant to the causal origin of homochirality ?.

Proc. Pakistan Acad. Sci. **32**, 1-12 .

**Is the Salam phase transition relevant to the
causal origin of homochirality ? (+)**

Julian Chela-Flores (*)
International Centre for Theoretical Physics
Miramare, P.O. Box 586;34100 Trieste, Italy
and
School of Theoretical Physics
Dublin Institute for Advanced Studies
10, Burlington Road, Dublin 4, Ireland.

(*) Also Instituto Internacional de Estudios Avanzados,
Apartado 17606 Parque Central, Caracas 1015A, Venezuela.

Abstract. From the point of view of (a) chemical evolution and (b) exobiology (the origin of life in the universe), we discuss the possibility that phase transitions of the type postulated by Salam in 1991 to occur in a racemic mixture of amino acids, may play instead the role of an amplification mechanism in the novel context of chirally-pure samples of amino acids. Salam's seminal contribution referred to the role of phase transitions in the induction of the observed homochirality of protein amino acids. This was assumed to occur in a racemic mixture that, as a consequence of the conjectured phase transition, turned homochiral beneath a certain critical temperature T_c . In the present review we discuss selected theoretical aspects of the question of homochirality. The relevance of starting a new generation of experiments is pointed out, as suggested by the present author in 1991. These experiments should test a form of condensed matter that has so far not been at the forefront of research from the point of view of the phase transitions. We discuss those aspects of the origin of homochirality which are still valid and require verification.

Key words for information retrieval:

Chirality, chemical evolution, exobiology, superconductivity, amino acids.

(+) To Professor Abdus Salam on the occasion of his 70th birthday. Submitted, by invitation, to the Proceedings of the Pakistan Academy of Sciences.

1. Introduction

1.1. CHEMICAL EVOLUTION AND THE ORIGIN AND EVOLUTION OF LIFE

The preliminary questions that are raised when we attempt to lay the scientific bases of the origin of life are: Where did the biomolecules come from?, Were they synthesized on Earth or, is there a need to study this problem in the context of exobiology (i.e., within a systematic search for life beyond the Earth)? In order to gain some insights into these fundamental questions it is important to recognise molecular relics of earlier stages of chemical and biological evolution. This review is devoted to one of the most important relics amongst them, but we should be aware from the beginning about the broad spectrum of conditions in which biomolecules can present themselves in the universe.

Although the extensive searches for extraterrestrial intelligence (SETI) have not been rewarded with a positive result, in spite of the enormous progress in the equipment used in the Project Phoenix, (Drake, 1996), interstellar organic molecules have in fact been detected by microwave and infrared techniques since 1969. (Prior to that date it was known that simple carbon compounds such as C_2 and CO were formed in the atmospheres of relatively cool stars.) Now we are aware of the existence of interstellar ammonia, water, formaldehyde and hydrogen cyanide. Altogether about one hundred molecules have been identified, about 75% are organic (Oro, 1994).

These discoveries have made the conjecture of Oro more plausible (Oro, 1961a), namely, that comets may have been relevant in the formation of biochemical compounds on the primitive Earth. From the old studies of the Alais meteorite in 1834, to the extensive studies of the meteorite that fell in Murchinson, Victoria (Australia) on September 28, 1969, important information about possible sources of life on Earth has been gathering. The Laboratory of Chemical Evolution (LCE) of the University of Maryland was directed by Ponnampereuma until his death in 1994. It played an important role in analyzing the Murchison meteorite, at a time when LCE was preparing to receive the first Moon samples (Ponnampereuma, 1995). The results of that investigation have an important bearing on the biochemical relic that we shall discuss presently (cf., Sec. 2.2).

1.2. CHEMICAL AND ATMOSPHERIC EVOLUTION IN THE SOLAR SYSTEM

In the early Earth atmosphere the most abundant element, hydrogen, may have combined with carbon to form methane (CH_4), with oxygen to form water (H_2O), and with nitrogen to form ammonia (H_3N). In such an environment, under the influence of an abundant source of energy, Calvin and co-workers were able to synthesize (in a model of a paleoatmosphere consisting of carbon dioxide, water and molecular hydrogen) some organic molecules including formaldehyde, which is relevant in the synthesis of sugars (ribose) and glycerol (cf., Sec. 2.1 and Garrison *et al.*, 1951). Other significant successes were the synthesis of amino acids (Miller, 1953), nucleotides (Oro, 1961b; Ponnampereuma *et al.*, 1963), and lipids (Oro, 1994). Another early success was the synthesis of protenoids by thermal heat in possible primitive Earth conditions (Fox and Harada, 1958). In other words, all the basic building blocks of life can, in principle, be synthesized in the laboratory. The question of how such synthesis may have occurred in the early Solar System is not an idle one:

Firstly, four satellites with atmospheres resembling the Earth's paleoatmosphere exist in the Solar System. Io is the first (closest) of the Galilean satellites of Jupiter (which has altogether

16 satellites). Io has a thin atmosphere of sulphur dioxide. The second Jovian moon, and the smallest of the Galilean satellites) Europa has some oxygen in its atmosphere (Hall *et al.*, 1995). Titan, perhaps the most intriguing component of the Solar System (cf. below), is a satellite of Saturn. It is surrounded by mainly molecular hydrogen atmosphere, but there is also a substantial component of methane. Finally, the last satellite known to have an atmosphere is Triton, the largest of the eight known moons of Neptune. It has a tenuous atmosphere with surface pressure of 15 microbars, mainly of nitrogen but with a trace of methane. It is remarkable that one of the components of the model paleoatmospheres of the chemical evolution experiments (ammonia) does not seem to be present in appreciable quantities in any of the satellites that have an atmosphere.

Secondly, chemical evolution is turning into an important chapter of the space sciences. This development may be illustrated with the questions raised by current studies of Titan. In the atmosphere of this satellite, the largest of Saturn's 18 confirmed satellites (bigger than even Mercury), biopolymers may possibly arise in this planet-like satellite. Indeed, under the action of sunlight it is possible that there may be liquid bodies of mixtures of ethane or methane on its surface. The Cassini-Huygens mission, under preparation at present, will attempt to obtain more reliable information on Titan. It includes a spacecraft (Cassini, built by NASA) bearing the atmospheric probe Huygens (built by the European Space Agency). The mission is due for launch in October, 1997 (Raulin *et al.*, 1995).

1.3. CHEMICAL EVOLUTION IN THE EARLY EARTH ATMOSPHERE

Chemical evolution on Earth seems to have taken place from 4.6-3.9 thousand million years before the present (Gyr bp), a geologic time interval which is known as the Hadean subera within the Archean era. It should be noticed, however, that considering the impacts by large asteroids in the early Earth does not necessarily exclude the possibility that the period of chemical evolution may have been considerably shorter. Indeed, the Earth may have been continuously habitable by non-photosynthetic ecosystems from a very remote date, namely since approximately 4.44 Gyr bp (Sleep *et al.*, 1989). In subsequent suberas of the Archean (3.9-2.5 Gyr bp) life, as we know it, certainly was present. This is well represented by fossils of the domain Bacteria, which is particularly well documented by many species of the phylum Cyanobacteria (Schopf, 1993).

Once again, it should be kept in mind that the content and the ratios of the two long-lived isotopes of reduced organic carbon in some of the earliest sediments (retrieved from the Isua peninsula, Greenland, some 3.8 Gyr bp), may convey a signal of biological carbon fixation (Schidlowski, 1995). This reinforces the expectation that chemical evolution may have occurred in a brief fraction of the Hadean subera, in spite of the considerable destructive potential of large asteroid impacts during the same geologic interval in all the terrestrial planets, as well as on the Moon.

2. The search for the origin of homochirality

2.1. WHAT IS HOMOCHIRALITY ?

This broad knowledge of early life has allowed a search for relics that may show the common thread of evolution from our last common ancestor. At the beginning of the 1990s Abdus Salam's deep interest in the role of asymmetry in the living state gave a strong impulse to the topic of the asymmetry of the biomolecules (Salam, 1991, Chela-Flores, 1991).

The first unifying principle in biochemistry (Crick, 1981) is that the key biomolecules have the same handedness: this phenomenon occurs when molecules are asymmetric, in such a

manner that they are able to exist in two configurations, both partners ('isomers') being mirror images of each other. These two possibilities are referred to, according to the response of a mixture of them to the incidence of a plane-polarized light. More precisely, the angle α through which the plane of polarization is rotated can be given a sign as follows: when looking towards the incoming beam of light, for a clockwise rotation of α , we would assign the term 'right-handed', or dextrorotatory (D); if such a rotation is anticlockwise we say that the rotation is 'left-handed', or laevorotatory (L).

Molecules that respond to beams of light in the above-mentioned manner are said to be *optically active*. Examples of single-handed biomolecules that will concern us in this review are the monomers of proteins, which are only L-amino acids, while the ribose and 2-deoxyribose monomers of the nucleic acids are all in the D-configuration. Phospholipids are no exception. These are biomolecules of the cell membrane having a phosphate group and one or more fatty acids. Indeed, the plasma membrane of the organisms of the domains Bacteria and Eukarya contain exclusively D-glycerol (Kandler, 1995).

Thus, the key biomolecules have the same handedness, or, we say more often that biomolecules have the same 'chirality' or, preferably that they are 'homochiral'. (These words are taken from the Greek language, as *cheir* means hand.) Finally, an equimolar mixture of both isomers is called a 'racemic mixture'. Chiral molecules have non-superposable D, or L three-dimensional mirror image structures or 'enantiomers' (once again, these are words derived from the Greek *enantios morphe*, whose meaning is 'opposite shape').

It is important to distinguish between the truly chiral influences (such as the electroweak interaction, cf., Sec. 2.3), from other physical effects that have been proposed since Pasteur's famous resolution of tartaric acid crystals. (These are crystalline naturally occurring carboxylic acids, with melting point of 170C, which Pasteur showed to be optically active). In fact, Barron defines a truly chiral system, as one in which there are two enantiomers that may be interconverted by space inversion, but not by time reversal (Barron, 1986). Optical rotation changes its sign under parity reversal, but not under time reversal, therefore satisfying Barron's criterion. The electroweak interactions provide an example of a truly chiral influence, and for this reason in this review we shall restrict our attention to this elementary interaction (cf., Sec. 2.3).

However, under kinetic control irreversibility destroys T-invariance, a situation which allows further possibilities in the context of the life sciences, since life itself is to be understood as a non-equilibrium phenomenon.

Homochirality is valid for all organisms, but some care must be taken with the concept of homochirality as we consider the highest taxa:

2.1.1. Exceptionally, in the domain Bacteria (encompassing thermotogales, the flavobacteria and relatives, the cyanobacteria, the purple bacteria, the Gram-positive bacteria, and the green nonsulfur bacteria) cell walls may contain D-amino acids, as in the case of *Bacillus brevis* or *Lactobacillus arabinosus*.

2.1.2. A second exception concerns another one of the three highest taxa, the domain Archaea encompassing the kingdoms Crenarchaeota (which, for instance, includes the genus *Pyrodictium* and the genus *Thermoproteus*) and Euryarchaeota (for instance, the extreme halophiles and archaeobacteria capable of producing methane as a byproduct of the reduction of carbon dioxide). Organisms in this domain contain L-glycerol in their membrane phospholipids, instead of the standard D-glycerol, as we mentioned above, which is characteristic of the phospholipids of the cellular membranes of the other two domains, Eukarya (the animals, the ciliates, the green plants, the fungi, the flagellates and the microsporidia) and Bacteria.

2.2. IS THE ORIGIN OF HOMOCHIRALITY EXTRATERRESTRIAL?

The property of homochirality is particularly relevant in the identification of microorganisms for the *in situ* search for extraterrestrial homochirality (SETH). This recent research proposal may eventually be implemented by means of an apparatus denominated the 'SETH cigar' (MacDermott and Tranter, 1995). The most useful physical techniques that may be used in this context are circular dichroism (CD) and optical rotatory dispersion (ORD, the functional relationship between the angle of rotation and the wavelength of the polarized light). Normally these techniques imply equipment that is too bulky. Both techniques, CD and ORD may allow the determination of the sign and magnitude of the optical rotation to be measured, as well as the identity of the putative homochiral molecule.

MacDermott and Tranter propose instead to miniaturize the equipment by just restricting the search to the optical rotation that would be produced by a chiral molecule. Their idea is based on the property of some crystals (dichroism) of being able to absorb selectively light vibrations in one plane, but allowing the vibrations at right angles to go through. Two polarized films at right angles are introduced in the presence of small electronic devices consisting of regions (p-n junctions) in which on semiconductors are present on either side, so that their conductivity is due to the flow of electrons ('n-type') and mobile holes ('p-type'), respectively. This device is called a light-emitting diode. In addition, a detector is used, which consists of a semiconductor diode that produces a significant photocurrent when illuminated (photodiode). With this arrangement no light should reach the detector unless the plane of polarization is rotated. In this manner racemic mixtures will not be capable of allowing light to reach the detector. Hence a signal in the detector is certain proof that the sample tested is homochiral.

Homochirality may be considered as a signature for life anywhere in the universe. Due to the progress in research and the availability of a whole series of forthcoming space missions (towards the end of this decade and the beginning of the next), the problem of chemical evolution has been virtually transferred from the laboratories of organic chemistry to the space sciences.

In this context we may recall the intriguing ideas of Bonner and Greenberg. These authors suggest, independently, that circularly polarized light from neutron stars on passing clouds of interstellar dust may selectively eliminate one or the other enantiomer in the dust mantle, if mirror image molecules are originally present (Bonner, 1991, Greenberg *et al.*, 1995). Another possible astrophysical source of homochirality may be the supernovas, according to some recent work (Cline *et al.*, 1995). We may state in general that living systems translate their genes, following the rules of the genetic code, into proteins composed of relatively a small number of L-amino acids. The strength of this statement may be illustrated with the discovery that out of the 74 amino acids that were found in the Murchison meteorite (cf., Sec. 1.1), only 55 have been found in extraterrestrial samples and none of them has been shown conclusively to have a chiral bias (Oro *et al.*, 1990). These numbers give us considerable insights, when we recall that the protein monomers are chosen from a very restricted set of twenty amino acids.

2.3. HOMOCHIRALITY AND THE ELECTROWEAK INTERACTION

The search for the origin of homochirality from the point of view of the effects of the weak interactions started in the 1950s. This has led to the proposal of various mechanisms that have been postulated in order to account for homochirality. Amongst them we should recall the

differential decomposition and resolution of enantiomers under beta-irradiation (the Vester-Ulbright hypothesis). The experimental confirmation of this theoretical conjecture has been much discussed over the last three decades (Garay, 1993). More ingenious mechanisms have been recently postulated, such as the possible relevance of molecular decay in the origin of homochirality (Li, 1994).

In the standard model of Glashow, Salam and Weinberg, the electroweak interactions arise by the exchange of photons, massive charged W^{+-} and neutral Z^0 bosons of spin 1 between quarks (u,d,s,c,b,t) and leptons (electron, muon, the tau particle and three types, or 'flavours', of neutrinos). Within the framework of this model, a substantial body of previous work has suggested the parity-violating electroweak neutral current (mediated by the Z^0 boson) as the main physical force inducing the observed homochirality (Keszthelyi, 1995). Neutral current weak interactions are predicted to induce small energy differences, equal in magnitude but opposite in sign between the enantiomers. Such a 'parity-violating energy difference' between the enantiomers (twice the small induced energy difference in each enantiomer) may induce energy shifts to both enantiomers, which in spite of being small effects they are, nevertheless, subject to calculation as well as to the eventual effect of amplification mechanisms (MacDermott, 1995). We shall return to this topic more fully in Sec. 5.1.

Salam's approach (1991) also invoked the electroweak interaction, but was original in appealing to further physical concepts that may apply during chemical evolution. He pointed out a particular cooperative phenomenon that may have taken place, namely, a phase transition beneath a certain critical temperature T_c . Amino acids that had been synthesized earlier from various precursors entered into a new phase assumed to be a Bose condensed mode.

These concepts: cooperative phenomena and condensation led to the suggestion that the electromagnetic force is not the only force producing chemical effects. The Z^0 component of the electroweak force, in spite of the fact that its effects appear to be negligible at low energies, may play an active role in biochemistry. The reasons for the proposed chemical role of the parity-violating weak interactions may be found in quantum chemistry calculations (QCCs) of, amongst others, Mason, MacDermott and Tranter (MacDermott, 1993) and, independently, by Bakasov, Ha and Quack (Bakasov *et al.*, 1996). We refer the reader to the reviews of Bonner (1991), Cline (1996) and Keszthelyi (1995) and for a comprehensive list of earlier references.

These calculations indicate that four amino acids, for which results have been obtained, are L-stabilized relative to their D-partners for configurations in aqueous media. (The above-mentioned amino acids are alanine, valine, serine, and aspartic acid.) Although the result of the calculations give only a small energy shift in the ground state energies of the L- and D-amino acids, nevertheless, in the conjectured phase transition, sufficiently large effects may still have to be taken into account, such as the inclusion of the top quark (t) and its antiparticle, in an enlarged version of the standard model (Chela-Flores, 1991).

2.4. PHYSICAL CRITERIA FOR THE GENESIS OF HOMOCHIRALITY

An important aspect of the origin of homochirality concerns the thermal conditions that may have existed at a time when chemical evolution took place. Although the geological record is unknown on Earth, as metamorphism has destroyed all evidence, the corresponding period in Moon stratigraphy is well documented (the Nectarian, 4.2-3.8 Gyr bp), when presumably the presently observed homochirality of amino acids originated on the Earth-bound biomolecules. Solid evidence of the heavy bombardment period is known from the rocks

retrieved by the Apollo missions. (Even samples from the pre-Nectarian period are known, as the Apollo 17 mission succeeded in retrieving a dunite-an intrusive igneous rock of ultramafic chemistry-some 4.4 Gyr bp.)

Since the ambient temperature of the Earth's surface is approximately 300K, if T_c is much smaller than this temperature, Salam's theory may not apply to the Earth, in which case some options are still open:

2.4.1. Pre-solar contributions may be necessary to get low enough temperatures. We may assume that such contributions may have arisen from a molecular cloud of dust and gas (predominantly in molecular form) out of which the Solar System originated. Since over seventy organic molecules have been detected in the last 25 years (cf., Sec. 1.1), it is possible that the effects that we are discussing in this review may have taken place in amino acids at a time prior to the formation of the Solar System. In those early stages the temperatures may indeed have been low enough (possibly in the range 10-20K, as in current small molecular clouds).

2.4.2. Major contributions come from the outer Solar System with ambient temperatures less than T_c . (For instance, the surface temperature of Uranus, which is some three billion km from the Sun, is well below -200C, a temperature in which both methane and nitrogen condense.) In the heavy-bombardment period (i.e., prior to about 3.8 Gyr bp) the Earth and Mars, for instance, may have exchanged matter and if life originated in Mars first, it is conceivable that it may have been transferred to the early Earth. So the assumption that chemical evolution of the Earth biomolecules may have been influenced by interplanetary exchange of chemicals cannot be ruled out.

2.4.3. If T_c is found to be considerably less than 300K, one would be led to accept that the Earth only acted as a junction place, where L-amino acids came together with D-sugars, so as to give rise to early replication. This option is in agreement with the Oro hypothesis (cf., Sec. 1.1).

It may be reasonable to assume that the Earth did accrete prebiotic organic molecules important for the origin of life from impacts from carbonaceous asteroids and comets during the period of heavy bombardment in the Hadean Earth (Chyba *et al.*, 1990). Remarkably, we may also assume that the alanine molecules, for instance, could withstand temperatures as high as about 700K for one second, whereas the other amino acids could withstand temperatures in the range 600 to 800K for a similar period of time, and remain stable and intact upon impact with the Earth.

Another physical question is whether phase transitions of the type of Bose condensation may occur in the living cell. We may recall that in the case of certain specific macromolecules relevant to living processes, this subject has been studied in both biochemistry, as well as in genetics, during the last thirty years.

In biochemistry, the ability of lipids to adopt a variety of phases is well documented (Luzzati, 1968; Eze, 1995). This has been called lipid polymorphism; an important aspect of this topic is that the macroscopic structure adopted by lipids depends on the experimental conditions; for instance, temperature is an important parameter which determines the macroscopic structure of hydrated membrane lipids (De Kruijff *et al.*, 1985). At present a conservative position is still that the possible biological significance of some of the polymorphic membrane transitions is striking, but we have not yet bridged the gap between these transitions and physiological phenomena.

On the other hand, evidence for phase transitions in the life sciences is not limited to biochemistry, but a separate line of research in genetics also suggests the occurrence of phase transitions in the living state: A factor that may influence the onset of DNA replication is intracellular *ion concentration* (Maino *et al.*, 1974; Dulbecco and Elkington, 1975). In fact,

experiments with rat liver-cell nuclei may indicate that chromatin structure and nuclear volume display abrupt transitions as function of ion concentration in the nuclear environment (Nicolini *et al.*, 1984): this experiment has suggested to its authors that the chromatin structural changes may be discussed in terms of phase transitions.

The phenomenon being proposed may be thought of as a quantum mechanical enhancing factor in molecular evolution from the initial racemic mixture to the present day chiral amino acids. The concept of condensation is not new in biology; indeed, it goes back some 30 years (Delbruck, 1963). In the context of the origin of life condensation in the nucleic acids had already been suggested to have played a role in the origin of Darwinian evolution (Chela-Flores, 1985).

3. Experimental tests of the putative superconducting-like phase transition in amino acids.

3.1. WHAT DIRECT EVIDENCE IS THERE FOR A CLEAR HALLMARK OF A PHASE TRANSITION THAT MAY UNDERLY THE ORIGIN OF HOMOCHIRALITY?

These results from the biochemistry of the cell membrane and from the genetics of chromatin structure lead us naturally to raise the question: What direct evidence is there for a clear hallmark of the phase transition underlying the origin of chirality? There are some alternatives:

3.1.1. Melanin - a dark brown pigment of many animals- is the product of tyrosine metabolism, which causes pigmentation of eyes, skin and hair in vertebrates. It is often located in melanophores (cells with permanent radiating processes lying superficially in vertebrates). Data taken from natural melanins, as well as from melanosomes isolated from human melanoma, may indicate that it undergoes a phase transition. In fact, its heat capacity shows a discontinuity near 1.9K. This anomaly, though relatively small, is significant beyond the scatter of the data points. It is probably associated with a magnetic transition, possibly from paramagnetism to antiferromagnetism (Mizutani *et al.*, 1976). In an analogous manner, a means of detecting the putative phase transition may be by measuring differences of specific heats and looking for anomalies in the curve for specific heat (a cubic polynomial) as has already been done for the above-mentioned non-amino acids.

3.1.2. One direct means of testing the new theory for the origin of chirality in amino acids is by taking a 50-50 racemic mixture of crystalline L- and D-amino acids and lowering the temperature. Salam has discussed extensively an experiment involving a 50-50 racemic mixture subjected to a temperature gradient ranging from the melting temperature for the amino acids to the absolute zero temperature.

3.1.3. In view of the discovery of organic superconductors with critical temperatures similar to metals and alloys, an alternative experimental procedure may be to test directly for conductivity and magnetic (Meissner effect) properties of amino acid crystals.

3.2. PHASE TRANSITIONS IN ORGANIC MATERIALS

The suggestion has been made that a superconducting-like phase transition may actually occur in amino acids. Physics has been rather slow in studying alternative candidates for superconductors. The first class of (low-temperature) superconductors was discovered in 1911; these early superconductors were unlike any form of matter that occurs in living systems. In fact, they were metals and alloys. The discovery of a radically new class of

superconductors had to wait some time. In 1986 a more interesting superconductor from the point of view of chemistry was identified in certain ceramics

The simple potassium-doped fullerene (Hebard *et al.*, 1991) has been shown to be superconducting with a reasonably high critical temperature i.e., the onset of the critical temperature is 18K. However, we should recall that organic superconductors have been known for some time, such as graphitic compounds with typical formulae C_8A ($A = K, Rb, Cs$) with values of T_c smaller than 1K (Hannay *et al.*, 1965).

Another example is that of the organic superconducting metal bis(ethylene dithiolo) tetrathiofulvalene triiodide with formula $(C_{10}H_8S_8)_2I_3$ (Yagubskii *et al.*, 1984) with a value of $T_c = 1.5K$ at normal pressure. Hence, the most we can say at present is that superconductivity-, or superfluidity-like effects may not be excluded from organic materials. This suggests to extend the search for quantum liquid behaviour observed in the organic materials to a particular form of organic materials, namely, amino acid crystals. This point will be discussed further in Sec. 5.5.

3.3. BIOMOLECULAR RELICS OF EVOLUTION

The search for physical mechanisms underlying the causal origin of homochirality are mainly motivated by the fact that the most characteristic of all relics of evolution is provided by the chiral nature of the biomolecules.

Other relics, however, are known. This may be illustrated with the widespread use of the standard genetic code by a very large number of organisms. (Exceptionally some protozoans and *Mycoplasma* use a modified code.)

The general question of whether it is possible for further relics of the earliest stages of evolution to have survived till the present has been raised in the past by various authors. Some well known examples are certain nucleotide sequences (introns) in genes of mainly eukaryotic organisms; they do not code for the gene product (neither proteins nor nucleic acids). The presumed antiquity of these sequences is based on the finding that some introns are ancient (Saldanha *et al.*, 1993). More recently, the question of further molecular relics from one of the earliest stages of the evolution of life on Earth (the RNA world) has been raised in the context of the origin of RNA plant pathogens, such as the viroids (Diener, 1993). The relevance of this putative relic has been discussed recently (Chela-Flores, 1994b; 1995).

Inserted in this broad range of problems regarding biomolecular relics, the all-important question of homochirality of amino acids as an evolutionary relic has been recently reviewed, particularly from the point of view of experiments (Bonner, 1994, Keszthelyi, 1995, Cline, 1995), while the relevance of chirality in biology has also been discussed extensively in an excellent review (Crossly, 1992).

The continued efforts to come to terms with the experimental question of Salam's two conjectures (cf., Sec. 5.3 below), have not been completed so far, but these set of experiments (Figureau *et al.*, 1993, 1995; Navarro-Gonzalez *et al.*, 1993; Wang *et al.*, 1995) are, nevertheless, illuminating an area of condensed matter physics which has so far not been extensively studied, and which potentially may be of wider interest than strictly within the domain of chemical evolution and the origin of life.

The preliminary tests of Wang and co-workers in Beijing (1995) have faced the first part of the task (to test for the specific heat curves; these experiments await confirmation and should be repeated), but the complementary magnetic properties should also be urgently explored and tested by independent groups. The intrinsic importance of testing new materials for the key technologies of the future should be emphasized, as technological innovation should not

stop at ceramics, which already produced the revolution of the high T_c superconductors in 1986 (Chela-Flores *et al.*, 1988), but biological materials should be thoroughly discussed and tested, as we already emphasized in our earlier paper (Chela-Flores, 1991).

4. Asymmetric adsorption of amino acids at crystal surfaces

4.1. L-AMINO ACIDS MAY BE ADSORBED ON L-QUARTZ CRYSTALS

There is a long history of experimental efforts with inorganic substances which have the ability to rotate the plane of polarization of light passing through a crystal (cf., Sec. 2.1). This ability is due to their chiral crystal structure. In this respect quartz, the most common of the six polymorphs of crystalline silica (SiO_2) is found in nature in the form of well defined enantiomorphous crystals, in which chirality is caused by the spiral structure of linked silicon-oxygen tetrahedra.

Quartz, therefore, is an obvious candidate for the stereoselective adsorption of a given enantiomer from a racemic mixture. A long series of experiments has led to the conclusion that asymmetric adsorption on quartz is a well authenticated phenomenon, but it is also equally clear that in this process, quartz acts merely as a passive 'filter' of homochiral domains, passive in the sense that quartz itself does not induce any catalytic process that may convert racemic or achiral reactants into optically active products (Bonner, 1991).

Asymmetric adsorption of amino acids at the surface of chiral crystals is a possible physical mechanism that may have led to the onset of homochirality in one class of biomolecules. It is well documented that about one per cent more L-amino acids than D-amino acids are adsorbed on L-quartz crystals (Bonner, 1979).

This experimental fact may be taken together with the theoretical work that has shown that due to the effect of the weak interactions, L-amino acids (and similarly, D-sugars) have smaller ground-state energy than their corresponding enantiomers, a property that has been shown by independent QCCs (cf., Sec. 2.3). The experiments on the stereoselective adsorption of amino acid enantiomers on the surface of either D-quartz or L-quartz suggest (together with the QCCs) that L-amino acids (which may be adsorbed in the surface of L-quartz) may serve as a reservoir of chirally-pure samples. Some amplification mechanism may have acted on, so as to enhance the small chiral bias that the QCCs suggest in favour of L-amino acids.

The analogous D-amino acids (that may be similarly adsorbed by the surface of D-quartz), presumably may not have benefitted from the small advantage due to the weak interactions in favour of the L-amino acids. Hence, it is forced upon us to discuss possible physical mechanisms that while acting on both L- and D-amino acid crystals may, nevertheless, act as amplification mechanisms on chirally-pure samples.

4.2. THE POSSIBLE RELEVANCE OF QUARTZ IN CHEMICAL EVOLUTION

The abundance of quartz and its presence in a variety of conditions typical of the Archean eon, militate in favour of the possibility that quartz may have provided early chirally-pure environments for amino acids.

We shall consider a few arguments that favour a physical mechanism in which L-quartz is the first step in the pathway that leads to the observed homochirality of protein amino acids: the environment in which quartz is found is ubiquitous. Although quartz is one of the thousand silicate minerals that make up the bulk of the Earth's outer crust, it is however one

of its commonest minerals (12 % by volume). Furthermore, all terrestrial planets have a crust similar to the silicate crust of the Earth. In the case of Mars, for example, samples ('fines') have been analyzed by the Viking missions. It was demonstrated that quartz is a prominent mineral in the fines that were retrieved from both the Chryse and the Utopia Planitiae (cf., Soffen, 1976, and the articles that follow his summary). Quartz is known to crystallize directly from igneous magma (i.e., from molten rock material). It occurs in deep-seated coarse-grained ('plutonic') rocks such as granites, as well as in volcanic rocks such as rhyolites, which are igneous rocks containing megascopic crystals of quartz.

An additional reason to focus our attention on quartz is that it is stable in sedimentary conditions: it may occur in marine and desert sands, but also as cement in compacted rocks such as sandstone, in which quartz appears as rounded grains between the diameters of 0.06 and 2 mm, with a variable content of other mineral grains. What seems even more pertinent is that quartz is also stable under low- and high-grade metamorphic conditions. In fact, a high-pressure, high-temperature form (coesite) is found in meteorite impact rocks. Another form (stishovite) is found to occur naturally in meteorite craters, such as Meteor Crater, Arizona.

Hence the probable presence of quartz in favourable environments during the heavy-meteorite impact-period (the Hadean) may have been a factor influencing the origin of life in the selection homochirality, which has persevered as a distinctive feature of the earth's biota.

5. On the breaking of chiral symmetry

5.1. AMPLIFICATION MECHANISMS

One of the principal difficulties in understanding the asymmetry of biomolecules is the question of the amplification mechanism that would boost a primordial small physical effect (Keszthelyi, 1995). New suggestions of amplification mechanisms should be considered in addition to the earlier possibilities such as those considered in the past:

5.1.1. Amplification of the Vester-Ulbricht processes, namely amplification mechanisms that may apply to the interaction of polarized particles with biomolecules.

5.1.2. Amplification of the Yamagata processes, namely those processes (such as those discussed in this work) in which the breakdown of the discrete P symmetry is considered as a possible source of biochirality.

5.1.3. Amplification known as the Kondepoudi-Nelson catastrophic mechanism (Kondepoudi and Nelson, 1985), According to this proposal the small electroweak energy difference between the enantiomers is sufficient to determine which of the two single-enantiomer reaction sequences was adopted at the metastable stage in a flow reactor that is off thermodynamic equilibrium.

The suggestion of Salam that Bose-Einstein condensation may act as a novel amplification mechanism still suffers from a difficulty: the underlying theory requires that large activation energy barriers be overcome in the process of turning a racemic mixture into optically pure amino acids. This requires that the phase transition may induce molecular changes that lead to the breaking of chemical bonds. Amongst various options that the phase transition hypothesis may be faced with is that the origin of life may be extraterrestrial, provided that the critical temperature is too small for the phase transition to have been supported by the environment of the early Earth.

5.2. DIFFERENT MECHANISMS FOR IMPLEMENTING PARITY VIOLATION

The possibility of an alternative model incorporating P violation of a third kind has been suggested in the past (Chela-Flores, 1994a), in which the phase transition acts on a racemic mixture of amino acids inducing spontaneous separation of the chiral phases due to their environment. The ground states of the two optical pure phases are taken to differ in energy (the L-amino acid having the lower ground-state energy), as discussed in Sec. 2.3.

In fact, it has been found that a small electroweak energy increment adds to the binding energy of a given state in one enantiomer and subtracts from that of the corresponding state of the other, giving an electroweak binding energy difference between the two optical isomers.

This rationalizes the observed bias of protein amino acids for the left-handed chirality, a phenomenon that has been preserved by biological (i.e., neodarwinian) evolution. The interpretation of the postulated phase transition provided in this note does not suffer from the difficulty of implying large activation energy barriers for the production of optically pure amino acid concentrations from an initial primordial racemic mixture.

If, on the other hand, extraterrestrial origin of homochirality is envisaged (cf., Sec. 2.2), a more compelling argument in favour of the present proposal is that, as pointed out above, molecules behave rather differently from macroscopic systems in the sense that, there are no sharp transitions (such as those considered in the original Salam proposal) between symmetric and asymmetric states in (free) molecules.

The BCS-like phase transition in the amino acids may occur at a certain critical temperature T_c . Preliminary experiments reporting negative results (cf., Sec.3.3) have attempted to find the value of T_c for the original theory. This situation motivates the search of further insights into the sources of homochirality. We suggested that one possible insight may be found in the connection between chiral symmetry breaking and pattern formation.

It has been observed that two-dimensional liquids composed of achiral units can spontaneously separate into right- and left-handed regions. This induces a breakdown of natural mirror symmetry, an effect which is related to ferroelectricity at the interfaces (Schwartz, 1993).

However, one of the important aspects of these experiments is that a new example has been provided, in which structure and symmetry at the molecular level have implications at the macroscopic level. This phenomenon has been observed at molecular length scales in some experiments (Eckhardt *et al*, 1993). The molecular lattice of a racemic organic monolayer has been imaged and was observed to spontaneously undertake separate chiral regions.

5.3. NOVEL AMPLIFICATION PROPOSALS

We assume that the connection between chiral symmetry breaking and pattern formation may occur in bulk three-dimensional systems such as amino acids. A temperature gradient would induce a racemic mixture of amino acids, through the Bose-Einstein condensation into a regime of spontaneously broken mirror symmetry, in which the chiral phases may be separated. This, we conjecture, should manifest itself physically through pattern formation.

The question of the amplification of the small effect of the electroweak interaction which gives the preferential stability to the L-amino acid, becomes a relevant problem due to the smallness of the coupling constant of the electroweak interaction. We shall refer to the corresponding parity-violating energy-difference, which constitutes a basic concept in the postulated nonlinear mechanism for amplification.

Salam made two different proposals, a point that has not been widely appreciated (Chela-Flores, 1994a):

5.3.1. He proposed the existence of a phase transition in a *racemic mixture* of amino acids in which large activation energies need to be overcome. This proposal does not refer to the environment in which the racemic mixture was inserted. This is the aspect of Salam's conjecture that was tested in two experiments (cf., Sec 4.1).

5.3.2. Salam went further in a separate proposal of a new mechanism for the amplification of the small effect of the neutral electroweak interactions on the ground state energy of the amino acids. This possibility was suggested on the bases of the QCCs (cf., Sec. 2.3). The macroscopic quantum mechanical amplification mechanism is analogous to Bose-Einstein condensation, currently assumed to occur in other forms of condensed matter, such as in liquid helium II and superconductors. It is this aspect of Salam's conjectures that has been preliminarily tested in the most recent experiment, which clearly requires independent confirmation (Wang et al., 1995).

The proposal (5.3.1) has the difficulty that large activation energies have to be overcome, so that chemical bonds are broken for a given right-handed amino acid to be converted to its left-handed form. The second proposal (5.3.2) will be discussed below.

5.4. ON SOME EXPERIMENTS TESTING THE QUESTION OF HOMOCHIRALITY

Unlike Salam's first suggestion which referred to a racemic mixture (Salam, 1990, 1991, 1992, 1993, 1995), we advocate the second part of his proposal (5.3.2). The main reason is the evident considerable difficulty in explaining the means of overcoming the large activation energies which would inhibit the phase transition in the racemic mixture of amino acids.

At present, in order to test the theory, one needs to address the question of the possible occurrence of a phase transition in a chirally-pure sample of L-amino acids. For instance, experiments could address the question of chirally-pure samples of amino acids that have been adsorbed preferentially on the surface of L-quartz crystals, as explained in the above illustration (cf., Sec. 4).

In spite of the difficulties in the original phase-transition theory of 1991, non-linear mechanisms of amplification of the small parity-violating energy-difference may still be relevant, but one should search those theories that are free from the considerable difficulty implied by the underlying large activation energies needed in the transition from the racemic mixture to the chirally-pure samples. One possibility of an acceptable phase transition theory, free from the above-mentioned difficulties, may be found in the generation of chirally-pure domains by the action of L-quartz surfaces, a process that was already discussed in Sec. 4.2.

5.5. CHIRAL AMINO-ACID CRYSTALS AND SUPERCONDUCTIVITY

Evidence for the phase transition in chirally-pure samples of L-amino acids should be searched for with similar experiments to those preliminary experiments which tested the possible existence of the phase transition, in which a range in temperature was scanned for the signature of the putative phase transition (cf., paragraph 5.3.1).

On the other hand, some evidence that should be kept in mind is that Salam and the present author (Salam, 1991, Appendix D; Chela-Flores, 1991) were already aware that in the special case of melanin, a derivative of the amino acid tyrosine, a phase transition may already have been detected experimentally (cf., Sec. 3.1). A similar experiment directly on a sample of the amino acid L-tyrosine would serve as a test of the conjectured unusual behaviour of amino acids. Further, the importance of testing for high-temperature superconductivity, in the present case of biological relics of the earliest steps in chemical evolution (homochirality in protein amino acids), was pointed out in the same above-mentioned section of our 1991 paper.

Having raised the question of spontaneous onset of homochirality, experimental attention finally seems justified at present to search for signs of a phase transition in single chirally-pure crystals of amino acids. As stated in Sec. 5.2, this mode of symmetry breaking should be distinguished from either Quack's *de facto* symmetry breaking (the ground state does not show the same symmetry as the hamiltonian), or from *de lege* symmetry breaking (parity violation occurs in the hamiltonian, as in the case Salam discussed in his 1991 paper). For more details of the first two types of parity violation we refer the reader to the original paper (Quack, 1993).

6. Conclusions

The postulated parity violation of a third kind (in which the homochiral domains may form spontaneously due to their environment) may be illustrated in the case of the chiral quartz crystals discussed in Sec. 4, or in the case of phenomena that is known to occur in the liquid crystal state (cf., Sec. 5.1). From what we have said above we feel that preliminary experiments testing parity violation of the third kind (such as the Wang *et al* experiments) are finally justified (i.e., tests of phase transitions in single amino acids).

In this manner we have been led naturally to suggest testing a form of condensed matter that has so far, from the present point of view, escaped the attention of experimentalists (although considerable information already exists on the crystal structures of amino acids). In this context it may be worth recalling the lesson learnt when ceramics surprised the scientific community by showing two aspects that normally complement the phenomenon: superconducting currents (with the associated signature in the specific heat curves, as well as the Meissner effect).

ACKNOWLEDGEMENTS

The author would like to thank Professor John T. Lewis for hospitality at the School of Theoretical Physics of the Dublin Institute for Advanced Studies, where the present work was completed.

References

- Bakasov, A., Ha, T.-K. and Quack, M. (1996). *Ab initio* calculation of molecular energies including parity violating interactions. In: Chemical Evolution: Physics of the Origin and Evolution of Life. Chela-Flores, J. and Raulin, F. (Eds.). (1996). (Proc. of the Fourth Trieste Conference on Chemical Evolution: The Cyril Ponnampereuma Memorial Conference, Trieste, Italy, 4-8 September, 1995). To be published by Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Barron, L.D. (1986). True and false chirality and parity violation. *Chem. Phys. Letters*. 221, 311-316.
- Bonner, W.A. (1979). Experiments on the abiotic origin and amplification of optical activity. In Walker (1979). pp. 5-20.
- Bonner, W.A. (1991). The origin and amplification of biomolecular chirality. *Origins of Life and the Evolution of the Biosphere* 21, 59-111.
- Bonner, W.A. (1994). Enantioselective autocatalysis-Spontaneous resolution and the prebiotic generation of chirality. *Origin of Life and the Evolution of the Biosphere* 24, 63-78
- Chela-Flores, J. (1985). Evolution as a collective phenomenon, *J. Theor. Biol.*, 117, 107-118.
- Chela-Flores, J. (1991). Comments on a novel approach to the role of chirality in the origin of life, *Chirality*, 3, 389-392
- Chela-Flores, J. (1994a). The origin of chirality in protein amino acids. *Chirality* 6, 165-168.
- Chela-Flores, J. (1994b). Are viroids molecular fossils of the RNA world?. *J. Theor. Biol.* 166, 163-166.
- Chela-Flores, J. (1995). Molecular relics from chemical evolution and the origin of life. In: *Chemical Evolution: Self-Organization of the Macromolecules of Life* Eds. J. Chela-Flores, M. Chadha, A. Negron-Mendoza, and T. Oshima. A. Deepak Publishing: Hampton, Virginia, USA. pp. 185-200.
- Chela-Flores, J., Das, M.P., and Saif, A.G. (1988). A Phenomenological Approach to High Tc Oxide Superconductors. *Solid State Commun.* 65, 77-80.
- Chyba, C.F., Thomas, P.J., Brookshaw, L., and Sagan, C. (1990). Cometary delivery of organic molecules to the early Earth. *Science* 249, 366-373.
- Cline, D. (1996). Determining the physical origin of homochirality in life. In: *Chemical Evolution: Physics of the Origin and Evolution of Life* Chela-Flores, J. and Raulin, F. (Eds.). (1996). To be published by Kluwer Academic Publishers, Dordrecht, The Netherlands.

Cline, D., Liu, Y. and Wang, H. (1995). Effect of a chiral impulse on the weak interaction induced handedness in a prebiotic medium. *Origins of Life and the Evolution of the Biosphere* 25, 201-209.

Crick, F. (1981). *Life Itself*. London: MacDonal & Co.

Crossly, R.(1992). The relevance of chirality to the study of biological activity, *Tetrahedron* 48, 8155-8178.

de Kruiff, B., Cullis, P.R., Verkleij, A.J., Hope, M.J. van Echteld, C.J.A., and Taraschi, T.F. (1985). Lipid polymorphism and membrane function. In: *The enzymes of biological membranes*. 2nd. ed. A. N. Martonosi (ed.). Vol 1, *Membrane Structure and Dynamics*. New York: Plenum Press. pp. 131-204.

Delbruck, M. (1963). In: *Commemoration of the 50th. Anniversary of Niels Bohr's Papers on Atomic Constitution, Session on Cosmos and Life*. Copenhagen: Institute for Theoretical Physics. pp. 41-67.

Diener, T.O. (1993). Small pathogenic RNAs of plants: Living fossils of the RNA world? In: *Chemical Evolution and the Origin of Life*. Eds. C. Ponnampereuma and J. Chela-Flores. Hampton, Virginia: Deepak Publishers.

Drake, F. (1996). Ongoing and future searches for extraterrestrial intelligent life. In: *Chemical Evolution: Physics of the Origin and Evolution of Life* Chela-Flores, J. and Raulin, F. (Eds.). (1996). (Proc. of the Fourth Trieste Conference on Chemical Evolution: The Cyril Ponnampereuma Memorial Conference, Trieste, Italy, 4-8 September,1995). To be published by Kluwer Academic Publishers, Dordrecht, The Netherlands.

Dulbecco, R. and Elkington, J. (1975). Induction of growth in resting fibroblastic cell cultures by Ca^{++} . *Proc. Natl. Acad. Sci. USA* 72, 1584-1588.

Eckart, C.J., Peachy, N.M., Swanson, D.R., Takacs, J.M., Khan, M.A., Gong, X., Kim, J.-H., Wang, J., and Uphaus, R.A. (1993). Separation of chiral phases in monolayer crystals of racemic amphiphiles. *Nature* 362, 614-616.

Eze, M. (1995). Membrane phase separations, asymmetry and implications in the origin of life. In: Chela-Flores, J., M. Chadha, A. Negron-Mendoza, and T. Oshima (Eds.).(1995). *Chemical Evolution: Self-Organization of the Macromolecules of Life*. A. Deepak Publishing: Hampton, Virginia, USA. pp. 131-137.

Figureau, A., Duval, E., and Boukenter, A., (1993). Search for phase transitions Changing Molecular Chirality. In: *Chemical Evolution: Origin of Life*. Eds. C. Ponnampereuma and J. Chela-Flores. A. Deepak Publishing: Hampton, Virginia, USA. pp. 157-164.

Figureau, A., Duval, E., and Boukenter, A. (1995). Can biological homochirality result from a phase transition?. *Origins of Life and Evolution of the Biosphere*. 25, 211-217.

Fox, S.W. and Harada, K. (1958). Thermal copolymerization of amino acids to a product resembling protein. *Science* 128, 1214.

- Garay, A.S. (1993). Theoretical and experimental studies of the possibility of chirality dependent time direction in molecules. In: *Chemical Evolution: Origin of Life*. Eds. C. Ponnampereuma and J. Chela-Flores. A. Deepak Publishing: Hampton, Virginia, USA (1993). pp. 165-179.
- Greenberg, J.M., Kouchi, A., Niessen, W., Irth, H., van Pardijs, J., de Groot, M., and Hermsen, W. (1995). Interstellar dust, chirality, comets and the origins of life: Life from dead stars? In: Ponnampereuma, C. and Chela-Flores, J. (Eds.). (1995). *Chemical Evolution: The Structure and Model of the First Cell* Kluwer Academic Publishers, Dordrecht, The Netherlands. pp. 61-70.
- Hall, D.T., Strobel, D.F., Feldman P.D., McGrath, M.A., and Weaver, H.A. (1995). Detection of an oxygen atmosphere on Jupiter's moon Europa. *Nature* 373, 677-679.
- Hannay, N.B., Geballe, T.H., Matthias, B.T., Anders, K., Schmidt, P., and MacNair, D. (1965). Superconductivity in graphitic compounds. *Phys. Rev. Letters* 14, 225-226.
- Hebard, A.P., Rosseinsky, M.J., Haddon, R.C., Murphy, D.W., Glarum, S.H., Palstra, T.T.M., Ramirez, A.P., and Kortan, A.R. (1991). Superconductivity at 18 K in potassium-doped fullerene K_3C_{60} . *Nature* 350, 600-602.
- Kandler, O. (1995). Cell wall biochemistry in Archea and its phylogenetic implications. In: Ponnampereuma, C. and Chela-Flores, J. (Eds.). (1995). *Chemical Evolution: The structure and model of the first cell*. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp. 165-169.
- Keszthelyi, L. (1995). Origin of the homochirality of biomolecules. *Quarterly Reviews of Biophysics* (In press).
- Kondepoudi, D.K. and Nelson, G.W. (1985). Weak neutral currents and the origin of biomolecular chirality. *Nature* 314, 438-441.
- Li, Th. (1994). A possible role of molecular decay in the origin of asymmetric emergence of biomolecules on earth. *J. Theor. Biol.* 170, 227-230.
- Luzzati, V. (1968). X-ray diffraction studies of lipid-water systems. In: *Biological Membranes*. Ed. D. Chapman. New York: Academic Press. pp. 71-12 .
- MacDermott, A. (1993). The weak force and the origin of life. In: *Chemical Evolution: Origin of Life* (Proc. Conf. Chemical Evolution and the Origin of Life, Trieste, Italy October 26-30, 1992). Eds. C. Ponnampereuma and J. Chela-Flores. A. Deepak Publishing: Hampton, Virginia, USA. pp. 85-99.
- MacDermott, A. (1995). The weak force and the origin of life and self-organization. In: Chela-Flores, J., M. Chadha, A. Negron-Mendoza, and T. Oshima (Eds.), 1995. *Chemical Evolution: Self-Organization of the Macromolecules of Life* A. Deepak Publishing: Hampton, Virginia, USA. pp. 237-249.

- MacDermott, A. and Tranter, G.E. (1995). Looking for the homochiral signature of life: the SETH Cigar. In: *Chemical Evolution: The structure and model of the first cell*. In: Ponnampereuma, C. and Chela-Flores, J. (Eds.). Kluwer Academic Publishers, Dordrecht, The Netherlands. pp. 77-81.
- Maino, V.C., Green, N.M., and Crumpton, M.J. (1974). The role of calcium ions in initiating transformation of lymphocytes. *Nature* 251, 324-327.
- Mason, S.F. (1992). *Chemical Evolution Origin of the Elements, Molecules, and Living Systems*. Clarendon Press: Oxford.
- Miller, S. (1953). A production of amino acids under possible primitive Earth conditions. *Science* 117, 528-529.
- Mizutani, U., Massalski, T.B., McGinnes, J.E., and Corry, P.M. (1976). Low temperature specific heat anomalies in melanins and tumor melanosomes. *Nature* 259, 505-507.
- Navarro Gonzalez, R., Khanna, R.K., and Ponnampereuma, C. (1993). Chirality and the origin of life. In: *Chemical Evolution: Origin of Life*. Eds. C. Ponnampereuma and J. Chela-Flores. A. Deepak Publishing: Hampton, Virginia, USA pp. 135-156.
- Nicolini, C., Carlo, P., Finollo, R., Vigo, F., Cavazza, B., Ledda, A., Ricci, E., and Brambilla, G. (1984). Phase transitions in nuclei and chromatin. *Cell Biophys.* 6, 183-196.
- Oro, J. (1961a). Comets and the formation of biochemical compounds on the primitive earth. *Nature* 190, 389-390.
- Oro, J. (1961b). Mechanism of synthesis of adenine from hydrogen cyanide under possible primitive Earth conditions. *Nature* 191, 1193-1194.
- Oro, J. (1994). Chemical synthesis of lipids and the origin of life. *J. Biol. Phys.* 20, 135-147.
- Oro, J., Miller, S.L., and Lazcano, A. (1990). The origin and early evolution of life on Earth. *Ann. Rev. Earth Planet Sci.* 18, 317-356.
- Ponnampereuma, C. (1995). The origin of the cell from Oparin to the present day. In: Ponnampereuma, C. and Chela-Flores, J. (Eds.). (1995). *Chemical Evolution: The Structure and Model of the First Cell* Kluwer Academic Publishers, Dordrecht, The Netherlands. pp. 3-9.
- Ponnampereuma, C., Lemmon, R.M., Mariner, R., and Calvin, M. (1963). Formation of adenine by electron irradiation of methane, ammonia, and water. *Proc. Natl. Acad. Sci. USA* 49, 737-740.
- Quack, M. (1993). Molecular quantum dynamics from high resolution spectroscopy and laser chemistry. *J. Mol. Struct.* 292, 171-195.
- Raulin, F., Bruston, P., Coll, P., Coscia, D., Gazeau, M-C., Guez, L. and de Vanssay, E. (1995). Exobiology on Titan. In: Ponnampereuma, C. and Chela-Flores, J. (Eds.). (1995).

Chemical Evolution: The Structure and Model of the First Cell Kluwer Academic Publishers, Dordrecht, The Netherlands. pp. 39-53.

Salam, A. (1990). Unification of the fundamental forces. Cambridge: Cambridge University Press. pp. 56-58.

Salam, A. (1991). The role of chirality in the origin of life. *J. Mol. Evol.*, 33, 105-113.

Salam, A. (1992). Chirality, phase transitions, and their induction in amino acids. *Phys. Lett. B288*, 153-160.

Salam, A. (1993). The origin of chirality, the role of phase transitions and their inductions in amino acids. In: *Chemical Evolution: Origin of Life*. Ponnampereuma, C. and Chela-Flores, J. (Eds.). A. Deepak Publishing: Hampton, Virginia, USA. pp. 101-117.

Salam, A. (1995). Biological macromolecules and the phase transitions they bring about. *Conceptual Tools for Understanding Nature*. G. Costa, G. Calucci, and M. Giorgi, (eds.). World Scientific Publishers: Singapore. pp. 209-220.

Saldanha, R., Mohr, G., Belfort, M., and Lambowitz, A.M. (1993). Group I and group II introns. *FASEB J.* 7, 15-24.

Schidlowski, M. (1995). Early terrestrial life: Problems of the oldest record. In: Chela-Flores, J., Chadha, M. Negron-Mendoza, A. and Oshima, T. (Eds.). *Chemical Evolution: Self-Organization of the Macromolecules of Life*. A. Deepak Publishing: Hampton, Virginia, USA. pp. 65-80.

Schopf, J.W. (1993). Microfossils of the Early Archean Apex Chert: New Evidence of the Antiquity of Life, *Science* 260, 640-646.

Schwartz, D.K. (1993). Instant patterns in thin films. *Nature* 362, 593-594.

Selinger, J.V., Wang, Z.-G., Bruinsma, R.F., and Knobler, C. M. (1993). Chiral symmetry breaking in Langmuir monolayers and smectic films. *Phys. Rev. Letters* 70, 1139-1142.

Sleep, N.H., Zahnle, K.J., Kasting, J.F., and Morowitz, H.J. (1989). Annihilation of ecosystems by large asteroid impacts on the early Earth. *Nature* 342, 139-142.

Soffen, G.A. (1976). Scientific results from the Viking Mission. *Science* 194, 1274-1276.

Yagubskii, E.B., Shchegdev, I.F., Laukhin, V.N., Konovich, P.A., Karatsovnik, M.V., Zvarykina, A.V., and Buravov, L.I. (1984). Normal pressure superconductivity in an organic metal (BEDT-TTF)₂I₃[bis(ethylene dithiolo) tetrathiofulvalene triiodide]. *JETP Lett.* 39, 12-16.

Walker, D.C. (1979). *Origins of optical activity in nature*. Amsterdam: Elsevier.

Wang, W. Q., X. Sheng, F. Jin, J. Wu, B. Yin, J. Li, Z. Zhao, H. Yang, Zhuang, Z., Lu, F., G. Yu, F., Shi, L. and Chen, Z. (1995). Susceptibility discontinuity and specific heat anomaly in

single crystal of alanine and valine. Paper presented at the Fourth Trieste Conference on Chemical Evolution: The Cyril Ponnampereuma Memorial Conference, Trieste, Italy, 4-8 September, 1995.

