

LAMP/92/1

**INTERNATIONAL CENTRE FOR
THEORETICAL PHYSICS**

**LAMP
SERIES REPORT**

(Laser, Atomic and Molecular Physics)

**OBSERVATION OF SQUEEZED LIGHT
AND QUANTUM DESCRIPTION
OF THE MACROSCOPICAL BODY MOVEMENT**

V.P. Bykov

MIRAMARE-TRIESTE



**INTERNATIONAL
ATOMIC ENERGY
AGENCY**



**UNITED NATIONS
EDUCATIONAL,
SCIENTIFIC
AND CULTURAL
ORGANIZATION**

International Atomic Energy Agency
and
United Nations Educational Scientific and Cultural Organization
INTERNATIONAL CENTRE FOR THEORETICAL PHYSICS

**LAMP
SERIES REPORT**
(Laser, Atomic and Molecular Physics)

**OBSERVATION OF SQUEEZED LIGHT AND QUANTUM DESCRIPTION
OF THE MACROSCOPICAL BODY MOVEMENT**

V.P. Bykov *

International Centre for Theoretical Physics, Trieste, Italy.

ABSTRACT

The possibility of a nondemolition measurement (observation) of macroscopical objects in widely distributed quantum mechanical states arises from the fact of the squeezed light observation. Macroscopical bodies – bodies of classical mechanics – are usually in states with narrow wave packets. It is shown that the absence of macroscopical bodies in widely distributed states is due to the focusing influence of the body's gravity field on its wave packet. An evidence that the gravity is essential in the classic limit of quantum mechanics is given.

MIRAMARE – TRIESTE

October 1992

* Permanent address: General Physics Institute of the Russian Academy of Sciences, Vavilova 38, 117942 Moscow, Russia.
Email address: BYKOV@OAK.FIAN.MSK.SU

In this paper we want to attract some attention to the consequences for some basic questions of quantum theory which follow from the observation of squeezed light. This observation is one of the most important achievements in optics in the last years and it will have many scientific continuations and practical applications. But we think that the most important consequence of the squeezed light observation will be some changes in our understanding of quantum mechanics. In essence, radical changes in the quantum measurement procedure appeared in the observation of squeezed light and these changes appeared imperceptibly.

1. QUANTUM DESCRIPTION OF THE MACROSCOPICAL BODY MOVEMENT.

We are concerned with only one problem - the quantum description of the macroscopical bodies movement. The macroscopical body is a body of significant mass, for example, 1 gramm (later we define more exactly this value). So we consider the possibility of a quantum description of body movement, which obeys, as is well known, the laws of classical mechanics. Let us consider a simple example - a uniform movement of macroscopical body with mass m and velocity \bar{v} in a free space. This movement is described by the wave packet [1]

$$\Psi(\bar{r}, t) = \frac{C}{(1+it/\tau)^{3/2}} \exp \left[-\frac{\bar{r}^2 - 2i\bar{v}\bar{r}t + i\bar{v}^2 \tau t}{2a^2(1+it/\tau)} \right] \quad (1.1)$$

with a characteristic size a , connected with the packet decay time τ by relation

$$\tau = ma^2/\hbar. \quad (1.2)$$

For a macroscopic body ($m \approx 1g$) and for a characteristic packet size of the order of the size of an atom ($a \approx 10^{-8}cm$), the packet (1.1) describes a well-defined rectilinear trajectory over great time interval, $\tau \approx 10^{11}sec$ ($\approx 3 \cdot 10^3 years$).

However, quantum mechanics does not prohibit large values of the parameter a , comparable, say, to the optical wavelength or even to the

geometric size of the macroscopic body at any value of its mass. Narrow wave packets, many times smaller than the geometric size of the body or even smaller than the typical optical wavelength (10^{-6} m), can be naturally attributed to macroscopical classical objects, since in this case the existence of the wave packet can be neglected generally considering it as a point.

However, to what can we attribute wide wave packets? No one macroscopical classical object with great quantummechanical uncertainty of its position was observed in the world surrounding us. If such objects could be seen we would not have such a science as a classical mechanics with its precisely definite trajectories. This fact, peculiar to a quantum mechanics, was observed long ago and in particular, gave initiative to a worldwide interpretation of the widespread solutions of the Schrödinger equation. These solutions were interpreted in an ensemble, statistical sense. It was acknowledged that in the measurement of the coordinate of some definite body, similar to that which was described by packet (1.1), one can obtain any value, but if one executes many measurements on bodies in identical states then the distribution of the probability of coordinate will be described by the squared module of the wave function $\Psi(\vec{r}, t)$. For example in the textbook of A. Messiah [2] in the discussion of the widespread solutions of the Schroedinger equation it is said: "In the classical approximation the function Ψ describes the "liquid" of classical noninteracting particles of mass m (statistical ensemble)...". A similar interpretation of the distributed solutions is given in the textbook of L. Schiff [1].

Just this ensemble, statistical interpretation of the widespread solutions can be disputed in connection with the lab observation of the squeezed light. We shall discuss it later and now show some additional examples of the distributed states of the macroscopic bodies.

2. MECHANICAL OSCILLATOR.

In quantum mechanics the hamiltonian [3]

$$H = \frac{1}{2m} P^2 + \frac{1}{2} \kappa Q^2 \quad (2.1)$$

describes a mechanical oscillator (fig.1) with operators of coordinate Q and conjugate momentum P subjected to the commutation relation

$$QP - PQ = i\hbar, \quad (2.2)$$

m is a mass and κ is an elasticity coefficient of the oscillator spring. Introducing variables $p = P/\sqrt{m}$, $q = Q/\sqrt{m}$ transform a hamiltonian to the more

usual form

$$H = \frac{1}{2}(p^2 + \omega^2 q^2), (\omega^2 = \kappa/m). \quad (2.3)$$

Operators of creation a^+ and annihilation a of the elementary oscillator excitations

$$a^+ = \frac{1}{\sqrt{2\hbar m \omega}} (\omega q - ip), \quad a = \frac{1}{\sqrt{2\hbar m \omega}} (\omega q + ip) \quad (2.4)$$

are important in the oscillator theory. In particular, coherent state of the oscillator $|z\rangle$ (fig.2a) is an eigenstate of the annihilation operator $a|z\rangle = z|z\rangle$; it is a solution of the Schroedinger equation at $z = z_0 e^{-i\omega t}$ and in a coordinate representation it is described by the wave function

$$\Psi_{\text{coh}}(q) = A \exp \left[-\frac{\omega}{2\hbar} \left(q - \sqrt{\frac{2\hbar}{m\omega}} z \right)^2 \right]. \quad (2.5)$$

The distribution for this state is shown in fig.2a; it is going periodically from right to left and back according to the harmonic law with amplitude $|z_0|$ frequency ω . Since the distribution width (or dispersion) of this state is essentially smaller than the oscillation amplitude at $|z| \gg 1$ ($Q \approx 1 \text{ cm}$, $\Delta Q/Q \approx 10^{-14}$) and going to zero with $\hbar \rightarrow 0$, this state naturally describes classical oscillations of the mechanical oscillator. Coherent state is a typical state with a narrow wave packet.

However there are also the stationary states $|n\rangle$, (fig.2b, n - whole numbers) which are the eigen states of the number particle operator (the number of the oscillator excitations): $a^+ a |n\rangle = n |n\rangle$. The state $|n(t)\rangle = e^{-in\omega t} |n\rangle$ is also a solution of the Schroedinger equation and in the coordinate representation is described by the wave function

$$\Psi_{\text{stat}}(q) = A H_n(\sqrt{\omega/\hbar} q) \exp(-\frac{1}{2}\omega q^2/\hbar), \quad (2.6)$$

where H_n is the Hermite polynomial of the n degree. The coordinate uncertainty in a coherent state is equal to $\Delta Q_{\text{coh}} = \sqrt{\hbar/2m\omega}$ and in a stationary state is equal to $\Delta Q_{\text{stat}} = \sqrt{(2n+1)\hbar/2m\omega}$.

Uncertainty in stationary state

$$\Delta Q \approx \omega^{-1} \sqrt{E/n}$$

is of the order of the oscillation amplitude in the coherent state at the equal energy ($n = |z|^2$), i.e. it is great at the great enough energy. So the stationary states at large n are typical widely distributed states. They are also macroscopical since they exist at high energies ($n \gg 1$) and great masses m . In classical mechanics there is no

appropriate movement of some specific object and the stationary states demand an ensemble, statistical interpretation which we are concerned with above.

3.A MACROSCOPIC BODY MOVEMENT IN THE GRAVITATIONAL FIELD.

The movement of the material point of the mass m in the gravitational field $U(\vec{r})$, which changes slowly in space, can be described by the gaussian wave packet [4]

$$\Psi(\vec{r}, t) = C(t) \exp \left[-(\vec{p}, F\vec{p}) + \frac{i}{\hbar} (\vec{p}_0(t)\vec{p} + E(t)) \right], \quad (3.1)$$

where $\vec{p} = \vec{r} - \vec{r}_0(t)$ and $\vec{r}_0(t), \vec{p}_0(t)$ are solutions of the classic hamiltonian equations

$$\dot{\vec{r}}_0 = \vec{p}_0/m, \quad \dot{\vec{p}}_0 = -\text{grad}_{\vec{r}_0} U(\vec{r}_0); \quad (3.2)$$

3×3 matrix $F(t)$ is symmetric and has complex elements, which are functions of time (real part of $F(t)$ defines geometric dimensions of the wave packet).

This wave packet is a solution of the Schroedinger equation if in the potential energy expansion near the \vec{r}_0 point

$$U(\vec{r}_0 + \vec{p}) = U(\vec{r}_0) + (\vec{p}, \text{grad}_{\vec{r}_0} U(\vec{r}_0)) + \frac{1}{2} (\vec{p}, U''\vec{p}) + \dots \quad (3.3)$$

the third order terms marked by dots can be neglected. The 3×3 matrix U'' of the second derivatives is also symmetric. Its real elements

$$U''_{\alpha\beta}(\vec{r}) = -\frac{\partial^2 U(\vec{r})}{\partial x_\alpha \partial x_\beta} \Big|_{\vec{r}=\vec{r}_0(t)} \quad (3.4)$$

are time-dependent through the vector $\vec{r}_0(t)$.

The wave packet (3.1) is moving along a classical trajectory (3.2) until its dimensions are so small that the third order terms in the potential energy (3.3) can be neglected. The matrix $F(t)$, which defines, in particular, the wave packet dimensions, must satisfy the matrix equation

$$i\hbar \dot{F} = \frac{2\hbar^2}{m} F^2 - \frac{1}{2} U'', \quad (3.5)$$

which is similar to the Ricatti equation. The values $C(t)$ and $E(t)$ are equal to

$$C(t) = C_0 \exp \left[-\frac{i\hbar}{m} \int_0^t dt (F_{11} + F_{22} + F_{33}) \right], \quad E(t) = \int_0^t dt \left[\frac{\vec{p}_0^2}{2m} - U(\vec{r}_0) \right]. \quad (3.6)$$

For macroscopical bodies the gravitational field is the most important. As is well-known the classic mechanics appeared from the observations on the macroscopical bodies - planets movement in the field of Sun. In this case the trajectory (3.2) does not depend on mass, since \vec{p}_0 and $U(\vec{r}_0)$ are both proportional to m . Introducing in (3.5) an evident

dependence $U(\vec{r}_0)$ on mass m , the equation

$$i\dot{F} = 2F^2 - \frac{1}{2} V, \quad (F = F\hbar/m, mV = U''), \quad (3.7)$$

in which all quantities do not depend on mass m , can be obtained. So one can see, that F is growing with the growth of mass m and it means that the wave packet dimensions are decreasing and the wave packet concentrates itself more and more on the classic trajectory (3.2). Consequently the wave packet (3.1) describes naturally, in the frame of quantum mechanics, the macroscopic body movement.

But this is not the only way to give a quantum description of the macroscopic body motion. Another way is connected with a well-known eikonal method [2]. When the mass m is large enough and consequently when the de Broglie wavelength $\lambda = \hbar/\sqrt{2mE}$ is small (E -energy of state) the stationary solution of the Schroedinger equation can be written in the form

$$\Psi(\vec{r}, t) = A(\vec{r}) \exp[i(S(\vec{r})/\hbar - Et/\hbar)], \quad (3.8)$$

where $S(\vec{r})$ - the eikonal function, which changes a little on the length of the order of the wavelength λ and which satisfies the equation

$$(\text{grad} S(\vec{r}))^2 = 1 - (U/E). \quad (3.9)$$

Amplitude $A(\vec{r})$ also changes a little on the length of the order of the wavelength λ and satisfies the equation

$$(\text{grad} S(\vec{r}) \text{ grad} A(\vec{r})) + \frac{1}{2} A \Delta S = 0. \quad (3.10)$$

In a gravitational field, U and E , both proportional to the mass m and then S_i do not depend on m . Curves orthogonal to the wave surfaces $S(\vec{r}) = \text{const}$ are the classical trajectories of the material body.

In this section basic elements of quantum description of the macroscopical body movement in a gravitational field [2,4] are given. As in the mechanical oscillator, there are two types of quasiclassical solutions. The narrow gaussian wave packet solution (3.1) naturally describes a classical movement of the macroscopical body. But there are also distributed solutions of the eikonal type. The eikonal function $S(\vec{r})$ is wide as in a longitudinal, so in the transversal directions to the trajectories. In the transversal direction it can be bound only by possible caustic surfaces (see, for instance, fig.3). If such/state is considered as a state of one definite body then there is nothing in classic mechanics corresponding to it.

To conclude this section it should be noted that in quantum mechanics the transition to a classic limit is not so automatic as, e.g. in the theory of relativity. In the case of relativity theory, it is sufficient for the body velocity divided by the light velocity to be small in order to obtain

classical mechanics laws from the relativity laws. But in quantum mechanics only some of the possible states, namely narrow wave packets, reduce to the classical motion with the growth of the mass. Another part of possible states - distributed solutions - can be interpreted in a classical style only in the ensemble, statistical sense. Narrow wave packets can also be interpreted in the ensemble, statistical sense of course, but for them this interpretation is not necessary.

4. SQUEEZED LIGHT AND ITS OBSERVATION.

Let us now look at a squeezed light. As a laser light it can be excited in an optical cavity. The cavity field is a superposition of the fields of the partial modes or resonances. It is well-known from practice that in a cavity one mode field excitation is possible. Since the space distribution of the mode field is defined by the boundary conditions, so the mode field can be considered as a system with one degree of freedom which coordinate is an electric field of the mode in some chosen point of the cavity. Then the quantum theory of the mode field coincides with the quantum theory of the mechanical oscillator. This field has the same hamiltonian (2.3), where coordinate q must be replaced by the mode electric field E in the chosen point of the cavity. In particular, different states - coherent and stationary, which were mentioned above - are possible for the mode field.

But these two state types are not only possible in mechanical or electromagnetic oscillators. Squeezed state gather in itself the properties of the narrow and widespread states [5,6]. In this state the electric field value distribution is described by the gaussian wave packet (fig.2a) as in the coherent state but the dispersion of this distribution is different from the dispersion of the coherent (or vacuum) state. It can be shown that in this case the dispersion changes periodically with the double frequency of the oscillator from the value smaller than the dispersion of the coherent state to the value greater than it.

Basic elements of the squeezed state theory are given below. Squeezed states of light [5,6] are the eigen states $|\zeta\rangle$ of the operator $\mu a + \nu a^\dagger$

$$(\mu a + \nu a^\dagger)|\zeta\rangle = \zeta|\zeta\rangle, \quad (|\mu|^2 - |\nu|^2 = 1), \quad (4.1)$$

where μ, ν, ζ - complex numbers. If μ and ν depend on time $\mu = \mu_0 e^{i\omega t}$, $\nu = \nu_0 e^{-i\omega t}$, then the states $|\zeta\rangle$ are the solutions of the Schroedinger equation. The most expressive representative of the squeezed states is the squeezed vacuum state corresponding to $\zeta=0$. In coordinate

representation it is described by the wave function

$$\Psi_\zeta(q) = A \exp\left[-\frac{\omega q^2}{2\hbar} \frac{\mu + \nu}{\mu - \nu}\right]. \quad (4.2)$$

The squeezed state dispersion

$$D^2 = \frac{\hbar}{2\omega} \left[|\mu|^2 + |\nu|^2 - 2|\mu||\nu| \cos(\Psi_0 + 2\omega t) \right] \quad (4.3)$$

oscillates with the double oscillator frequency. It can be smaller than the coherent state dispersion $D_{\min}^2 = \frac{\hbar}{2\omega} (|\mu| - |\nu|)^2 < D_{\text{coh}}^2 = \frac{\hbar}{2\omega}$ and can be greater than it $D_{\max}^2 = \frac{\hbar}{2\omega} (|\mu| + |\nu|)^2 > D_{\text{coh}}^2 = \frac{\hbar}{2\omega}$ (fig.4). The instants when the dispersion is small gave initiative to the term - squeezed state. A squeezing coefficient

$$K = D_{\text{coh}}/D_{\min} = \sqrt{D_{\max}/D_{\min}} = |\mu| + |\nu| = (|\mu| - |\nu|)^{-1} \quad (4.4)$$

is a characteristic of the squeezed state. There is an energetic limitation for the squeezing coefficient. At the average photon number N the maximum squeezing coefficient is equal to

$$K_{\max} = \sqrt{N+1} + \sqrt{N}. \quad (4.5)$$

The qualitative picture of the squeezed state can be accepted from fig.4. Fig.4a gives the coordinate oscillations versus time in the coherent state. The constant dispersion of this state is shown by the thickness of sine line. At the usual laser intensity, say, at the 1 joule energy, stored in the laser cavity, the ratio of the dispersion to the oscillation amplitude is small, $\approx 10^{-9} - 10^{-10}$. Fig.4b shows oscillations in the squeezed state. Here the dispersion changes with time and at some instants is of the same order of value as amplitude. The points, where the dispersion is minimal, suggested the name of squeezed states. Fig.4c shows "oscillations" in squeezed vacuum state. One should be fully aware that now there are no oscillations with ω frequency, but only the dispersion changes with the double frequency. We emphasise that although the oscillations are absent the squeezed vacuum is a highly excited, macroscopical state with big energy.

Parametric excitation of the oscillator is described by the Schroedinger equation

$$i\hbar \frac{\partial \Psi(q,t)}{\partial t} = \frac{1}{2} [p^2 + \Omega^2(t)q^2] \Psi(q,t).$$

As can be seen a parameter of the oscillator - its resonant frequency Ω - depends on time. The solution of this equation, similar to the squeezed vacuum state, is

$$\Psi(q, t) = A(t) \exp \left[\frac{i \dot{\epsilon}(t) q^2}{2 \hbar \epsilon(t)} \right], \quad (4.6)$$

where $\epsilon(t)$ is the solution of the equation

$$\ddot{\epsilon}(t) + \Omega^2(t) \epsilon(t) = 0 \quad (4.7)$$

which satisfies initial conditions $\epsilon(0)=1$, $\dot{\epsilon}(0) = i\Omega_0$ (Ω_0 is the average oscillator frequency). The dispersion of the state (4.6) equals $D = |\dot{\epsilon}(t)| \sqrt{\hbar/2\Omega_0}$. The squeezing coefficient grows with time if $\epsilon(t)$ is not bound solution of the equation (4.7).

Squeezed light was already observed in some laboratories [7,8]

One of the possible devices was realized by a group of investigators from the Texas University and is given in fig.5. Although this experiment was not the first one, it is the most fruitful one from the engineering point of view. It consists of three major parts - the source of parametric pumping, the parametric generator and the detector-analyzer of squeezed states. The first two parts are similar to those which were used in the first experiments on the parametric generation [9,10]. The source of the parametric pumping power is the solid-state ring laser with a YAG:Nd³⁺ and with an intracavity excitation of the second harmonic in the nonlinear crystal. The second harmonic of the Nd-laser is used for pumping the parametric generator, where the first harmonic appears again but now in the squeezed vacuum state. The receiver-analyzer compares the signals from laser and parametric generator and measures the dispersion of the squeezed light in the instants close to the maximums of the amplitude of the laser signal. The results of observation are shown in fig.6. As one can see the parametric signal dispersion changes twice for a period of the laser signal: its period is π , but not 2π . One can see also that at some phases θ the squeezed light dispersion becomes smaller than the dispersion of the coherent light (dashed line). It is just evidence that the light is in the squeezed state.

Take notice the unusual character of the measurement procedure. In it there are no partial measurements of the field strength, which in usual quantummechanical measurement procedure provides the distribution $|\Psi(E)|^2$. The result of a measurement now is just the state parameter, in our case - the state dispersion. Another peculiarity of measurement is its nondemolition (or nondisturbing) character. Indeed the object of measurement is the light beam, coming from the parametric generator cavity. This beam does not come back into the cavity independently whether the measurement was made with it or not. If such measurement was made

the information about it cannot reach the cavity and therefore the field state in the cavity cannot be changed. But the results of measurement of the light beam, coming from the cavity, give us information about the state of the field in the cavity.

Note that due to the dynamical equilibrium between pumping and losses the parametric generator field is stationary. Therefore if we made a measurement, we can be sure that the field in the cavity remains

in the squeezed vacuum state for a long time. So the squeezed light observation tells us that there is a macroscopical (number of photons is big) quantummechanical object - the electromagnetic field of the picked out cavity mode - which dispersion we can measure, leaving the object itself in the state in which it was measured before.

The time instances when the dispersion of the field is smaller than the dispersion of the coherent state extremely attracted attention of the investigators. It is natural since the quantummechanical uncertainties, just as the noise, prevent the precise measurement of corresponding quantities. The smaller this uncertainty the more precisely corresponding quantity can be measured and small uncertainty is an attractive feature of the light squeezed states. But from the point of view of the researcher the states with a great uncertainty are more interesting; they are less classical and more quantummechanical. For this paper they are important since its observation, taking into account what was said above about the squeezed light measurements, shows that the measurement of the distributed state dispersion is possible in one object without disturbing the distributed state.

Naturally if such measurements are possible with electromagnetic oscillator, they are possible also with a mechanical oscillator. This idea we develop in the next section.

5. MACROSCOPIC BODIES IN DISTRIBUTED STATES AND THEIR NONDEMOLITION MEASUREMENT.

So it was shown something more in experiments with squeezed light than its generation possibility. In addition it is shown that the quantummechanical state of one macroscopical object (field of a particular cavity mode) can be learned in a nondisturbing manner. In particular, it can be learned that this object is in the widespread state when the uncertainty of its coordinate is of a macroscopical value approximately equal to the oscillation amplitude in the coherent state at equal energies of both states.

If such learning is possible with electromagnetic oscillator it is

also possible with a mechanical oscillator and even with a macroscopic body freely moving in the space since it can be seen as an oscillator with a zero frequency. Such conclusion is a natural consequence of a similar description of the electromagnetic and mechanical oscillators (see also below).

One can ask - why there are no macroscopic bodies in widespread states in the world surrounding us. Quantum mechanics does not prohibit such states (see first section), and in some cases they are preferable.

One possible explanation of it is concluded in supposition that at the observation of such bodies its wave packet reduction occurs. To exclude such possibility we considered the electromagnetic signal reflection from the body (mirror) which is in the widely distributed state [11]. Major results of this consideration are given below.

We suppose [11] that the mirror has two degrees of freedom (fig.7). The first degree of freedom, transversal oscillator describes the motion of the negative charges (bounded electrons) along the mirror (displacement Q , conjugate momentum P , surface mass density ρ , surface charge density σ). This motion leads to the reflection of the electromagnetic signal. The second degree of freedom, longitudinal oscillator describes the motion of the mirror along the direction of the electromagnetic wave propagation (coordinate q , conjugate momentum p , surface mass density μ). It is supposed also that there are only waves of normal incidence to the mirror. The mirror is infinitely thin and perpendicular to the z -axis along which waves propagate. The electrical field and the charge displacement directed along x -axis. The region of space occupied by the field has the same section form as a form of the mirror and its area is equal to s . Then quantum system mirror+field is described by hamiltonian

$$H = \frac{1}{2\rho s} \left[P - \frac{\sigma s}{c} A(q) \right]^2 + \frac{1}{2} s K Q^2 + \frac{1}{2} s \int dz \left[\frac{1}{4\pi} \left(\frac{\partial A}{\partial z} \right)^2 + 4\pi c^2 \Pi^2 \right] + \frac{1}{2\mu s} p^2 + \frac{1}{2} s \kappa q^2 \quad (5.1)$$

where K and κ - the transversal and longitudinal oscillator elasticities, $A(z)$ - vector-potential of the electromagnetic field and $\Pi(z)$ - its canonically conjugate momentum.

The field oscillators (plane waves) in the initial instant are in coherent states with such phases that the field forms a rectangular pulse filled with a high frequency oscillations

$$\langle E_{in}(z, t) \rangle = E_0 \sin \omega_0 \bar{t} \Pi(\tau_0, \bar{t}), \quad (5.2)$$

where $\Pi(\tau_0, \bar{t})$ - a function describing the rectangular form of the

in-pulse of the $2\tau_0 = 4\pi n_0 / \omega_0$ duration ($2n_0$ - number of wavelength in the pulse), $t = \bar{t} - z/c$ and \bar{t} - the instant of the arriving of the pulse at the origin. The transversal oscillator initially is in the ground or vacuum state and the longitudinal oscillator is in a squeezed state described by the wavefunction

$$\Psi(q) = (2\pi \bar{q}^2)^{-1/4} \exp(-q^2/4\bar{q}^2), \quad (5.3)$$

where $\bar{q}^2 = \hbar / (2s\mu\nu\sqrt{K})$ and K is the squeezing coefficient. The additional parameter in the wavefunction (5.3) - the coefficient of squeezing - provides the possibility to change the longitudinal oscillator parameters μ, ν in future without changing its distribution; in particular the transition to the motion in the free space ($\nu \rightarrow 0$) will become possible.

The investigation of this problem shows that the field averaged value in the reflected signal in the stationary conditions and in the resonance ($\Omega = \omega_0$) is

$$\langle E_R(z, t) \rangle = \frac{1}{2} E_0 e^{-2q^2/\bar{q}^2} \sin \Omega(\tau - \bar{t}), \quad (5.4)$$

where $\bar{q} = \lambda/2\pi = c/\Omega$ and $\tau = t + z/c$. The average value of the squared field in the reflected signal is

$$\langle E_R^2(z, t) \rangle = \frac{1}{8} E_0^2 \left[1 - e^{-8q^2/\bar{q}^2} \cos 2\Omega(\tau - \bar{t}) \right]; \quad (5.5)$$

this value is proportional to the electrical energy density. As can be seen when the position uncertainty of the mirror is small ($\bar{q} \ll \lambda$) the reflected signal keeps the properties of the coherent state; in particular $\langle E_R \rangle^2 \approx \langle E_R^2 \rangle$. When the position uncertainty is large ($\bar{q} \gg \lambda$) the field average value is close to zero but the average value of the squared field is not close to zero and keeps the finite value; only double frequency oscillations of energy density are close to zero. One can say that the amplitude reflection coefficient goes to zero when the position uncertainty grows, but the intensity reflection coefficient keeps finite value in the same case. It means that the reflected signal being a macroscopical one is in essentially quantum state, as only in such states the relation $\langle E \rangle^2 \ll \langle E^2 \rangle$ is possible.

It can be shown also that the length of the reflected signal is greater than the length of the incident signal approximately on \bar{q} .

Consequence of these results is the conclusion that it is possible to find out experimentally the macroscopical body (mirror) to be in a distributed quantummechanical state as a result of one pulse reflection

from this body without essential change of its state. Indeed the reflected pulse, as we could see, bears information about the distributed state of the mirror and being the macroscopical one can be analysed with the existing experimental means such as used for example to analyse the squeezed light [7,8]. At the same time it can be shown [13] that the reflection process does not change the mirror state essentially. Such possibility contradicts the usual interpretation of the distributed state as describing an ensemble of objects [1,2].

We call attention also to the lengthening of the reflected pulse $\langle E_R^2 \rangle$ compared with the incident one, due to the partial reflection of the incident pulse from different layer of the mirror distribution over the longitudinal coordinate. The human eye with its low temporal resolving power cannot, of course, notice such a lengthening (of the order \bar{q}) of the pulse. For oblique incidence of light from a point source on a mirror, however, this partial reflection from different distribution layers is converted into an angular distribution of different rays, and the angular resolution of the human eye is high enough. A naked eye, therefore, will see bodies with large quantum-mechanical uncertainty simply as blurred. Consequently, were macroscopic bodies with large quantum-mechanical uncertainty to exist in the world surrounding us, they could be simply seen with the naked eye.

6. WHY THERE ARE NO MACROSCOPIC BODIES IN DISTRIBUTED STATES.

The absence of macroscopic bodies in widespread states could be explained by their formation in the concentrated state at some initial time and their subsequent gradual diffusion. But there are no reasons why they should be formed just in the concentrated state at the initial time.

The concentration of the macroscopic body wavepackets cannot be explained in terms of their interaction with some fields. In fact, since bodies consist of charged particles, they must interact with the electromagnetic field. However, such interaction leads only to the formation of an associated (nonradiative) field, since all particles in uniform rectilinear motion have such fields. A similar argument also applies to other fields. In all cases, the packet (1.1) describes the motion of the center of inertia of the macroscopic body and its associated fields, and consequently, the problem of large values of the parameter α cannot be removed in this way.

An exception is the interaction of the macroscopical body with the gravitational field [12]. According to the general theory of relativity,

the gravitational field created by the body can be considered as a deformation of space, a deviation it from Euclidian, with this deformation occurring both in the region occupied by the body and in its vicinity. If the mass related to some part of the wave packet deforms space, the remaining parts of the packet move in this deformed space. In general, any part of the wave packet moves in the space deformed by this part itself and by all the other parts. Consequently, there is an effect of the wave packet on itself, or a self-effect of the wave packet caused by its interaction with the gravitational field. As is shown below this self-effect leads to the formation of an effective potential well within which the wave packet preserves permanently its concentrated form. Moreover, the formation of such a potential well is obviously energetically favored, and this explains the concentration of the wave packets of macroscopic bodies.

Since we are dealing with macroscopic bodies of ordinary density (of the order of 1 g/cm^3), the gravitational potential is weak, and therefore the Newtonian expression for the potential is sufficient for calculation.

Let us examine a uniform spherical body of radius R and density ν . Since we will calculate the gravitational effect of one part of the wave packet on another, and since the wave packet is much smaller than the geometric size of the body, we will first examine the attraction of two mutually penetrating massive spheres whose centers are displaced by a distance s which is much smaller than R (fig.8). As can be seen, the attraction between the spheres can be calculated by taking into account the attraction by the first sphere of a layer of thickness $2s$ which cover half of the first sphere. Considering only the projection of the forces of attraction of unit masses along the line joining centers of the spheres, we obtain the following expression for the force of attraction of the spheres:

$$F = \frac{16\pi^2}{3} G \nu^2 R^3 \int_0^{\pi/2} \sin\theta \cos\theta d\theta = \frac{8\pi^2}{3} G \nu^2 R^3 s. \quad (6.1)$$

Consequently, the potential energy of the interaction of the two spheres is

$$U_s = \beta s^2, \quad \beta = \frac{4}{3} \pi^2 G \nu^2 R^3. \quad (6.2)$$

Taking into account this expression for the potential energy, we obtain a Schrodinger equation describing the wave packets of the macroscopic bodies

$$i\hbar \frac{\partial \Psi(\vec{r}, t)}{\partial t} = -\frac{\hbar^2}{2m} \Delta \Psi(\vec{r}, t) + U(\vec{r}) \Psi(\vec{r}, t), \quad (6.3)$$

where

$$U(\vec{r}) = \beta \int d\vec{r}' (\vec{r} - \vec{r}')^2 |\Psi(\vec{r}', t)|^2. \quad (6.4)$$

This equation shows that the gravitational self-effect of the wave packet of a macroscopic body is similar to the optical self-focusing effect. The main difference is that optical self-focusing occurs in the two directions normal to the direction of wave propagation, whereas the gravitational self-effect occurs in all three directions.

Let us find the steady-state, spherically symmetric solution of Eq.(6.3). For $\Psi = \Psi(\vec{r})$ and $\text{Im}\Psi = 0$, we have, according to (6.4),

$$U(\vec{r}) = \alpha + \beta r^2, \quad (6.5)$$

where $\alpha = \beta \int dr' r'^2 |\Psi(r')|^2$ is an unimportant constant. Thus, to determine $\Psi(r)$, we have the equation

$$-\frac{\hbar^2}{2m} \frac{1}{r^2} \frac{d}{dr} \left(r^2 \frac{d\Psi}{dr} \right) + \beta r^2 \Psi = E\Psi, \quad (6.6)$$

whose solution is

$$\Psi(r) = \frac{1}{\sqrt{r}} \exp\left(-\frac{r}{2r_0^2}\right), \quad (6.7)$$

where

$$r_0 = \sqrt{\frac{2\hbar^2 R^3}{3Gm^3}} \quad (6.8)$$

is the characteristic size of the wave packet due to the gravitational self-effect. This concentrated wave packet can be considered as a soliton solution of the Schroedinger equation (6.3). Some values are presented in the following table:

Table

m, g	R, cm	r_0 , cm
10^{28}	$1.3 \cdot 10^9$	$1.25 \cdot 10^{-20}$
10^{12}	$6.2 \cdot 10^3$	$1.25 \cdot 10^{-18}$
1	0.62	$1.25 \cdot 10^{-12}$
10^{-12}	$6.2 \cdot 10^{-3}$	$1.25 \cdot 10^{-6}$
10^{-15}	$6.2 \cdot 10^{-6}$	$4 \cdot 10^{-5}$

The mass, geometric size of the macroscopic body with density $\nu \approx 1 \text{g/cm}^3$ and size of its wave packet are given in the respective columns. Data for a typical macroscopic body ($m=1\text{g}$) is given in the middle row of Table. The top row presents data for a body with a mass of the order of the mass of the Earth, while the last row is that for a

mass of 10^{-15}g , for which the geometric size is comparable to the size of the wave packet. The size corresponding to the condition $r_{cr} = r_0 = R$ can be determined from (6.8):

$$r_{cr} = \sqrt[10]{\frac{9\hbar^2}{32G\nu^3}}. \quad (6.9)$$

The condition (6.9) gives a quantitative criterion for dividing bodies according to their size into macroscopic and microscopic. A criterion for such division by mass can also be obtained from (6.9). Bodies with mass greater than

$$m_{cr} = \sqrt[10]{\frac{32\pi\nu\hbar^6}{81G^3}} \quad (6.10)$$

should be considered macroscopic.

Thus, for masses greater than 10^{-12}g , the size of the wave packet is negligible not only compared to the geometric size of these masses, but also compared to the typical optical wavelength ($10^{-4} - 10^{-5} \text{cm}$).

This discussion explains why there are no macroscopic bodies in states with large quantum-mechanical uncertainty of their center of mass. But this explanation was reached by a very high prize, namely by giving up the superposition principle and the linearity of the Schroedinger equation [13], as can be seen from (6.3). However the violation of the superposition principle is not very profound. In particular, it will not influence the microscopical, atomic effects - the basic field of the quantum mechanics application.

It may be that the giving up the superposition principle is not definite. After the creation of the quantum theory of gravity our considerations can appear as similar to a semiclassical theory in quantum electrodynamics in which the medium is described quantummechanically and the radiation - classically. In quantummechanical description of both the gravity and medium superposition principle can be restored.

The potential well discussed above is not very deep. By expanding some energy a macroscopic body can therefore be changed to a widespread state with a large wave packet. Study of the distributed states of macroscopic bodies would be of great scientific value.

7. CONCLUSION

Considerations stated above show that in mechanics of macroscopical bodies gravity plays an important role. Due to gravity the wave packets of macroscopical bodies are concentrated and the classical

mechanics is mechanics of concentrated but not widespread packets.

These considerations also show that macroscopical bodies can be in the widespread states and it is a matter of importance to observe them practically. Bodies in such states will be new objects of physical investigation. A natural way to excite such states is a parametric excitation of the oscillator. An oscillator excited in this way twice in a period becomes widely quantummechanical distributed.

It is clear also why the widespread quantum states were observed first for the electromagnetic field but not for mechanical system. The electromagnetic field mass is very small at usual intensities and it cannot deform the space, So the concentrated states of field do not have an advantage over the widespread states.

In connection with what has been said above it is correct to refer to the A. Einstein paper [4], who was the first to raise a question about the correct description of the macroscopic body movement.

ACKNOWLEDGMENTS

The author would like to thank A.M. Prokhorov and F.V. Bunkin for support and also F. Persico and L. Pitaevskii for critical remarks. Thanks are also due to Professor Abdus Salam, the International Atomic Energy Agency and UNFSCO for hospitality at the International Centre for Theoretical Physics, Trieste.

REFERENCES

1. L.I. Schiff, 1955, Quantum mechanics, N.Y., McGraw-Hill Book Company, Inc., N.Y.
2. A. Messiah, Mécanique quantique, 1959, v.1,2, Paris, Dunod
3. W.H. Louisell, 1964, Radiation and noise in quantum electronics, N.Y., McGraw-Hill Book Comp., Inc., N.Y.
4. V.P. Bykov, Lebedev Inst. Reports 1989, No.1, p.3.
5. K. Husimi, Progr. Theor. Phys., 1953, v.9, p.381
6. D. Stoler, Phys. Rev., 1970, v.D1, p.3217; 1971, v.D4, p.1925
7. R.E. Slusher, L.W. Hollberg, V. Yurke, C. Mertz, J.F. Valley, Phys. Rev. Lett., 1985, v.55, p.2409
8. Ling-An Wu, H.J. Kimble, J.L. Hall, Huifa Wu
Phys. Rev. Lett., 1986, v.57, p.2520
9. S.A. Akhmanov, R. Khokhlov, Sov. Phys. JETP, 1962,
10. J.A. Giordmaine, R.C. Miller, Phys. Rev. Lett., 1965, v.14, p.973
11. V.P. Bykov, Bull. Acad. Sci. USSR, Phys. Ser., v.55, N2, p.15, (1991)
OP AC Review, v.1, N2, p.173, (1990-1991)
12. V.P. Bykov, 1989, Sov. Phys. Dokl., 1989, v.34(10), p.911
13. V.P. Bykov, 1991, Journ. Sov. Laser Res., v.12, N1, p.35
14. A. Einstein, Scientific Papers, presented to Max Born, 1953,
Edinburgh, Oliver Boyd, p.33

FIGURE CAPTIONS

1. A mechanical oscillator.
2. Quantummechanical states of oscillator:
 - a. Coherent state
 - b. stationary states
3. An orbit system for the gravitational field which forms an eikonal solution of the Schroedinger equation:
 - a. trajectory, b. wave front c. caustic lines
4. Field oscillations versus time:
 - a. coherent state b. squeezed state c. squeezed vacuum state
5. A lab device for observation of the squeezed light:

RL - ring Nd-laser with intracavity generation of the second harmonics, OPO - optical parametric oscillator, LO - local oscillator, PD - photodiode, FH - first harmonics, SH - second harmonics
6. A dispersion of the squeezed light versus the phase of the coherent signal
7. A reflection of light from the mirror, which quantummechanical state is widely distributed, IP initial pulse.
8. Attraction of two mutually penetrating balls.




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8