

United Nations Educational, Scientific and Cultural Organization
and
International Atomic Energy Agency

THE ABDUS SALAM INTERNATIONAL CENTRE FOR THEORETICAL PHYSICS

URANIUM GROUNDWATER ANOMALIES AND ACTIVE NORMAL FAULTING

Wolfgang Plastino^{a,b}, Giuliano Francesco Panza^{c,j}, Carlo Doglioni^d,
Maria Luce Frezzotti^e, Angelo Peccerillo^f, Pierino De Felice^g, Francesco Bella^a,
Pavel P. Povinec^h, Stefano Nisiⁱ, Luca Ioannucciⁱ, Piergiorgio Apriliⁱ, Marco Balataⁱ,
Maria Letizia Cozzella^g and Matthias Laubensteinⁱ

^a*Department of Physics, University of Roma Tre, Rome, Italy*

^b*National Institute of Nuclear Physics, Section of Roma Tre, Rome, Italy*

^c*Department of Earth Sciences, University of Trieste, Trieste, Italy*

^d*Department of Earth Sciences, University of Roma "La Sapienza", Rome, Italy*

^e*Department of Earth Sciences, University of Siena, Siena, Italy*

^f*Department of Earth Sciences, University of Perugia, Perugia, Italy*

^g*ENEA, Department of Environment, National Institute for Metrology of Ionizing Radiations,
C.R. Casaccia, Rome, Italy*

^h*Department of Nuclear Physics and Biophysics, Comenius University, Bratislava, Slovakia*

ⁱ*National Institute of Nuclear Physics, Gran Sasso National Laboratory, Assergi (AQ), Italy*

^j*Abdus Salam International Centre for Theoretical Physics, Trieste, Italy*

MIRAMARE – TRIESTE

August 2010

Abstract

The ability to predict earthquakes is one of the greatest challenges for Earth Sciences. Radon has been suggested as one possible precursor, and its groundwater anomalies associated with earthquakes and water-rock interactions were proposed in several seismogenic areas worldwide as due to possible transport of radon through microfractures, or due to crustal gas fluxes along active faults. However, the use of radon as a possible earthquake's precursor is not clearly linked to crustal deformation. Here we show that uranium groundwater anomalies, which were observed in cataclastic rocks crossing the underground Gran Sasso National Laboratory, can be used as a possible precursor of earthquakes in domains where continental lithosphere is subducted. Measurements evidence clear, sharp anomalies from July, 2008 to the end of March, 2009, related to a preparation phase of the seismic swarm, which occurred near L'Aquila, Italy, from October, 2008 to April, 2009. On April 6th, 2009 an earthquake ($M_w=6.3$) occurred at 01:33 UT in the same area, with normal faulting on a NW-SE oriented structure about 15 km long, dipping toward SW. In the framework of the geophysical and geochemical models of the area, these measurements indicate that uranium may be used as a possible strain meter in extensional tectonic settings similar to those where the L'Aquila earthquake occurred.

Introduction

Earthquake prediction, due to its societal relevance and the intrinsic complexity of the problem, has been the subject of several controversial discussions and reviews (1-4).

The geodynamic processes leading to earthquakes can modify radon (Rn) migration patterns in groundwater that could be used as a potential strain meter probe (5). Radon (Rn) as a possible candidate for earthquake's precursor has been studied for a long time, but there is no clear evidence that it is really a precursor (6). It has been suggested as one of several possible early signals, and its air or groundwater anomalies associated with earthquakes and air-rock or water-rock interactions were detected in several seismogenic areas worldwide (7-18). The physical processes associated with Rn anomalies are based on the changes of Rn emanation rates occurring due to strain signal near to the earthquake's nucleation point. Particularly, its behaviour before, during and after the main shock, considering the consolidated scheme for Rn release due to stress-strain processes in the rock is unclear. In the geological environment, the Rn concentration depends on the isotopic abundance of its parent radionuclides (^{238}U and ^{226}Ra), and on their geochemical patterns with reference to environmental redox and pH characteristics. The geodynamic processes induced by earthquakes can modify Rn migration patterns as a potential indicator of strain. However, the activity of Rn in fractured lithologies is difficult to predict, and the Rn concentration does not uniquely constrain the rock deformation or the chemical inhomogeneity nor its relationship with the transient crustal strain signals from 'aseismic' fault slip, near to the earthquake's nucleation point (6). Moreover, non-tectonic factors related to variations of chemical and physical groundwater parameters may be of importance and proper geological, hydrological and hydrogeological settings (16) are required to isolate the variations induced by stress-strain processes.

Results and Discussion

The Gran Sasso National Laboratory of the National Institute of Nuclear Physics (LNGS-INFN) is located in the Gran Sasso massif (central Apennines, Italy) inside the largest aquifer of central Italy, within Mesozoic carbonates. These rocks have been deformed and fractured during an earlier Miocene-Pliocene compressive phase. Since Pleistocene, the massif has been cross-cut by normal faults (17).

Within the framework of the INFN's scientific program ERMES (Environmental Radioactivity Monitoring for Earth Sciences) Rn (17), radiocarbon (^{14}C) (19) and tritium (^3H) (20) have been monitored in the groundwater inside the LNGS-INFN, and different chemical, physical and fluid dynamical characteristics of groundwater have been measured. The uranium (U) groundwater monitoring started on June 2008 with the aim of better defining the Rn groundwater transport processes through the cataclastic rocks (17), as well as to check its contribution to the neutron background at the LNGS-INFN (21). The U measurements (Fig. 1) carried out between June 2008 and May 2010 at four different sites (E1, E4 and E3, E3dx)

show the presence of two different water groups, whose existence is confirmed by stable isotope of hydrogen and oxygen ($\delta^2\text{H}$, $\delta^{18}\text{O}$), ^{14}C and ^3H analyses (22).

A seismic swarm, whose main shock occurred at 01:33 UT on April 6th, 2009, has affected the area under investigation from October 2008 to April 2009. The main ($M_w=6.3$) event is located about 18 km far from LNGS-INFN. After 6th April, 2009 the seismicity continued and migrated to the N and SE sectors, with 2 aftershocks of $M_w=5.2$ and $M_w=5.4$. Until 31st May 2010 more than 15,000 events with magnitudes greater than 1.0 with depth between 6-20 km have been recorded in the area 42.00° - 42.75° N and 12.75° - 13.75° E.

Simultaneous measurements of the following parameters were performed for each water sample at the monitoring sites inside the LNGS-INFN: U, sodium (Na), magnesium (Mg), potassium (K), calcium (Ca), Electrical Conductivity (EC), Oxidation-Reduction Potential (ORP), and pH. Some 110 parallel readings were recorded and each series was normalized to the corresponding average.

The U, Na, Mg, K and Ca readings show a long-term variability, around the unit value, limited to $\pm 20\%$. In addition to this background, in the period A, from 23rd June 2008 to 31st March 2009, these readings show a clear short-term 40-80% peak structure. In this period, relative to site E3dx for example, the experimental relative standard deviations (RSD) are about 20-25% for U, Na, Mg and Ca, and 35% for K that shows the largest peaked short-term variations, as well. Readings of EC, ORP and pH show relatively smooth fluctuations within $\pm 10\%$. RSD values vary between 2% and 5% and these quantities are excluded from further analysis.

Readings of Na, Mg, Ca and K were all normalized to the relevant U readings, in order to remove the common peak structure. The RSD of these ratios are 10% (Na), 9% (Mg), 7% (Ca) and 3% (K). Considering the Na/U ratio, with RSD of 0.10 and the two individual Na and U variability, both about 0.24, the Na-U covariance was estimated to be about 0.05. This is confirmed by the experimental covariance between each reading and the U ones, obtained from the set of data: 0.049 (Na), 0.046 (Mg), 0.047 (Ca). It is then evident that U, Na, Mg, Ca and K readings are correlated (Table 1).

A complete different situation can be observed in period B, from 10th April 2009 to 31st May 2010. The systematic peak structure for U, Na, Mg, Ca and K readings is no longer present, and only few isolated peaks of K and Na can now be observed. This is confirmed by the statistics reported in (Table 1). As an example, relative again to site E3dx, the RSD are now reduced to about 5% for U, Mg and Ca, and 15% for K and Na. These values agree well with the corresponding RSD of normalized (to U) readings in period A, typical of the constant background variability of periods A and B.

Period B readings of EC, ORP and pH show a similar behavior in period A, with RSD values between 2% and 5%. In both periods the pH reported the minimum RSD of 1.7% and 1.1%. The covariance analysis confirms this finding. Indeed, because of the loss of the peak structure, the correlation coefficients between each reading and the U ones are reduced from 0.88 (Na), 0.91 (Mg), 0.75 (K), 0.93 (Ca) to 0.31 (Na), 0.50 (Mg), 0.19 (K), 0.56 (Ca).

The water-rock interaction, modulated by percolation processes (due to meteoric events)

above the water table of the Gran Sasso aquifer, do not justify the U anomalies (22). Therefore, the progressive marked increase of U, along with the enrichment in calcium (Ca), magnesium (Mg), and sodium (Na) observed in groundwater located close to the main fault, crossing the deep underground LNGS-INFN during the preparation phases of the recent Aquila earthquake of 6th April 2009, is interpreted as chemical interactions with endogenic, upper mantle-derived carbon dioxide (CO₂) fluids, generated by recycling and melting of carbonaceous sediments during the ongoing subduction of the continental Adriatic plate beneath the Italian peninsula (23).

Recent evidences (24) indicate a shallow upper mantle convection/circulation (25) related to metasomatism and melting, and confirm the “west to east” flow of the mantle relative to the lithosphere, as suggested by the Apennines slab eastward retreat (26). This flow is consistent with the shear-wave splitting analysis that indicates an E–W upper mantle anisotropy (27, 28), due to the olivine crystals preferred orientation induced by mantle flow. The changes in the crustal stress pattern and stress magnitude are likely to be caused by buoyancy driven mantle circulation in the region rather than by gravitational potential energy differences (due to lateral density variations) in the crust itself (29). The combined analysis of petrological, geochemical and geophysical data reveals a surprisingly consistent picture showing that variations of seismic waves velocity in the mantle could be related to compositional differences of mantle sources, and not only to pressure and temperature variations (30). Petrological and geophysical modeling (31) explains how carbon is efficiently cycled in the upper mantle beneath Italy and the Western Mediterranean region at the Ma scale, via low-fractions of carbonate melts (about 0.1 wt%) generated by melting of carbonate-rich lithologies of the subducted Adria lithosphere, induced by the progressive rise of mantle temperatures behind the eastward-retreating subducting plate. The extraction of small-degree carbonate melts produces an abundant source of metasomatic fluids. Fluxes of carbonate melts are Na-bearing dolomitic (MgCaCO₃) in composition and contain from a hundred up to a thousand times the concentration in noble gases, heat-producing elements [e.g., U, thorium (Th), and potassium (K)] and other very incompatible elements [caesium (Cs), rubidium (Rb), and barium (Ba)] than the primitive mantle (23).

Experimental petrology data (32) indicate that the generation of carbonate-rich melts beneath the western Mediterranean and Italy commences at depths of 150-120 km. Due to their low density and viscosity, such melts are extremely mobile and migrate upward through the mantle, forming a carbonated partially molten CO₂-rich mantle, recorded by tomographic images between 120 and 60 km depth (30, 31). The upwelling of carbonate-rich melts to depths less than 60-70 km (i.e., below their P-T stability field) induces a massive outgassing of CO₂ in the lithosphere. Buoyancy forces, probably favored by fluid overpressures, are able to allow its migration to the exosphere through deep lithospheric faults, especially in the zones of thinned continental crust (31) (e.g., “non volcanic” CO₂ soil degassing in Tuscany and Latium).

However, not all of the CO₂ formed at mantle depth may reach the surface: some could remain trapped beneath the Moho and within the lower crust, in zones characterized by a thickened continental crust and lithosphere. Crustal CO₂ confinement might naturally give rise to over-pressurized reservoirs enriched in rare gases, Th, U and K, and other very incompatible elements, which are known to facilitate, in normal fault areas, seismogenesis, like the Colfiorito 1997 earthquake (33, 34).

The mechanic of faults is strongly influenced by fluids, their pore pressure, chemistry (e.g., CO₂, H₂O, etc.) and the resulting variation of friction along the fault plane (34, 35). Fluids may be actively triggering the rupture, and they are passively squeezed out during seismic events. Mantle CO₂ confined in the lower-middle crust is a chemically powerful agent for shallow aquifers. Discontinuous inputs of very low amounts (in the order of 1 mole % or less), induced by the progressive increase of CO₂ fluxes in zones of active faulting, are sufficient to give the aquifers the capability to dissolve carbonate bed/wallrocks, which should result in a spiked load of Mg and Ca, associated to an enrichment in U and other incompatible elements, similar to the geochemical patterns evidenced by the monitoring discussed here.

GPS data (36) show that, at global scale, a large part of deformation is absorbed by viscous shear in the lower crust. Such steady-state deformation contrasts with the stick-slip behavior of faults such as the one that generated the April 6th, 2009 event. Therefore, during the interseismic period, the lower crust, below the brittle-ductile transition, is constantly sheared, while faults in the upper crust are locked. During this time interval, the volume of rocks located at the brittle-ductile transition should suffer dilatancy, generated by the differential motion of the lower crust relative to the inactive upper crustal segment of the fault. Fluids are expected to fill in the fractures formed during the interseismic period. The separation produces a weak volume of rocks that sustain the hanging wall of the normal fault only until the load of the brittle upper crust will not overtake the strength of the dilated rocks. At that point, the hanging wall suddenly falls down to refill the stretched volume. This process implies the increase in pore pressure and the squeezing out of fluids. The hanging wall is expected to gradually adjust to its new position, generating a long sequence of aftershocks as those observed after the M_w 6.3 April 6th, 2009 L'Aquila earthquake. Some 15 km SW from the fault zone where LNGS-INFN is located, the area is cross-cut by several NW-SE trending active normal faults (37) cross-cutting the carbonatic massif of the Gran Sasso, where a sudden increase of fluid discharge has been observed during the main earthquake.

Although U concentrations and ²³⁴U/²³⁸U isotope ratios have been monitored in thermal waters (38-40), in order to assess the validity of U isotopes as fluid phase earthquake precursors, this monitoring at LNGS-INFN was performed in a shallow aquifer with a high dynamic behavior due to the high permeability of the Mesozoic limestones that form part of the Gran Sasso massif (17).

Conclusions

We propose that the U groundwater anomalies observed before the seismic swarm and the main shock, which occurred on 6th April, 2009 in L'Aquila, provide a key geochemical signal of a progressive increase of deep CO₂ fluxes at middle-lower crustal levels. Repeated sharp U enrichments in groundwater, that can be directly associated with the geodynamics of the earthquake, represent a much more precise strain-meter than Rn, whose presence could be modulated by U content during the preparation phase of the earthquake, and only successively released by microfracturing, during the main shock and aftershocks. The U groundwater anomalies observed before the L'Aquila earthquake will trigger further research on geodynamics and hydrogeochemistry of earthquakes, important for the better understanding of preparatory and final stages of earthquakes, and for possible routine use of groundwater earthquake precursors.

Acknowledgments

The authors greatly acknowledge the support by National Scientific Committee Technology of the National Institute of Nuclear Physics and the Chemistry and Chemical Plants Service of the Gran Sasso National Laboratory, and the Istituto Nazionale di Geofisica e Vulcanologia – Italian Seismic Bulletin for the seismic data.

Author Contributions

W.P. conceived the study, designed/coordinated the experiment. F.B., M.L.C., P.D.F., M.L., W.P. and P.P.P. provided comments on the manuscript for nuclear measurements and metrology. C.D. provided comments on the manuscript for geodynamic modeling. M.L.F. and A.P. provided comments on the manuscript for Earth carbon dioxide degassing and volcanism modeling. G.F.P. provided comments on the manuscript for tectonophysics modeling. W.P. and P.D.F. analyzed the data. P.A. and M.B. supported the logistic. L.I. performed groundwater sampling and assisted in ICP-MS measurements. S.N. performed the ICP-MS measurements. W.P., G.F.P., C.D., M.L.F., A.P., and P.D.F. wrote the paper.

Materials and Methods

Sampling. Groundwater to be analyzed flows in pipes introduced into the rock up to a depth ranging from three to six meters. Water samples were collected weekly in four sites located inside the LNGS-INFN. Each one-liter sample was stored, in cleaned and rinsed polyethylene bottles, after five minutes of water flushing at maximum flow.

ICP-MS. The water samples were diluted 10 times and acidified with 2.5% of nitric acid to stabilize traces in the sample. During sampling reagents of trace analysis grade (HNO_3 super pure by Carlo Erba® Reagenti), ultra pure water (produced by Millipore MilliQ®-Element), plastic containers and ancillary equipment were used because long-lived radionuclides Th and U at trace levels had to be measured. The ICP-MS measurements were carried out using a quadrupole mass spectrometer from Agilent® Technologies, model 7500a. The tuning of the instrument was optimized in order to reach high sensitivity, stable signal, and low background. A Babbington nebulizer was used during the measurements. The concentration values were determined in quantitative mode using an external calibration curve, because the matrix effect of the ten times diluted groundwater samples is negligible. During the measurements a multi-element solution was used as the internal standard to correct for possible instability and drift of the ICP-MS device. This solution was added on line using the third line of the peristaltic pump. The calibration curve of U response was corrected using ^{209}Bi . The accuracy reached in this way, as a rule, was better than 5%, which is reasonable for the ICP-MS technique.

References

1. Geller RJ, et al. (1997) Earthquakes cannot be predicted. *Science* 275:1616–1617.
2. Wyss M (1997) Cannot earthquakes be predicted?. *Science* 278:487–488.
3. Nature debates (1999)
http://www.nature.com/nature/debates/earthquake/equake_frameset.html.
4. Peresan A, et al. (2005) Intermediate-term middle-range earthquake predictions in Italy: a review. *Earth Sci. Rev.* 69:97–132.
5. Hauksson E (1981) Radon content of groundwater as an earthquake precursor: evaluation of worldwide data and physical basis. *J. Geophys. Res.* 86:9397-9410.
6. Roeloffs E (1999) Radon and rock deformation. *Nature* 399:104-105.
7. Ulomov VI, Mavashev B Z (1968) On a precursor of a strong tectonic earthquake. *Dokl. Akad. Sci. USSR, Earth Sci. Sec.* 174:9–11.
8. Wakita H, Nakamura Y, Notsu K, Noguchi M, Asada T (1980) Radon anomaly: a possible precursor of the 1978 Izu-Oshima-kinkai earthquake. *Science* 207:882-883.
9. King CY (1981) Do radon anomalies predict earthquakes? *Nature* 293:262.
10. Wakita H, Igarashi G, Nakamura Y, Sano Y, Notsu K (1989) Coseismic radon changes in groundwater. *Geophys. Res. Lett.* 16:417-420.
11. Torgersen T, Beniot J, Mackie D (1990) Controls on groundwater Rn-222 concentrations in fractured rock. *Geophys. Res. Lett.* 17:845–848.
12. Monnin MM, Seidel JL (1992) Radon in soil-air and groundwater related to major geophysical events: a survey. *Nucl. Instr. Meth. A* 314:316-330.
13. Virk HS, Singh B (1994) Radon recording of Uttarkashi earthquake. *Geophys. Res. Lett.* 21:737–740.
14. Igarashi G, et al. (1995) Groundwater radon anomaly before the Kobe earthquake in Japan. *Science* 269:60-61.
15. King CY, Koizumi N, Kitagawa Y (1995) Hydrogeochemical anomalies and the 1995 Kobe earthquake. *Science* 269:38–39.
16. Trique M, Richon P, Perrier F, Avouac JP, Sabroux JC (1999) Radon emanation and electric potential variations associated with transient deformation near reservoir lakes. *Nature* 399:137-141.
17. Plastino W, Bella F (2001) Radon groundwater monitoring at underground laboratories of Gran Sasso (Italy). *Geophys. Res. Lett.* 28:2675-2678.
18. Richon P, et al. (2003) Radon anomaly in the soil of Taal volcano, the Philippines: a likely precursor of the M 7.1 Mindoro earthquake (1994). *Geophys. Res. Lett.* 30:1481-1484.
19. Plastino W, Kaihola L, Bartolomei P, Bella F (2001) Cosmic background reduction in the radiocarbon measurements by liquid scintillation spectrometry at the underground laboratory of Gran Sasso. *Radiocarbon* 43:157-161.
20. Plastino W, et al. (2007) Tritium in water electrolytic enrichment and liquid scintillation counting. *Radiat. Meas.* 42:68-73.

21. Plastino W, et al. (2009) Environmental radioactivity in the ground water at the Gran Sasso National Laboratory (Italy): a possible contribution to the variation of the neutron flux background. *J. Radioanal. Nucl. Chem.* 282:809-813.
22. Plastino W, et al. (2010) Uranium groundwater anomalies and L'Aquila earthquake, 6th April 2009 (Italy). *J. Environ. Radioact.* 101:45-50.
23. Peccerillo A (2005) Plio-Quaternary volcanism in Italy. Petrology, geochemistry, geodynamics. *Springer, Heidelberg*, 2005:1-365.
24. Panza GF, Raykova RB, Carminati E, Doglioni C (2007) Upper mantle flow in the western Mediterranean. *Earth. Planet. Sci. Lett.* 257:200-214.
25. Faccenna C, Jolivet L, Piromallo C, Morelli A (2003) Subduction and the depth of convection in the Mediterranean mantle. *J. Geophys. Res.* 108:2099-2111.
26. Gueguen E, Doglioni C, Fernandez M (1998) On the post-25 Ma geodynamic evolution of the western Mediterranean. *Tectonophysics* 298:259–269.
27. Margheriti L, Lucente FP, Pondrelli S (2003) SKS splitting measurements in the Apenninic–Tyrrhenian domain (Italy) and their relation with lithospheric subduction and mantle convection. *J. Geophys. Res.* 108:2218-2237.
28. Barruol G, Deschamp A, Coutant O (2004) Mapping upper mantle anisotropy beneath SE France by SKS splitting indicates Neogene asthenospheric flow induced by Apenninic slab rollback and deflected by the deep Alpine roots. *Tectonophysics* 394:125–138.
29. Ismail-Zadeh A, Aoudia A, Panza GF (2010) Three-dimensional numerical modeling of contemporary mantle flow and tectonic stress beneath the Central Mediterranean. *Tectonophysics* 482:226-236.
30. Panza GF, Peccerillo A, Aoudia A, Farina B (2007) Geophysical and petrological modeling of the structure and composition of the crust and upper mantle in complex geodynamic settings: the Tyrrhenian Sea and surroundings. *Earth Sci. Rev.* 80:1-46.
31. Frezzotti ML, Peccerillo A, Panza GF (2009) Carbonate metasomatism and CO₂ lithosphere-asthenosphere degassing beneath the Western Mediterranean: an integrated model arising from petrological and geophysical data. *Chem. Geol.* 262:108-120.
32. Thomsen TB, Schimdt MW (2008) Melting of carbonated pelites at 2.5–5.0 GPa, silicate-carbonatite liquid immiscibility, and potassium–carbon metasomatism of the mantle. *Earth Planet. Sci. Lett.* 267:17–31.
33. Chimera G, Aoudia A, Saraò A, Panza GF (2003) Active tectonics in Central Italy: constraints from surface wave tomography and source moment tensor inversion. *Phys. Earth Planet. Int.* 138:241–262.
34. Miller S, et al. (2004) Aftershock driven by a high pressure CO₂ source at depth. *Nature* 427:724-727.
35. Sibson RH (1992) Implications of fault-valve behaviour for rupture nucleation and recurrence. *Tectonophysics* 211:283-293.

36. Kreemer C, Holt WE, Haines AJ (2002) The global moment rate distribution within plate boundary zones. *Geodynamics Series* 30:173-189.
37. Vezzani L, Ghisetti F (1998) Carta Geologica dell'Abruzzo - scala 1/100.000. *Regione Abruzzo-Selca*.
38. Kuleff I, Petrov P, Kostadinov K (1980) Uranium content variations in thermal waters from the west Rhodope crystalline massif during earthquakes in the Chepino valley (south Bulgaria). *J. of Rad. Chem.* 58:267–274.
39. Finkel RC (1981) Uranium concentrations and $^{234}\text{U}/^{238}\text{U}$ activity ratios in fault associated groundwater as possible earthquake precursors. *Geophys. Res. Lett.* 8:453–456.
40. McCaig A (1989) Fluid flow through fault zones. *Nature* 340:600.

Table 1. Summary of statistical analysis (Relative Standard Deviation, Mean, Covariance and Correlation with U) for each site and simultaneous readings of U, Na, Mg, K, Ca, EC, ORP, and pH, during the periods A (from 23rd June 2008 to 31st March 2009) and B (from 10th April 2009 to 31st May 2010). Each series was normalized to the corresponding average.

		A								B								
		U	Na	Mg	K	Ca	EC	ORP	pH	U	Na	Mg	K	Ca	EC	ORP	pH	
E1	RSD	0.145	0.148	0.134	0.272	0.130	0.026	0.063	0.024	0.041	0.036	0.035	0.088	0.036	0.026	0.054	0.008	
	MEAN	1.073	1.079	1.015	1.081	1.030	0.967	0.987	1.014	0.956	0.952	0.991	0.952	0.982	1.020	1.008	0.991	
	COV (UX)	0.020	0.019	0.017	0.027	0.016	0.000	-0.002	-0.001	0.002	0.001	0.000	0.000	0.000	0.000	0.000	0.000	0.000
	CORR (UX)	1.000	0.935	0.877	0.714	0.886	-0.095	-0.236	-0.174	1.000	0.400	0.181	0.141	0.309	0.002	0.173	-0.102	
E3	RSD	0.139	0.158	0.137	0.419	0.132	0.024	0.047	0.025	0.049	0.040	0.034	0.108	0.038	0.021	0.057	0.012	
	MEAN	1.072	1.075	1.022	1.105	1.040	0.973	1.038	1.002	0.957	0.956	0.986	0.928	0.977	1.016	0.977	0.999	
	COV (UX)	0.019	0.019	0.016	0.029	0.017	0.000	0.000	0.000	0.002	0.000	0.000	0.000	0.000	0.000	0.000	0.000	
	CORR (UX)	1.000	0.874	0.851	0.517	0.837	-0.062	-0.033	0.023	1.000	0.115	0.074	-0.010	-0.081	0.138	-0.148	-0.033	
E3dx	RSD	0.241	0.240	0.221	0.366	0.219	0.022	0.046	0.017	0.056	0.132	0.043	0.157	0.040	0.026	0.053	0.011	
	MEAN	1.134	1.099	1.066	1.143	1.092	0.987	1.032	1.003	0.920	0.940	0.961	0.919	0.946	1.008	0.982	0.998	
	COV (UX)	0.057	0.050	0.047	0.065	0.048	0.000	0.000	-0.001	0.003	0.002	0.001	0.002	0.001	0.000	0.001	0.000	
	CORR (UX)	1.000	0.884	0.906	0.752	0.831	-0.019	0.019	-0.130	1.000	0.309	0.496	0.193	0.554	0.226	0.302	-0.460	
E4	RSD	0.183	0.171	0.150	0.403	0.165	0.030	0.053	0.021	0.069	0.045	0.041	0.103	0.040	0.023	0.062	0.021	
	MEAN	0.987	1.069	1.033	1.175	1.051	0.979	1.076	0.999	1.009	0.959	0.980	0.904	0.970	1.013	0.955	1.001	
	COV (UX)	0.033	0.024	0.023	0.046	0.026	0.000	0.002	0.000	0.005	0.002	0.002	0.003	0.001	0.001	0.001	0.000	
	CORR (UX)	1.000	0.775	0.872	0.645	0.896	0.080	0.267	-0.127	1.000	0.624	0.554	0.419	0.470	0.356	0.271	-0.158	




Figure 1. Time variation of U concentration in groundwater sampled at E1, E3, E3dx, and E4 from June 2008 to May 2010. In the period A, from 23rd June 2008 to 31st March 2009, these readings show a clear short-term 40-80% peak structure. In the period B, from 10th April 2009 to 31st May 2010, the systematic peak structure for U is no longer present.

A seismic swarm started in the area under investigation, whose main shock occurred at 01:33 UT on April 6th, 2009. The cumulative seismic moment (CSM) from January 2008 to May 2010 in the area 42.00°-42.75° N and 12.75°-13.75° E has been estimated from seismic data of the Italian Seismic Bulletin, Istituto Nazionale di Geofisica e Vulcanologia: the clear jump in the CSM coincides with the main shock on April, 6th, 2009.




Figure 2. Present-day architecture of mantle-CO₂ degassing beneath L'Aquila (Central Italy).

The generation and evolution of deep mantle-derived CO₂ (A) has been delineated integrating surface wave tomography (B) with experimentally determined melting relationships for carbonated peridotite and crustal lithologies, relevant to recent mantle processes in the western Mediterranean region (C) (23, 30, 31). (A) Geochemical reconstruction (0-200 km depth) beneath the Latium - Abruzzo (L'Aquila) region (Central Italy). (B) Models (24) of shear-wave velocities (km s⁻¹) vs depth for Latium (b2) and Abruzzo (L'Aquila) zones (b3); the shadowed areas indicate the range of variability of the thickness of the layers, the red dashed lines mark the range of variability of the Moho depth; this representation is used to evidence that the boundaries between layers can well be transition zones in their own right. (C) Pressure-temperature diagram showing the effects of CO₂ on upper mantle lithologies (31). Peridotite - CO₂ *solidus*; Peridotite - CO₂ and Dry Peridotite *solidi* in the CMSA system; Eclogite - CO₂ *solidus*. KNCFMASH - CO₂ *solidus* (carbonated pelite + 1.1 wt.% H₂O + 4.8 wt.% CO₂). Blue area = estimated standard present-day mantle temperatures at the inferred pressures.

Melting of sediments and/or continental crust of the subducted Adriatic lithosphere at pressures greater than 4 GPa (120 km) and temperatures of 1200 °C (C - blue area) generate carbonate-rich melts. Such melts, migrating upward through the mantle, form a carbonated partially molten layer recorded by tomographic images between 70 and 160 km of depth (B - cell b2) (30). Further upwelling of carbonate-rich melts induces massive outgassing of CO₂ in the lithospheric mantle (C - carbonate - CO₂ field boundary).

High velocities in the uppermost mantle beneath Abruzzo (L'Aquila) support evidence for a LID initiating to subduct toward the west. Beneath Latium, at the same depth, an asthenospheric mantle wedge is present, as indicated by the quite low V_s between 40 and 80 km of depth (LID; B - cell b3) (30).