

---

NUSKY 11, Trieste June 22, 2011

# Restricting cosmogenic neutrino fluxes with Fermi 3 years data

Oleg Kalashev (INR RAS)  
Dmitry Semikoz (APC Paris)  
Graciela Gelmini (UCLA)

---

---

NUSKY 11, Trieste June 22, 2011

# Restricting cosmogenic neutrino fluxes with Fermi 2.5 years data

Oleg Kalashev (INR RAS)  
Dmitry Semikoz (APC Paris)  
Graciela Gelmini (UCLA)

---

\* e-mail: [kalashev@ms2.inr.ac.ru](mailto:kalashev@ms2.inr.ac.ru)

# Restricting cosmogenic neutrino fluxes

## Some works on the subject:


- V. S. Berezinsky and A. Yu. Smirnov, *Astrophys. Sp. Sci.* 32 461 (1975)
- M. Ahlers, L. A. Anchordoqui, M. C. Gonzalez-Garcia, F. Halzen and S. Sarkar, *Astropart. Phys.* 34, 106 (2010) [arXiv:1005.2620 [astro-ph.HE]]
- V. Berezinsky, A. Gazizov, M. Kachelriess and S. Ostapchenko, *Phys. Lett. B* 695, 13 (2011) [arXiv:1003.1496 [astro-ph.HE]]
- X. Wang, R. Liu, F. Aharonian, arXiv:1103.3574 [astro-ph.HE]

# Restricting cosmogenic neutrino fluxes with Fermi 2.5 years data


## *Overview*

- Propagation of Ultra High Energy Cosmic Rays (UHECR)
- Fitting HiRes spectrum by models with proton primaries
- Secondary photons and limit on the diffuse gamma ray flux from FERMI LAT
- Secondary neutrino fluxes
- Conclusion


# Main Factors influencing UHECR and $\gamma$ -ray propagation



# Main Factors influencing UHECR and $\gamma$ -ray propagation



# Attenuation lengths



# Simulations of cosmic rays propagation

- Monte Carlo based simulations
  - Random extragalactic magnetic field is taken into account
- Transport equation approach (rectilinear propagation)
  - Fast calculation (good for parameter space scanning)
  - Gives reasonable result for homogeneous source distribution with density  $n \geq 10^{-4} Mpc^{-3}$  for energies

$$E \geq 10^{18} eV \times Z \times \frac{B}{10^{-10} G}, \quad L_{cor} = 1 Mpc$$



# Fitting experimental data

- Energy spectrum  $j(E)$ 
  - Binned maximum likelihood function is used
  - Poisson probability of the observed event set is maximized

$$L(\mathbf{n}; \boldsymbol{\nu}) = \prod_i^N \frac{\nu_i^{n_i}}{n_i!} e^{-\nu_i}$$

- Goodness of fit defined as fraction of hypothetical experiments which result in worse agreement with the theory than the real data having the same total number of events

## Phenomenological source model:

$$F(E, z) = \Phi(E) S(z); \quad \Phi(E) = f E^{-\alpha} \text{Exp}(-E/E_{max}) \text{Exp}(-E_{min}/E)$$
$$S(z) = (1+z)^m \Theta(z-z_{min}) \Theta(z_{max}-z)$$

$z$  – red shift,  $\Theta(x)$ -step function

# Phenomenological source model:

$$F(E, z) = \Phi(E) S(z)$$

$$\Phi(E) = f E^{-\alpha} \text{Exp}(-E/E_{\max}) \text{Exp}(-E_{\min}/E)$$


$$S(z) = (1+z)^m \Theta(z-z_{\min}) \Theta(z_{\max}-z)$$

$z$  – red shift,  $\Theta(x)$ -step function

Parameter	Name	Typical Values
Power of the Injection Spectrum, $E^{-\alpha}$	$\alpha$	$2 \leq \alpha \leq 2.7$
End point of the Energy Spectrum	$E_{\max}$	$10^{20} \leq E_{\max}/\text{eV} \leq 10^{21}$
Evolution factor: $(1+z)^{3+m}$	$m$	$0 \leq m \leq 5$
Red shift of the nearest source	$z_{\min}$	$0 < z_{\min} < 0.01$
Maximal source redshift	$z_{\max}$	$z_{\max} = 2$
Minimal injection spectrum energy	$E_{\min}$	$10^{17}\text{eV}$

# Goodness of fit plots HiRes Spectrum


$$E_{fit} > 10^{18} \text{ eV}$$



$$z_{min} = 0.00125 \quad (\sim 5 \text{ Mpc})$$


# HiRes Spectrum fit example

$$E_{fit} > 10^{18} \text{ eV}$$



# Goodness of fit plots HiRes Spectrum

$$E_{fit} > 10^{18} \text{ eV}$$



Restricting cosmogenic neutrino fluxes with Fermi 2.5 years data

# Sample source evolution models

**Star Formation Rate:** H. Yuksel, M. D. Kistler, J. F. Beacom and A. M. Hopkins, *Ap. J.* 638 L5 (2008)


$$S_{SFR}(z) \propto \begin{cases} (1+z)^{3.4}, & z < 1 \\ (1+z)^{-0.3}, & 1 < z < 4 \\ (1+z)^{-3.5}, & z > 4 \end{cases}$$

**GRB:** H. Yuksel and M.D. Kistler *Phys. Rev. D* 75, 083004 (2007)

$$S_{GRB}(z) \propto \begin{cases} (1+z)^{4.8}, & z < 1 \\ (1+z)^{1.1}, & 1 < z < 4 \\ (1+z)^{-2.1}, & z > 4 \end{cases}$$

**AGN:** G. Hasinger, T. Miyaji, M. Schmidt, *Astron. and Astroph.* 441 417 (2005);  
M. Ahlers, L. A. Anchordoqui and S. Sarkar, *Phys. Rev. D* 79, 083009 (2009)

$$S_{AGN}(z) \propto \begin{cases} (1+z)^{5.0}, & z < 1.7 \\ \text{constant}, & 1.7 < z < 2.7 \\ 10^{(2.7-z)}, & z > 2.7 \end{cases}$$




# Sample source evolution models

$$z_{min} = 0.00125$$


**Star Formation Rate:** H. Yuksel, M. D. Kistler, J. F. Beacom and A. M. Hopkins, *Ap. J.* 638 L5 (2008)

$$S_{SFR}(z) \propto \begin{cases} (1+z)^{3.4}, & z < 1 \\ (1+z)^{-0.3}, & 1 < z < 4 \\ (1+z)^{-3.5}, & z > 4 \end{cases}$$



**GRB:** H. Yuksel and M.D. Kistler *Phys. Rev. D* 75, 083004 (2007)


$$S_{GRB}(z) \propto \begin{cases} (1+z)^{4.8}, & z < 1 \\ (1+z)^{1.1}, & 1 < z < 4 \\ (1+z)^{-2.1}, & z > 4 \end{cases}$$



**AGN:** G. Hasinger, T. Miyaji, M. Schmidt, *Astron. and Astroph.* 441 417 (2005);

M. Ahlers, L. A. Anchordoqui and S. Sarkar, *Phys. Rev. D* 79, 083009 (2009)

$$S_{AGN}(z) \propto \begin{cases} (1+z)^{5.0}, & z < 1.7 \\ \text{constant}, & 1.7 < z < 2.7 \\ 10^{(2.7-z)}, & z > 2.7 \end{cases}$$



# Fermi Gamma-ray Space Telescope



**Energy range (LAT):** 20 MeV - 300 GeV

**Field of view:** 20% of the sky at any instant; expose all parts of sky for ~30 minutes every 3 hours



# Fermi Gamma-ray Space Telescope




**Energy range (LAT):** 20 MeV - 300 GeV

**Field of view:** 20% of the sky at any instant; expose all parts of sky for ~30 minutes every 3 hours

**Launched** from Cape Canaveral Air Station  
11 June 2008

# Fermi Gamma-ray Space Telescope

## The Spectrum of the Isotropic Diffuse Gamma-Ray Emission



1 year data based bound:  
Fermi collaboration  
*Phys.Rev.Lett.* 104:101101,2010

2.5 years data based bound:  
*A.Neronov, D.V.Semikoz* 2011

# Interactions

## ■ Protons ,neutrons and nuclei

Pion production

$$p \gamma_b \rightarrow p \pi \dots$$

$e^+ e^-$  pair production

$$p \gamma_b \rightarrow p e^+ e^-$$

neutron  $\beta$ -decay

$$n \rightarrow p e^- \bar{\nu}_e$$


## ■ Electron-photon cascade

Inverse Compton

$$e \gamma_b \rightarrow e \gamma$$

$e^+ e^-$  pair production

$$\gamma \gamma_b \rightarrow e^+ e^-$$



# Interactions

## ■ Protons ,neutrons and nuclei

Pion production

$$p\gamma_b \rightarrow p\pi\dots$$

$e^+ e^-$  pair production

$$p\gamma_b \rightarrow p e^+ e^-$$

neutron  $\beta$ -decay

$$n \rightarrow p e^- \bar{\nu}_e$$

## ■ Electron-photon cascade

Inverse Compton

$$e\gamma_b \rightarrow e\gamma$$

$e^+ e^-$  pair production

$$\gamma\gamma_b \rightarrow e^+ e^-$$


Synchrotron losses

Double pair production

$$\gamma\gamma_b \rightarrow e^+ e^- e^+ e^-$$


$e^+ e^-$  pair production by  $e$

$$e\gamma_b \rightarrow e e^+ e^-$$



# Previous example fit with interaction products included

$$E_{fit} > 10^{18} \text{ eV}$$




$$\alpha = 2.45$$

$$m = 3.5$$

$$E_{max} = 10^{21} \text{ eV}$$

$$Z_{min} = 0.00125$$




# Fitting of the UHECR spectrum with Fermi bound imposed on secondary photon flux

$\chi^2$  statistics is used to obtain goodness of joint fit





Bins with small number of events are combined into larger bins

For Fermi bound only bins with photon flux exceeding bound are included in  $\chi^2$

Goodness of spectrum fit in the bins with small number of events is calculated separately on base of Poisson statistics



# Goodness of joint fit plots



$$\alpha = 2.45$$

$$m = 3.5$$

$$E_{max} = 10^{21} \text{ eV}$$

$$Z_{min} = 0.00125$$

# Sample source models

Star Formation Rate: Yuskel et al. (2008)

$$S_{SFR}(z) \propto \begin{cases} (1+z)^{3.4}, & z < 1 \\ (1+z)^{-0.3}, & 1 < z < 4 \\ (1+z)^{-3.5}, & z > 4 \end{cases}$$

GRB: Yuksel & Kistler (2007)


$$S_{GRB}(z) \propto \begin{cases} (1+z)^{4.8}, & z < 1 \\ (1+z)^{1.1}, & 1 < z < 4 \\ (1+z)^{-2}, & z > 4 \end{cases}$$

AGN: Hasinger et al. (2005)

$$S_{AGN}(z) \propto \begin{cases} (1+z)^{5.0}, & z < 1.7 \\ \text{constant}, & 1.7 < z < 2.7 \\ 10^{(2.7-z)}, & z > 2.7 \end{cases}$$

1 yr Fermi bound imposed

$z_{min} = 0$






# Energy scale shift as attempt to solve problem

Extra parameter:  $\Delta$ -energy scale shift for UHECR data

$$E \rightarrow E\Delta \quad 1.3^{-1} \leq \Delta \leq 1.3$$

# 1 yr Fermi

SFR




$$z_{min} = 0$$


$$E \rightarrow E\Delta$$

$$1.3^{-1} \leq \Delta \leq 1.3$$

GRB




AGN



1 yr Fermi

2.5 yr Fermi


SFR





$$z_{min} = 0$$

$$E \rightarrow E\Delta$$



$$1.3^{-1} \leq \Delta \leq 1.3$$



GRB



AGN




Restricting cosmogenic neutrino fluxes with Fermi 2.5 years data

---

## 95% CL secondary neutrino flux range calculation


- Using only models with goodness of fit  $P > 0.05$
- For each energy bin we find model predicting highest/lowest neutrino flux

# 95% CL secondary neutrino flux range calculation




Dependence on  $E_{\text{max}}$  for models with evolution  $S(z)=(1+z)^m$

# 95% CL secondary neutrino flux range calculation




# Photon diffuse flux dependence on EBL



IR/Optic background models used here:  
Old: F.Stecker et al. astro-ph/0510449  
New: Kneiske et al. astro-ph/1001.2132v1

# Comparison with other simulation



Reference:

M. Ahlers, L. A. Anchordoqui, M. C. Gonzalez-Garcia, F. Halzen and S. Sarkar, *Astropart. Phys.* 34, 106 (2010)

Fig. B7



---





# Conclusions

- Fermi LAT diffuse gamma ray flux limits become to constrain possible set of allowed UHECR source models and cosmogenic neutrino fluxes
- Pure proton source models with strong evolution are disfavored.
- Problems with strongly evolving sources can be avoided if one assumes nuclei primaries or local source overdensity


---

# Appendix

# $e^+e^-$ pair production on p is the main source of diffuse photon background



# Attenuation lengths (EM cascade)



# Simulations of cosmic rays propagation

Sample transport equation for electrons (includes only pair production PP and inverse Compton scattering ICS)

$$\begin{aligned} \frac{d}{dt} N_e(E_e, t) = & -N_e(E_e, t) \int d\epsilon n(\epsilon) \int d\mu \frac{1 - \beta_e \mu}{2} \sigma_{\text{ICS}}(E_e, \epsilon, \mu) + \\ & \int dE'_e N_e(E'_e, t) \int d\epsilon n(\epsilon) \int d\mu \frac{1 - \beta'_e \mu}{2} \frac{d\sigma_{\text{ICS}}}{dE_e}(E_e; E'_e, \epsilon, \mu) + \\ & \int dE_\gamma N_\gamma(E_\gamma, t) \int d\epsilon n(\epsilon) \int d\mu \frac{1 - \mu}{2} \frac{d\sigma_{\text{PP}}}{dE_e}(E_e; E_\gamma, \epsilon, \mu) + Q(E_e, t) \end{aligned}$$

---

# Interactions

- Protons and neutrons

  - Deflection by EGMF

$B < 10^{-9}$  G P. Kronberg, Rept. Prog. Phys. 57, 325 (1994).

$B > 10^{-17}$  G A. M. Taylor, I. Vovk, A. Neronov, Astronomy & Astrophysics, vol 529, 2011

$B \sim 10^{-12}$  G K. Dolag, D. Grasso, V. Springel and I. Tkachev, JCAP 0501, 009 (2005)

# Interactions

- Protons and neutrons

Pion production

$$N\gamma_b \rightarrow N'\pi\dots$$

$e^+e^-$  pair production

$$p\gamma_b \rightarrow pe^+e^-$$

neutron  $\beta$ -decay

$$n \rightarrow pe^-\bar{\nu}_e$$

# Interactions

## ■ Protons and neutrons

### Pion production

$$N\gamma_b \rightarrow N'\pi\dots$$

$$E_{th} = \frac{m_\pi(m_p + m_\pi/2)}{\epsilon} \simeq 7 \times 10^{16} \left(\frac{\epsilon}{eV}\right)^{-1} eV$$


For MWB ( $\epsilon \simeq 10^{-3} eV$ ):  $E_{th} \simeq 70 E eV$

SOPHIA event generator

*A.Mucke et al., Comp.Phys.Comm.124,290(2000)*

### $e^+e^-$ pair production

$$p\gamma_b \rightarrow pe^+e^-$$



### neutron $\beta$ -decay

$$n \rightarrow pe^- \bar{\nu}_e$$



# Interactions

## ■ Protons and neutrons

### Pion production

$$N\gamma_b \rightarrow N'\pi\dots$$

$$E_{th} = \frac{m_\pi(m_p + m_\pi/2)}{\epsilon} \simeq 7 \times 10^{16} \left(\frac{\epsilon}{eV}\right)^{-1} eV$$

For MWB ( $\epsilon \simeq 10^{-3} eV$ ):  $E_{th} \simeq 70 EeV$

SOPHIA event generator

*A.Mucke et al., Comp.Phys.Comm.124,290(2000)*

### $e^+e^-$ pair production

$$p\gamma_b \rightarrow pe^+e^-$$

$$E_{th} = \frac{m_e(m_A + m_e)}{\epsilon} \simeq 5 \times 10^{14} \left(\frac{\epsilon}{eV}\right)^{-1} eV$$


For MWB ( $\epsilon \simeq 10^{-3} eV$ ):  $E_{th} \simeq 5 \times 10^{17} eV$

*p energy loss rate:*

*M.J.Chodorowski et al. Astrophys.J.400,181(1992)*

### neutron $\beta$ -decay


$$n \rightarrow pe^-\bar{\nu}_e \quad \tau = 900s$$



# Some references on UHECR propagation

<b><math>\pi</math> production</b>	A.Mucke et al., Comp.Phys.Comm.124,290(2000)
<b>Photodisintegration</b>	F.Stecker et al. Astrophys.J. 512 (1999) 521-526. E.Khan et al. Astropart.Phys. 23 (2005) 191-201
<b><math>e^+e^-</math> pair production</b>	M.J.Chodorowski et al. Astrophys.J.400,181(1992)
<b>Extragalactic magnetic field</b>	K.Dolag et al., astro-ph/0410419  F.Stecker et al. astro-ph/0510449
<b>Infrared background</b>	Kneiske et al. astro-ph/1001.2132v1
<b>Radio background</b>	T.A. Clark, L.W. Brown, and J.K. Alexander, Nature 228, 847 R.J. Protheroe, P.L. Biermann, Astropart. Phys. 6, 45

# Energy loss lengths



$p$  and  $\gamma$  energy loss lengths  
(minimal RB assumed)

# Interactions

## ■ Electron-photon cascade

Inverse Compton

$e^+ e^-$  pair production

Synchrotron losses

Double pair production


$e^+ e^-$  pair production by  $e$

$$e \gamma_b \rightarrow e \gamma$$

$$\gamma \gamma_b \rightarrow e^+ e^-$$


$$\gamma \gamma_b \rightarrow e^+ e^- e^+ e^-$$

$$e \gamma_b \rightarrow e e^+ e^-$$



# Deflection and synchrotron radiation

Gyroradius:  $R_g = \frac{E}{qeB_{\perp}} \simeq 110 \times \frac{1}{Z} \left( \frac{E}{10^{19} \text{ eV}} \right) \left( \frac{B_{\perp}}{10^{-10} \text{ G}} \right)^{-1} \text{ Mpc}$



Synchrotron loss length:

$$\frac{dE}{dt} = -\frac{4}{3} \sigma_T \frac{B^2}{8\pi} \left( \frac{qm_e}{m} \right)^4 \left( \frac{E}{m_e} \right)^2$$

$$E_{\gamma} \simeq \frac{3eB}{2m_e} \left( \frac{E_e}{m_e} \right)^2 \simeq$$

$$2.2 \times 10^{14} \left( \frac{E_e}{10^{21} \text{ eV}} \right)^2 \left( \frac{B}{10^{-9} \text{ G}} \right) \text{ eV}$$

The gyroradius and the synchrotron loss rates of electrons for various strengths of the EGMF

# Observation of photons from distant blazars

HESS	1ES 1101-232 (z=0.186)	$\Gamma = 2.88 \pm 0.17$	$\frac{dN}{dE} \sim E^{-\Gamma}$
	H2356-309 (z=0.165)	$\Gamma = 3.06 \pm 0.21$	
	<i>Nature 440 (2006) 1018-1021</i>	$0.5 < E / \text{TeV} < 10$	
VERITAS	3C66A (z = 0.444)	$\Gamma = 4.1 \pm 0.4_{stat} \pm 0.6_{sys}$	
		<i>Astrophys. J. Lett. 693 (2009) L104-L108</i>	