

Unstable periodic orbits for the Navier-Stokes equations

Luis Fazendeiro¹, Bruce M. Boghosian²,
Peter V. Coveney¹, Jonas Lätt³ and Spencer Smith²

1. Centre for Computational Science, UCL, London, UK
2. Tufts University, Medford, USA
3. Ecole Polytechnique Federale de Lausanne, Switzerland

6th August 2009, ICTP, Trieste, Italy

2nd International Conference and Advanced School, Turbulent Mixing and Beyond


Contents

- 1 – Unstable Periodic Orbits**
- 2 – Dynamical Zeta Function**
- 3 – Lower-dimensional systems**
- 4 – Lattice Boltzmann Method**
- 5 – HYPO4D**
- 6 – Turbulent flow**
- 7 – 4D numerical relaxation**
- 8 – Prospects & Challenges**

1 – Unstable Periodic Orbits

- Role of UPO's has been acknowledged since the work of Poincaré (“founder” of modern dynamical systems theory);
- Typical trajectory will wander incessantly in a sequence of close approaches to the UPOs;
- Analogy from statistical mechanics in physics: set of UPO's can be viewed as the microstates from which a macroscopic description of the system can be calculated;
- Accuracy of predictions is limited by the (non-composite) UPO of smaller period which we fail to include;

1 – Unstable Periodic Orbits



- Abstract “dynamical landscape” with peaks representing the UPOs;
- Chaotic trajectory can be visualized as being the motion of a ball rolling on this abstract dynamical landscape;
- Motion will be strongly affected by the sharpness of the peak (stability of the UPOs);
- Strange attractor is closure of the set of all the UPOs (in the neighborhood of which the system will spend most of the time)

from: http://www.scholarpedia.org/article/Unstable_periodic_orbits

Curator: Dr. Paul So, George Mason University, Fairfax, VA

1 – Unstable Periodic Orbits

- Kawahara & Kida¹ found 2 periodic solutions in plane Couette flow, simulated using spectral methods;
- Novel efficient algorithm to locate UPOs (Boghosian *et al*, preprint), based on previous work by Lan and Cvitanovic²;
- Tested on Lorenz model and several low-dimensional systems; good convergence to the UPOs, including 2D fluid;
- Plot $\Delta(t,T)$: find initial guess (minimum) for whole trajectory, and value of T;

$$\Delta(t,T) = \left\| r(t+T) - r(t) \right\| \geq 0$$
- Numerically relax minima (in 4D) towards finding UPO;

2– Dynamical Zeta function

- In “traditional” fluid turbulence, observables are computed as averages over many time frames:

$$\langle A \rangle = \lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t A(\rho(\tau)) d\tau.$$

- Alternative approach would be to using a generating function:

$$s(\beta) = \lim_{t \rightarrow \infty} \frac{1}{t} \ln \langle e^{\beta \cdot A^t} \rangle,$$

$$\langle e^{\beta \cdot A^t} \rangle = \frac{1}{|\mathcal{M}|} \int_{\mathcal{M}} dx e^{\beta \cdot A(f^t(x))}.$$

- Then the moments of the PDF of an observable A are:


$$\left. \frac{\partial s}{\partial \beta} \right|_{\beta=0} = \langle A \rangle,$$

$$\left. \frac{\partial^2 s}{\partial \beta^2} \right|_{\beta=0} = \langle A^2 \rangle - \langle A \rangle^2$$

2 – Dynamical Zeta function

- DZF expression $\zeta^{-1}(s, \beta) = \lim_{T_{\max} \rightarrow \infty} \prod_{\{P(T < T_{\max})\}} 1 - e^{-sT_p + \beta A_p - \log(\Lambda_p)}$
 - product runs over all UPO's with period $T < T_{\max}$
 - T_p : period of the UPO.
 - A_p : average value of the observable on the UPO.
 - Λ_p : stability eigenvalue of the UPO.
- Generating function: $s = s(\beta) : \quad \zeta^{-1}(s(\beta), \beta) = 0.$
- Computation of expectation value $\left. \frac{\partial s}{\partial \beta} \right|_{\beta=0} = \langle A \rangle$

Example of a strange attractor: Lorenz system



$$\frac{dx}{dt} = \sigma(y - x)$$

$$\frac{dy}{dt} = x(\rho - z) - y$$

$$\frac{dz}{dt} = xy - \beta z$$

ρ = Prandtl number

σ = Rayleigh number


$\rho = 28$

$\sigma = 10$

$\beta = 8/3$

x: proportional to intensity of convection

y: proportional to temperature difference



SINGLE CONVECTION CELL



- each of the lobes corresponds to a steady state
- but there are transitions, including reversal of direction: hot air descends, cold air ascends!

“Deterministic nonperiodic flow”

E. N. Lorenz, J. Atmos. Sci. 20 130 (1963)

Picture: http://en.wikipedia.org/wiki/Lorenz_attractor

2 – Dynamical Zeta function



- Comparison between the 2 approaches for the Lorenz equations: exact polynomial expansion (left) versus “noisy” time-integration (right).

2 – Dynamical Zeta function

(Main) Advantages in this approach:

- Degree of accuracy is high and converges quickly, with the number of known lower period UPOs
- No need to redo initial value problem every time we wish to compute the average of some quantity
- Averages are no longer stochastic in nature, we have an exact expansion to compute them

3 – Lower-dimensional systems


Dynamical system :

$$\dot{\rho}(t) = f(\rho(t))$$

Lorenz equation


State ρ : point in 3D space

Orbit : Time-evolution of 3D point.




2D force-driven turbulent flow

State ρ : velocity distribution $u(x, y)$ at a given time (infinite-dimensional space).




Orbit : Time-evolution of Fluid.



3 – Lower-dimensional systems: 2D fluid



- Magnetohydrodynamic forcing in a thin layer of conducting fluid



- Fluid flow simulated using OpenLB, with a sinusoidal force, to reproduce experimental work (quasi-2D)¹. Colours represent the vorticity field: red is large negative value (counterclockwise), blue is large positive value (clockwise).

3 – Lower-dimensional systems: 2D fluid

- Magnetohydrodynamic forcing in a thin layer of conducting fluid




- Fluid flow simulated using OpenLB, with a sinusoidal force, to reproduce experimental work (quasi-2D)¹. Colours represent the vorticity field: red is large negative value (counterclockwise), blue is large positive value (clockwise).

3 – Lower-dimensional systems: 2D fluid revisited

- Same system as previous slide. Red denotes large negative vorticity (counterclockwise rotation), blue is large positive vorticity (clockwise rotation). Periodic solution shown here (found analytically) is highly unstable.

3 – Lower-dimensional systems: 2D fluid revisited



- Same system as previous slide. Red denotes large negative vorticity (counterclockwise rotation), blue is large positive vorticity (clockwise rotation). Periodic solution shown here (found analytically) is highly unstable.

4 – Lattice Boltzmann Method

(or why stream and collide is all it takes)

- Fluid flow (nearly incompressible NSE) is simulated using the Lattice Boltzmann Method^{1,2}

$$f_i(\vec{r} + \vec{c}_i, t + 1) = f_i(\vec{r}, t) + \Omega_i(\vec{r}, t),$$

$$\rho = \sum_{i=1}^b f_i, \rho \vec{u} = \sum_{i=1}^b f_i \vec{c}_i.$$

$$\sum_{i=1}^b \Omega_i = 0, \sum_{i=1}^b \Omega_i \vec{c}_i = \vec{0}.$$

$$f_i(\vec{r} + \vec{c}_i, t + 1) - f_i(\vec{r}, t) = \mathcal{S}_{ij}(f_j(\vec{r}, t) - f_j^{eq}(\vec{r})),$$

$$f_i(\vec{r} + \vec{c}_i, t + 1) - f_j(\vec{r}, t) = -\frac{1}{\tau}(f_i(\vec{r}, t) - f_i^{eq}(\vec{r})).$$




FIG. 1. Lattice structure of d3q15 (left) and d3q18 (right).

$$f_i^{eq}(\rho, \vec{u}) = \rho[a + b\vec{c}_i \cdot \vec{u} + c(\vec{c}_i \cdot \vec{u})^2 + du^2],$$

$$\sum_{i=1}^b f_i^{eq} = \rho, \sum_{i=1}^b f_i^{eq} \vec{c}_i = \rho \vec{u},$$

$$\nu = (\tau - \frac{1}{2})c_s^2 \Delta t,$$

1. S. Chen, G. D. Doolen, Annu. Rev. Fluid Mech., 30, 329-364, 1998



2. S. Succi, "The LBE for Fluid Dynamics and Beyond", Oxford University Press, Oxford, 2001

4 – Lattice Boltzmann Method

- Basic premise: macroscopic dynamics is the result of collective behavior of many microscopic ones and the macroscopic dynamics not sensitive to underlying details is microscopic physics;
- How did we go from discrete velocities to hydrodynamics? Multi-scale (Chapman-Enskog) expansion, assuming diffusion time scale much slower than convection time scale;
- Main areas of application: complex boundaries, interfacial dynamics, multiphase flows; Codes: OpenLB, LB3D (amphiphilic fluids), HemeLB (cerebral blood flow);
- Main advantages of LBM:
 - local collision operator + linear streaming operator (almost “embarrassingly” parallel);
 - minimal set of velocities, from which macroscopic quantities are computed;
 - no need for extra term for extra equation for pressure term, which so often requires extra numerical treatments (pressure is now eq. of state);

5 – HYPO4D¹

- Communication pattern between processors: only the halo values need to be sent to nearest neighbors →





- Scalability tests at NGS and HPCx (Gold Star award)

(i.e., ~1.9 speed-up for cubic lattice with $L=1024$, when going from 512 to 1024 processors)

5 – HYPO4D: Ranger + Intrepid scaling

- In the halo-exchange step, all MPI communications are non- blocking, in order to prevent dead-lock;
- No aggressive optimization pursued, so that code can be deployed (almost) seamlessly on different platforms;
- Ranger @TACC: Sun Constellation Linux Cluster, 62976 AMD cores, 123 TB memory, 579.4 TFlops (theoretical) peak;
- Intrepid @Argonne: IBM BlueGene/P, 163 840 cores, 80 TB memory, 557 TFlops (theoretical) peak;

Ranger scalability tests: Hard versus Soft scaling



- Linear scaling up to 16K on Ranger (left) and 33K on Intrepid (right). Also close to linear up to 33K on Ranger and 65K on Intrepid;


6 – Turbulent flow

- Current approach: fluid starts from rest, periodic boundary conditions in all directions (isotropic, homogeneous turbulence), ABC type force applied:

$$\begin{aligned}\vec{\nabla} \cdot \vec{F} &= 0 \\ \vec{\nabla} \times \vec{F} &= \vec{F}\end{aligned}$$

- Numerical stability tests applied at all sites in the lattice at every time step, convergence tests and other quantities measured at regular intervals;
- After a certain threshold of LB viscosity, (weak) turbulent behavior sets in;
- HYPO4D = initial value problem (LB) + minimum search + 4D numerical relaxation (variational principle applied to LBE) + post-processing tools;


6 - Turbulent flow: Convergence test



- Cubic lattice, 64^3 , varying kinematic (LB) viscosity;
- Maximum $Re \sim 500$, taking velocity averaged over many time, steps, but DNS, no modeling included;



- Convergence test for the velocity field, taken between consecutive time steps. Notice sharp increase after turbulent regime sets in (vertical axis scale is logarithmic).

6 - Turbulent flow: Energy spectrum $E(q)$




- Kolmogorov picture of turbulent flow divides it (roughly) into 3 different scales: kinetic, inertial, dissipative;


- Left plot shows energy spectrum $E(q)$ before transition to time-dependent behavior;



- Right plot shows spectrum for two different time snapshots after the transition, with Kolmogorov fitting for visual aid;




7 - 4D Numerical Relaxation: search for minima



- Vertical axis is T, horizontal axis is t, color code gives “delta” quantity:

$$\Delta(t, T) \equiv \sqrt{\sum_{\mathbf{r}} \sum_{i=1}^l (f_i(\mathbf{r}, t+T) - f_i(\mathbf{r}, t))^2},$$

7 - 4D Numerical Relaxation: search for minima




- Detail of previous plot, showing several (purplish) minima/areas of interest, as well as hinting at some regularity in their distribution

7 - 4D Numerical Relaxation: search for minima

- New minima for $\nu = 0.01$

Delta(t,T) - sampling rate of 500 LB time-steps - $\tau=0.53$



7 - 4D Numerical Relaxation

- Memory critical resource for the full 4D relaxation procedure;
- For the minimum shown $T \sim 26.5K$, we have to keep in memory at least $64^3 * 19$ (number of LB velocities) $* 8$ (double precision) $* 2.65 * 10^4$ variables ~ 1 TB; Then SD=5 copies and CG=8!!!!!!
- Minimization algorithm:

- define functional $\mathcal{F} \equiv \frac{1}{2} \sum_{t=0}^{T-1} \sum_{\vec{r}} \sum_{i=1}^b |\phi_i(\vec{r}, t)|^2$ with

$$\phi_i(\vec{r}, t) = f_i(\vec{r} + \vec{c}_i, t + 1) - f_i(\vec{r}, t) - \Omega_i(\vec{r}, t)$$

- compute gradient $\frac{\delta \mathcal{F}}{\delta f_k(\vec{s}, q)}$

7 - 4D Numerical Relaxation: algorithm

- Fill (4D) lattice with $f_i(\vec{r}, t)$ from $t = t_0, T - 1$
- 4D collide and stream (periodic time BC)
- Compute ϕ and initial \mathcal{F}
- Combination of SD and/or CG until \mathcal{F} no longer varies
- SD algorithm :


$$f_i(\vec{r}, t)_{-} = \alpha * \frac{\partial \mathcal{F}}{\partial f_i(\vec{r}, t)}$$

- α found through golden mean search procedure, done at every step


7 - 4D Numerical Relaxation: failed experiment :(

- Left: initial 26.5K trajectory;
Right: after SD minimization, and also time-forwarding from checkpoint on same machine;
- Y-axis: total energy in both cases;
- At every single SD step, \mathcal{F} decreases!

L=64, ABC force, visc = 0.01



L=64, ABC force, visc = 0.01



8 - Prospects & Challenges

- Relaxation procedure in order to identify first (smaller period) UPOs in 3D NSE;
- Characterization of the UPOs (stability eigenvalues, but also energy dissipation rate, vorticity, enstrophy, etc);
- Compare averages from one or more UPOs with time forward averaging;
- Possible heteroclinic connections between periodic solutions, along which bursting (intermittency) may occur (K&K);
- Classification of the (prime) UPOs identified + creation and maintenance of a digital library of such orbits;
- Could a symbolic dynamics be devised from some of these UPOs, for 3D NSE turbulence?

Acknowledgements

