

OUTLINE(1)

- CHAOS and its characterization (Lyapunov Exponents)
 - Classical Dynamical Systems (CDS)
 - An equivalent indicator of chaos for CDS:
 - the Kolmogorov–Sinai (metric) Entropy [Pesin Thm.]
-

- Algebraic description of CDS
- Systems with finite number of states:
 - Quantum Dynamical Systems (QDS)

ABSENCE OF CHAOS FOR THESE SYSTEMS

CHAOS over times windows (entropy production analysis)

- Quantization (Weyl and Anti–Wick)
- Classical Limit (of the phase–space quantization)
- Coherent States on the torus

OUTLINE(2)

- Quantum Dynamical Entropies (QDE)
 - ALF Entropy
 - CS Entropy
 - Relation between the QDE and the KS entropy
 - QDE show absence of chaos for systems with finite number of states
-

- Classical (Continuous) limit and Temporal Evolution

“““VIOLATION OF THE CORRESPONDENCE PRINCIPLE”””

BREAKING TIME

- Analytical and Numerical estimation of the Breaking Time by means of QDE production analysis
- CONCLUSION

The problem of defining quantum chaos


Classical Chaos: exponential amplification of small errors

$$|\delta x| \xrightarrow[n]{} |\delta x(n)| = \lambda^n |\delta x| = e^{n \log \lambda} |\delta x|$$

$\log \lambda$ is **Lyapunov Exponent**

$|\delta x(n)|$ increase no longer in **compact** systems...

$$\log \lambda := \lim_{n \rightarrow +\infty} \lim_{\delta x \rightarrow 0} \frac{1}{n} \log \frac{|\delta x(n)|}{|\delta x|}$$



... and $|\delta x|$ cannot be let go to zero in **quantum** or **discrete** systems!

Indeed:

- for a **discrete** system $|\delta x| > a$ ($a = \frac{1}{N}$ = lattice spacing);
- for a **quantum** system $|\delta x| > \frac{\hbar}{|\delta p|}$ ($|\delta p|$ bounded on **compact**).


$$\begin{aligned} \forall \hbar : \log \lambda(\hbar) = 0 \quad \text{but} \quad \lim_{n \rightarrow \infty} \lim_{\hbar \rightarrow 0} \log \lambda(\hbar) > 0 \\ \forall N : \log \lambda(N) = 0 \quad \text{but} \quad \lim_{n \rightarrow \infty} \lim_{N \rightarrow \infty} \log \lambda(N) > 0 \end{aligned}$$

An equivalent indicator of chaos in classical continuous systems

- **Entropy** $[S(n)]$: information on the evolving system up to time n .
- Loosely speaking, the **Kolmogorov-Sinai metric entropy** is
$$h_\mu(T) := \lim_{n \rightarrow \infty} \frac{S(n)}{n} \text{ (entropy per unit time).}$$

Theorem 1 (Pesin)

Ergodicity $\implies h_\mu(T) = \sum$ positive Lyapunov exponent



Classical Dynamical Systems (\mathcal{X}, μ, T)

$$\left\{ \begin{array}{l} \textcolor{red}{\mathcal{X}} : \text{Measurable space (Torus } \mathbb{T}^2 := \mathbb{R}^2 / \mathbb{Z}^2) \\ \textcolor{red}{\mu} : \text{normalized measure } (\mu(\mathcal{X}) = 1) \text{ (Lebesgue } \mu(d\mathbf{x}) = dx_1 dx_2) \\ \textcolor{red}{T} : \text{an invertible measurable map } T : \mathcal{X} \mapsto \mathcal{X} \text{ (described later)} \\ \mu\text{-invariant } (\mu \circ T = \mu) \end{array} \right.$$

Partition over \mathcal{X}

$$\textcolor{red}{\mathcal{E}} := \{\textcolor{red}{E}_\ell\}_{\ell=1,2,\dots,D} \quad \text{such that} \quad \left\{ \begin{array}{l} E_\ell \subset \mathcal{X} \\ \bigcup_{\ell=1}^D E_\ell = \mathcal{X} \\ E_\ell \cap E_k = \emptyset \quad , \quad \forall \ell \neq k \end{array} \right.$$



Evolved partition at time j

$$\textcolor{red}{T}^j(\mathcal{E}) := \{T^{-j}(E_\ell)\}_{\ell=1,2,\dots,D}$$

Definition

Ω_D^n is the set of strings $\textcolor{red}{i} := \{i_0, i_1, \dots, i_{n-1}\}$,
 i_j belonging to the alphabet $\{1, 2, \dots, D\}$

$$\mathcal{E} = \{E_{\text{Blue}}, E_{\text{Gray}}, E_{\text{Magenta}}, E_{\text{Yellow}}\}$$



All trajectories starting in $E_{\text{GYMBB}} := E_G \cap \dots \cap T^{-4}(E_B)$ are **encoded** by the same **string** $\{\text{GYMBB}\}$ (up to time 4)

Their **probability** is:
 $\mu_{\text{GYMBB}} := \mu(E_{\text{GYMBB}})$

Refined partition (up to time n)

$$\mathcal{E}_{[0,n-1]} := \{E_{\mathbf{i}}\}_{\mathbf{i} \in \Omega_D^n} \quad , \quad E_{\mathbf{i}} := \bigcap_{j=0}^{n-1} T^{-j}(E_{i_j})$$

strings $\mathbf{i} \in \Omega_D^n$ encode trajectories $\{T^k \mathbf{x}\}$, $\mathbf{x} \in E_{\mathbf{i}}$

Richness in trajectories of $E_{\mathbf{i}} \in \mathcal{E}_{[0,n-1]}$ is measured by the volume $\mu_{\mathbf{i}} := \mu(E_{\mathbf{i}})$.

Kolmogorov metric entropy

With the probabilities $\mu_{\mathbf{i}}$ we can compute the SHANNON ENTROPY

$$S_{\mu}(\mathcal{E}_{[0,n-1]}) := - \sum_{\mathbf{i} \in \Omega_D^n} \mu_{\mathbf{i}} \log \mu_{\mathbf{i}}$$

the entropy production per time step

$$h_{\mu}(T, \mathcal{E}) := \lim_{n \rightarrow \infty} \frac{1}{n} S_{\mu}(\mathcal{E}_{[0,n-1]})$$

and the KS entropy

$$h_{\mu}(T) := \sup_{\mathcal{E}} h_{\mu}(T, \mathcal{E})$$


The classical Dynamics: the Unimodular Group

$$\mathbb{T}^2 \ni \mathbf{x}_n \mapsto \mathbf{x}_{n+1} := \mathbf{T} \mathbf{x}_n \pmod{1} \in \mathbb{T}^2$$


$$T := \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \quad a, b, c, d \in \mathbb{Z}, \quad \det(T) = 1$$

\mathbf{T} is a toral automorphism and a generalization of the so called [Arnold Cat Map](#). Depending on $\text{Tr}(T)$ we have two kind of dynamics:

- $|\text{Tr}(T)| > 2$ [Chaotic Systems](#) $\log \lambda = \log \left(\frac{\text{Tr}(T) + \sqrt{\text{Tr}(T)^2 - 4}}{2} \right)$
- $|\text{Tr}(T)| \leq 2$ Regular Systems



Systems with finite number of states :: Chaos...



...but also Periodicity

Algebraic description of Dynamical Systems

1) Classical Dynamical Systems (CDS $(\mathcal{M}_\mu, \omega_\mu, \Theta)$)

$$\left\{ \begin{array}{l} \mathcal{M}_\mu : \text{Algebra of observables} \quad (\mathcal{C}^0(\mathbb{T}^2), L_\mu^\infty(\mathbb{T}^2)) \\ \omega_\mu : \text{State over } \mathcal{M}_\mu \quad (\omega_\mu(f) := \int_{\mathbb{T}^2} f(\mathbf{x}) d^2\mathbf{x}) \\ \Theta : \text{Discrete group of } \omega_\mu\text{-preserving} \quad (\Theta^k(f(\mathbf{x})) := f(T^{-k}\mathbf{x})) \\ \quad \text{automorphisms of } \mathcal{M}_\mu \end{array} \right.$$

2) Quantum Dynamical Systems (QDS $(\mathcal{M}_N, \omega_N, \Theta_N)$)

$$\left\{ \begin{array}{l} \mathcal{M}_N : \text{Finite dimensional algebra} \quad (N \times N \text{ Full-matrix algebra}) \\ \omega_N : \text{State over } \mathcal{M}_N \quad (\Theta_N\text{-invariant}) \quad (\omega_N(m) := \frac{1}{N} \text{Tr}(m)) \\ \Theta_N : \text{Unitary dynamics on } \mathcal{M}_N \quad (\Theta_N(m) := U m U^\dagger) \end{array} \right.$$

QUANTIZATION

To quantize a CDS $(\mathcal{M}_\mu, \omega_\mu, \Theta)$ means to find two linear maps $\mathcal{J}_{N,\infty}$ and $\mathcal{J}_{\infty,N}$ such that:

$$\begin{array}{ll} \mathcal{J}_{N,\infty} : \mathcal{M}_\mu \longmapsto \mathcal{M}_N & ; \quad \mathcal{J}_{N,\infty}(f) = M_N \\ \mathcal{J}_{\infty,N} : \mathcal{M}_N \longmapsto \mathcal{M}_\mu & ; \quad \mathcal{J}_{\infty,N}(M_N) = f \end{array}$$

$$\text{Classical limit is: } \mathcal{J}_{\infty,N} \circ \mathcal{J}_{N,\infty} \xrightarrow{N \rightarrow \infty} \mathbb{1}_{\mathcal{M}_\mu}$$

WEYL GROUP

- On compact phase space, we cannot make a finite dimensional quantization with CCR

$$[\hat{Q}, \hat{P}] = i \hbar \mathbb{1} ;$$

- We can find U_N and V_N that behave as $e^{2\pi i \hat{P}}$, respectively $e^{-2\pi i \hat{Q}}$

$$N = \frac{1}{\hbar}$$

WEYL OPERATORS

$W_N(\mathbf{n}) = e^{2\pi i(n_1 \hat{P} - n_2 \hat{Q})} = e^{\frac{i\pi}{N} n_1 n_2} V_N^{n_2} U_N^{n_1}$ provide the so-called

WEYL QUANTIZATION:

$$f(\mathbf{x}) = \sum_{\mathbf{n} \in \mathbb{Z}^2} \hat{f}_{\mathbf{n}} e^{2\pi i \sigma(\mathbf{n}, \mathbf{x})} \quad \sigma(\mathbf{n}, \mathbf{x}) = n_1 x_2 - n_2 x_1$$

can be mapped in $M_f = \sum_{\mathbf{n} \in \mathbb{Z}^2} \hat{f}_{\mathbf{n}} W_N(\mathbf{n})$

by means of the Weyl quantization operator

$$\mathcal{J}_{N,\infty}^W : \mathcal{M}_\mu \longmapsto \mathcal{M}_N \quad ; \quad \mathcal{J}_{N,\infty}^W(f) = M_f$$

DYNAMICAL EVOLUTION OF THE WEYL OPERATORS:

$$\Theta_N^j(W_N(\mathbf{n})) = W_N(T^j \cdot \mathbf{n}) \quad .$$

Such a relation guarantees:

$$\left(\Theta_N^j \circ \mathcal{J}_{N,\infty}^W \right) (f) = \left(\mathcal{J}_{N,\infty}^W \circ \Theta^j \right) (f) \quad .$$

CLASSICAL LIMIT FOR THE DYNAMICS

$$\begin{array}{ccc} \text{Quantum} & \xrightarrow[\text{LIMIT}]{\text{CLASSICAL}} & \text{Classical} \\ \text{evolution} & & \text{evolution} \end{array}$$

But this holds **ONLY IF** time–evolution does not cross a **BREAKING TIME**, depending on N !

In particular, depending on the Dynamical System considered, it exists an α such that for any given $f \in L_\mu^\infty(\mathbb{T}^2)$ it holds true

$$\lim_{\substack{k, N \rightarrow \infty \\ k < \alpha \log N}} \left\| \left(\Theta^k - \mathcal{J}_{\infty, N} \circ \Theta_N^k \circ \mathcal{J}_{N, \infty} \right) (f) \right\|_2 = 0$$

where $\|\cdot\|_2$ is the $L_\mu^2(\mathbb{T}^2)$ norm $\|g\|_2 := \sqrt{\int_{\mathbb{T}^2} |g|^2 \mu(d\mathbf{x})}$

ANTI–WICK QUANTIZATION

Using a “well defined” set of Coherent States (CS):

$$\begin{aligned} \mathcal{J}_{N\infty}(f) &:= N \int_{\mathcal{X}} \mu(d\mathbf{x}) f(\mathbf{x}) |C_N(\mathbf{x})\rangle \langle C_N(\mathbf{x})| \\ \mathcal{J}_{\infty N}(X)(\mathbf{x}) &:= \langle C_N(\mathbf{x}), X C_N(\mathbf{x}) \rangle \end{aligned}$$

Our CS family $\{|C_N(\mathbf{x})\rangle \mid \mathbf{x} \in \mathcal{X}\}$

$$\begin{aligned} |C_N(\mathbf{x})\rangle &:= W_N(\lfloor N\mathbf{x} \rfloor) |C_N\rangle \\ |C_N\rangle &= \sum_{j=0}^{N-1} C_N(j) |j\rangle \quad C_N(j) := \frac{1}{2^{(N-1)/2}} \sqrt{\binom{N-1}{j}}. \end{aligned}$$

Properties of $\{|C_N(\mathbf{x})\rangle \mid \mathbf{x} \in \mathcal{X}\}$

1. **Measurability**: $\mathbf{x} \mapsto |C_N(\mathbf{x})\rangle$ is measurable on \mathcal{X} ;
2. **Normalization**: $\|C_N(\mathbf{x})\|^2 = 1, \mathbf{x} \in \mathcal{X}$;
3. **Completeness**: $N \int_{\mathcal{X}} \mu(d\mathbf{x}) |C_N(\mathbf{x})\rangle \langle C_N(\mathbf{x})| = \mathbb{1}$;
- 4'. **Localization**: given $\varepsilon > 0$ and $d_0 > 0$, there exists $N_0(\varepsilon, d_0)$ such that for $N \geq N_0$ and $d(\mathbf{x}, \mathbf{y}) \geq d_0$ one has

$$N |\langle C_N(\mathbf{x}), C_N(\mathbf{y}) \rangle|^2 \leq \varepsilon.$$

- 4''. **Dynamical localization**:

There exists an $\alpha > 0$ such that for all choices of $\varepsilon > 0$ and $d_0 > 0$ there exists an $N_0 \in \mathbb{N}$ with the following property: if $N > N_0$ and $k \leq \alpha \log N$, then $N |\langle C_N(\mathbf{x}), U_N^k C_N(\mathbf{y}) \rangle|^2 \leq \varepsilon$ whenever $d(T^k \mathbf{x}, \mathbf{y}) \geq d_0$.

(U_N^k is the single step unitary evolution operator).

RESULTS

PROPERTIES: 1 2 3 4' \implies Classical limit

1 2 3 4' 4'' \implies Classical limit
of the dynamic

The Alicki Lindblad Fannes Dynamical Entropy

Given a Quantum Dynamical System $(\mathcal{M}_N, \omega_N, \Theta_N)$ we introduce:

- $\mathcal{Y} := \{y_\ell\}_{\ell=1}^D$; $\sum_{\ell=1}^D y_\ell^\dagger y_\ell = \mathbb{1}_{\mathcal{M}_0}$ – PARTITION OF UNIT
 $y_\ell \in \mathcal{M}_0 \subseteq \mathcal{M}_N$; \mathcal{M}_0 (subalgebra) s.t. $\Theta_N(\mathcal{M}_0) = \mathcal{M}_0$
- **EXAMPLE:** Partition of 2 elements ($D = 2$)

$\mathcal{M}_N ::$ $N \times N$ Full-matrix algebra

$\omega_N ::$ $\omega_N(m) := \frac{1}{N} \text{Tr}(m)$

$\Theta_N ::$ $\Theta_N(m) := U m U^\dagger$

$\mathcal{Y} ::$ $\{M_0, M_1\}$ (fulfilling $M_0^\dagger M_0 + M_1^\dagger M_1 = \mathbb{1}_N$)

- the time-evolving partition of unit: $\Theta_N^k(\mathcal{Y}) := \{\Theta_N^k(y_i)\}_{i=1}^D$

- **EXAMPLE:** with the partition $\{M_0, M_1\}$

$\Theta_N^k(\mathcal{Y}) ::$ $\{\Theta_N^k(M_0), \Theta_N^k(M_1)\} = \{U^k M_0 U^{\dagger k}, U^k M_1 U^{\dagger k}\}$

- the refined partition:

$\mathcal{Y}_{\Theta_N}^{[0, n-1]} = \left\{ \Theta_N^{n-1}(y_{i_{n-1}}) \Theta_N^{n-2}(y_{i_{n-2}}) \cdots \Theta_N(y_{i_1}) y_{i_0} \right\}_{i \in \Omega_D^n}$

- **EXAMPLE:** with the partition $\{M_0, M_1\}$ and $n = 3$

$$\mathcal{Y}_{\Theta_N}^{[0,2]} :: \left\{ M_{(000)}, M_{(001)}, M_{(010)}, M_{(011)}, \right. \\ \left. M_{(100)}, M_{(101)}, M_{(110)}, M_{(111)} \right\}$$

$$\left\{ \begin{array}{l} M_{(000)} := \Theta_N^2(M_0) \Theta_N(M_0) \quad M_0 = U^2 M_0 U^\dagger \quad U M_0 U^\dagger \quad M_0 = U^2 M_0 U^\dagger M_0 U^\dagger M_0 \\ M_{(001)} := \Theta_N^2(M_1) \Theta_N(M_0) \quad M_0 = U^2 M_1 U^\dagger \quad U M_0 U^\dagger \quad M_0 = U^2 M_1 U^\dagger M_0 U^\dagger M_0 \\ M_{(010)} := \Theta_N^2(M_0) \Theta_N(M_1) \quad M_0 = U^2 M_0 U^\dagger \quad U M_1 U^\dagger \quad M_0 = U^2 M_0 U^\dagger M_1 U^\dagger M_0 \\ M_{(011)} := \Theta_N^2(M_1) \Theta_N(M_1) \quad M_0 = U^2 M_1 U^\dagger \quad U M_1 U^\dagger \quad M_0 = U^2 M_1 U^\dagger M_1 U^\dagger M_0 \\ M_{(100)} := \Theta_N^2(M_0) \Theta_N(M_0) \quad M_1 = U^2 M_0 U^\dagger \quad U M_0 U^\dagger \quad M_1 = U^2 M_0 U^\dagger M_0 U^\dagger M_1 \\ M_{(101)} := \Theta_N^2(M_1) \Theta_N(M_0) \quad M_1 = U^2 M_1 U^\dagger \quad U M_0 U^\dagger \quad M_1 = U^2 M_1 U^\dagger M_0 U^\dagger M_1 \\ M_{(110)} := \Theta_N^2(M_0) \Theta_N(M_1) \quad M_1 = U^2 M_0 U^\dagger \quad U M_1 U^\dagger \quad M_1 = U^2 M_0 U^\dagger M_1 U^\dagger M_1 \\ M_{(111)} := \Theta_N^2(M_1) \Theta_N(M_1) \quad M_1 = U^2 M_1 U^\dagger \quad U M_1 U^\dagger \quad M_1 = U^2 M_1 U^\dagger M_1 U^\dagger M_1 \end{array} \right.$$

- the $D^n \times D^n$ density matrices $\rho[\mathcal{Y}_{\Theta_N}^{[0,n-1]}]$ with elements

$$\left[\rho[\mathcal{Y}_{\Theta_N}^{[0,n-1]}] \right]_{i,j} := \omega_N \left(\left[\mathcal{Y}_{\Theta_N}^{[0,n-1]} \right]_j^\dagger \left[\mathcal{Y}_{\Theta_N}^{[0,n-1]} \right]_i \right).$$

- **EXAMPLE:** with the partition $\mathcal{Y}_{\Theta_N}^{[0,2]}$

$$\left[\rho[\mathcal{Y}_{\Theta_N}^{[0,2]}] \right]_{i,j} := \omega_N \left(M_{(j_0, j_1, j_2)}^\dagger M_{(i_0, i_1, i_2)} \right) \quad \text{and so} \\ \left[\rho[\mathcal{Y}_{\Theta_N}^{[0,2]}] \right]_{(010), (100)} := \omega_N \left(M_{(100)}^\dagger M_{(010)} \right) \\ := \frac{1}{N} \text{Tr} \left(M_1^\dagger U M_0^\dagger U M_0^\dagger M_0 U^\dagger M_1 U^\dagger M_0 \right)$$

- the Von Neumann Entropy:

$$H_{\omega_N, \mathcal{M}_0}[\mathcal{Y}_{\Theta_N}^{[0,n-1]}] = - \text{Tr} \left(\rho[\mathcal{Y}_{\Theta_N}^{[0,n-1]}] \log \rho[\mathcal{Y}_{\Theta_N}^{[0,n-1]}] \right).$$

Then, the **ALF**–entropy of $(\mathcal{M}_N, \omega_N, \Theta_N)$ is given by:

$$h_{\omega_N, \mathcal{M}_0}^{\text{ALF}}(\Theta_N) := \sup_{\mathcal{Y} \subset \mathcal{M}_0} h_{\omega_N, \mathcal{M}_0}^{\text{ALF}}(\Theta_N, \mathcal{Y}) ,$$

where
$$h_{\omega_N, \mathcal{M}_0}^{\text{ALF}}(\Theta_N, \mathcal{Y}) := \limsup_n \frac{1}{n} H_{\omega_N} \left[\mathcal{Y}^{[0, n-1]} \right] .$$

PROPOSITION 1

Let $(\mathcal{A}_X, \omega_\mu, \Theta)$ represent a classical dynamical system. Then,

$$h_{\omega_\mu, \mathcal{A}_X}^{\text{ALF}}(\Theta) = h_\mu(T)$$

PROPOSITION 2

If $(\mathcal{M}_N, \omega_N, \Theta_N)$ be a quantum dynamical system with \mathcal{M}_N finite dimensional, then

$$h_{\omega, \mathcal{M}_N}^{\text{ALF}}(\Theta_N) = 0$$

h^{ALF}

- behave as $h_\mu(T)$ for CDS
- (so) - test CHAOS as $h_\mu(T)$ do
- reveal **NO CHAOS** on finite dimensional systems

- All these quantities are computed in the $n \longrightarrow \infty$ limit...
- What about the running Von Neumann entropies?

RESULT (For the ALF entropy)

Finding a correspondence (actually the more natural) between

- a classical partition \mathcal{E} KS
- a partition of unit \mathcal{Y} ALF






and using the dynamical localization condition, we get

$$\lim_{\substack{k, N \rightarrow \infty \\ k \leq \alpha \log N}} \frac{1}{k} \left| H_{\omega_N}[\mathcal{Y}_{\Theta_N}^{[0, k-1]}] - S_{\mu}(\mathcal{E}^{[0, k-1]}) \right| = 0.$$


– Remember –

$$\begin{array}{ccccc}
 S_{\mu}(\mathcal{E}^{[0, n-1]}) & \xrightarrow[\textcolor{red}{n}]{\textcolor{red}{1} \lim_{n \rightarrow \infty}} & h_{\mu}(T, \mathcal{E}) & \xrightarrow[\textcolor{red}{\mathcal{E}}]{\textcolor{red}{\sup}} & h_{\mu}(T) \\
 \uparrow \lim_{\substack{k, N \rightarrow \infty \\ k \leq \alpha \log N}} \frac{1}{k} \cdots & & & & \\
 H_{\omega_N, \mathcal{M}_0}[\mathcal{Y}_{\Theta_N}^{[0, n-1]}] & \xrightarrow[\textcolor{red}{n}]{\textcolor{red}{1} \limsup_{n \rightarrow \infty}} & h_{\omega_N, \mathcal{M}_0}^{\text{ALF}}(\Theta_N, \mathcal{Y}) & \xrightarrow[\textcolor{red}{\mathcal{Y}}]{\textcolor{red}{\sup}} & h_{\omega_N, \mathcal{M}_0}^{\text{ALF}}(\Theta_N)
 \end{array}$$

\mathcal{Y} : partition of D elements that maximize the entropy rate

Time n	1	2	3	...	m	m'
$\text{Size}(\rho_{\Theta}^{[0,n-1]})$	$D \times D$	$D^2 \times D^2$	$D^3 \times D^3$	\dots	$D^m \times D^m$	$D^{m'} \times D^{m'}$
Max. number of eigenvalues of $\rho_{\Theta}^{[0,n-1]}$ different from 0	D 	D^2 	D^3 	\dots	D^m 	N^2 
$\langle \text{Eigenvalues} \rangle$	$\frac{1}{D}$	$\frac{1}{D^2}$	$\frac{1}{D^3}$	\dots	$\frac{1}{D^m}$	$\frac{1}{N^2}$
$H_{\tau_N, \mathcal{M}_0} [\mathcal{Y}_{\Theta}^{[0,n-1]}]$	$\log D$	$2 \log D$	$3 \log D$	\dots	$m \log D$	$2 \log N$
$h_{\tau_N, \mathcal{M}_0}^{\text{ALF}} [\mathcal{Y}, \Theta]$	$\log D$	$\log D$	$\log D$	\dots	$\log D$	$\frac{2 \log N}{m'}$

BREAKING TIME



Given a Quantum Dynamical System $(\mathcal{M}_N, \omega_N, \Theta_N)$ we introduce:

$$\mathcal{Y} := \{y_\ell\}_{\ell=1}^D ; \quad \sum_{\ell=1}^D y_\ell^* y_\ell = \mathbb{1}_{\mathcal{M}_0} \quad - \quad \text{PARTITION OF UNIT (PU)}$$

$$\text{WAVE PACKET REDUCTION POSTULATE} \implies \rho \xrightarrow{\text{MEASURE}} \mathcal{I}_{\mathcal{Y}}(\rho) := \sum_j y_j \rho y_j^*$$

- The map $\mathcal{I}_{\mathcal{Y}}$ is called an **instrument**;
- it describe the change in the state ρ caused by the measure;
- $\omega[y_j \rho y_j^*]$ is the probability that the measure select the i^{th} value.

The CS Instrument

$$\mathcal{E} := \{E_\ell\}_{\ell=1}^D ; \quad \begin{cases} \bigcup_{\ell=1}^D E_\ell = \mathcal{X} \\ E_\ell \cap E_k = \emptyset \end{cases} \quad - \quad \text{CLASSICAL PARTITION (CP)}$$

With the family of CS $\{|C_N(\mathbf{x})\rangle \mid \mathbf{x} \in \mathcal{X}\}$ and $P_{\mathbf{x}} := |C_N(\mathbf{x})\rangle\langle C_N(\mathbf{x})|$:

- the map $\mathcal{I}(E_\ell)(\rho) := \mathcal{N} \int_{E_\ell} P_{\mathbf{x}} \rho P_{\mathbf{x}} \mu(d\mathbf{x})$ is called a **CS-instrument**;
- it describe the change in the state ρ caused by the E_ℓ -dependent measurement process;
- $\omega[\mathcal{I}(E_\ell)(\rho)]$ is the probability that the measure gives values in E_ℓ , when the pre-measurement state is ρ .

Time-stroboscopic CS measurement

$$\mathcal{P}_i^{\text{CS}} = \mathcal{P}_{i_0, i_1, \dots, i_{n-1}}^{\text{CS}} := \omega[\mathcal{I}(E_{i_{n-1}}) \circ \Theta \circ \mathcal{I}(E_{i_{n-2}}) \circ \Theta \circ \dots \circ \mathcal{I}(E_{i_1}) \circ \Theta \circ \mathcal{I}(E_{i_0})(\rho)]$$

is the probability that several measure, taken stroboscopically at times $t_0 = 0$, $t_1 = 1$, ..., $t_{n-1} = n-1$, give values in $E_{i_0}, E_{i_1}, \dots, E_{i_{n-1}}$.

CS Quantum Entropies

With the probabilities μ_i we can compute the SHANNON ENTROPY

$$S(U, \mathcal{I}, \mathcal{E}, \rho, n) := - \sum_{i \in \Omega_D^n} \mathcal{P}_i^{\text{CS}} \log \mathcal{P}_i^{\text{CS}} \quad ;$$

its production per time step, is defined as CS quantum entropy

$$H(U, \mathcal{I}, \mathcal{E}, \rho) := \lim_{n \rightarrow \infty} \frac{1}{n} S(U, \mathcal{I}, \mathcal{E}, \rho, n)$$

and it is decomposable in two part: the Measurement CS Quantum Entropy

$$H_{\text{meas}}(\mathcal{I}, \mathcal{E}, \rho) := H(1_{\mathcal{N}}, \mathcal{I}, \mathcal{E}, \rho) \quad ,$$

and the remaining part, which is supposed to incorporate the dynamics

$$H_{\text{dyn}}(U, \mathcal{I}, \mathcal{E}, \rho) := H(U, \mathcal{I}, \mathcal{E}, \rho) - H_{\text{meas}}(\mathcal{I}, \mathcal{E}, \rho) \quad .$$

PROPOSITION 1

Consider the Classical Dynamical System (\mathcal{X}, μ, T) endowed with a classical partition \mathcal{E} . Then it is possible to define the automorphism U and the classical instrument \mathcal{I} in such a way that

$$H(U, \mathcal{I}, \mathcal{E}, \rho) = h_{\mu}(T, \mathcal{E})$$

holds true.

PROPOSITION 2

For finite dimensional systems

$$H(U, \mathcal{I}, \mathcal{E}, \rho) = 0$$

RESULT (For the CS Quantum Entropy)

If we assume that dynamical localization condition holds, and we take for ρ the tracial state $\frac{1}{N} \mathbf{1}_N$, we find an α such that it holds true

$$\lim_{\substack{n, N \rightarrow \infty \\ n < \alpha \log N}} \frac{1}{n} \left| S(U, \mathcal{I}, \mathcal{E}, \rho, n) - S_\mu(\mathcal{E}_{[0, n-1]}) \right| = 0 \quad .$$

Moreover, this effect is purely related to the dynamic component of the entropy, indeed it exists an α' such that

$$\lim_{\substack{n, N \rightarrow \infty \\ n < \alpha' \log N}} \frac{1}{n} S(\mathbf{1}_N, \mathcal{I}, \mathcal{E}, \rho, n) = 0 \quad .$$

– Remember –

$$\begin{array}{ccccc}
 S_\mu(\mathcal{E}^{[0, n-1]}) & \xrightarrow[\textcolor{red}{n}]{\frac{1}{n} \lim_{n \rightarrow \infty}} & h_\mu(T, \mathcal{E}) & \xrightarrow[\textcolor{red}{\mathcal{E}}]{\sup} & h_\mu(T) \\
 \uparrow \lim_{\substack{k, N \rightarrow \infty \\ k \leq \alpha \log N}} \frac{1}{k} \cdots & & & & \\
 S(U, \mathcal{I}, \mathcal{E}, \rho, n) & \xrightarrow[\textcolor{red}{n}]{\frac{1}{n} \limsup_{n \rightarrow \infty}} & H(U, \mathcal{I}, \mathcal{E}, \rho) & \longrightarrow & \begin{cases} H_{\text{meas}}(\mathcal{I}, \mathcal{E}, \rho) \\ H_{\text{dyn}}(U, \mathcal{I}, \mathcal{E}, \rho) \end{cases}
 \end{array}$$

CONCLUSION

- We used QDE to find footprint of CHAOS in quantum (or discrete) systems, obtained from classical continuous one.
 - We found that the correspondence between Classical and Quantum Dynamics (Breaking Time BT) lasts much less than the Heisenberg time.
 - The BT scales logarithmically in the dimension of the Hilbert space, moreover it is inversely proportional to the Lyapunov exponent.
 - For the Quantum Cat Maps we exactly determined $BT = \frac{1}{2} \frac{\log N}{\log \lambda}$
-
- We showed how Quantum Dynamical Entropies can be profitably used in a Classical Discretized context.